

Department for Environment, Food and Rural Affairs

The Expert Committee on Pesticide Residues in Food (PRiF)

Report on the pesticide residues monitoring programme: Results of Quarter 1 2021



OGL

© Crown copyright 2021

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit [The National Archives](#) or email PSI@nationalarchives.gov.uk

This publication is available at [Expert Committee on Pesticide Residues in Food](#)

For further details on information contained in this report, previous surveys or information concerning pesticide residues in food, please contact:

Expert Committee on Pesticide Residues in Food
HSE's Chemicals Regulation Division
Mallard House
Kings Pool
Peasholme Green
York
YO1 7PX

prif@hse.gov.uk

Contents

Summary: Quarter 1 2021 Findings	4
Chair's summary of results	4
Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF)	9
2021 Survey Design	12
Summary of Table of Results	14
Summary of samples sent to FSA for follow up	17
Summary of MRL Exceedances	19
Section 1: findings by food	21
Aubergine (GB)	21
Aubergine (NI)	23
Banana (GB)	24
Banana (NI)	26
Beans with pods (GB)	27
Beans with pods (NI)	31
Beef (GB)	33
Beef (NI)	34
Berries and small fruit (GB)	35
Berries and small fruit (NI)	37
Broccoli (GB)	39
Broccoli (NI)	42
Cheese (soft) (GB)	43
Cheese (soft) (NI)	45
Eggs (GB)	47
Eggs (NI)	48
Fish (white) (NI)	49
Grapes (GB)	51
Grapes (NI)	53
Melon (GB)	54
Melon (NI)	56
Milk (GB)	58
Milk (NI)	59
Mushrooms (GB)	60
Mushrooms (NI)	62
Peppers (GB)	63
Peppers (NI)	65
Potatoes (GB)	67
Potatoes (NI)	68
Raspberry (GB)	69
Raspberry (NI)	71
Rice (GB)	73
Rice (NI)	75
Section 2: Sample details and supplier responses	77
Sample details	77
Supplier responses	79
Section 3: HSE assessment of risk	80
Assessment of Risk to Human Health	89
Section 4: issues arising in this report and updates on previous reports	112
Issues arising in this report	112
In our next report	116
Section 5: background and reference	117
Glossary	125

Summary: Quarter 1 2021 findings

Chair's comments on results

This is our Quarter 1 report for 2021.

During this year's surveillance programme, we are measuring up to 395 different pesticides in each of the foods we survey. The Quarter 1 programmes surveyed 531 samples of 17 different foods (see [contents page](#) for a full list).

Of 531 samples, we found residues in 220 of them and, of these, 15 samples contained residues over the Maximum Residue Level (MRL). Most of the residues detected did not cause health concern.

HSE undertakes screening and detailed risk assessments, as required, for the pesticide residues found. This is to determine whether the residues present could lead to someone eating an amount above a level that is considered safe. HSE also produces [detailed risk assessments](#) for every case where the actual residue level found could lead to an intake above the safety levels.

We needed to consider the potential short term health effects of some of the residues found in more detail. In these cases, we concluded that effects on health were either unlikely or not expected, or, in the case of chlorpyrifos in one sample of tenderstem broccoli, not clear if the chlorpyrifos level found may cause any adverse effect; noting too that the HSE's assessment of the risk is precautionary.

We also needed to consider the potential genotoxic health effects of some of the residues found. We concluded that at the levels present, a risk of an adverse effect on health due to genotoxicity would be low.

These detailed considerations as well as links to underlying information are covered in our reports for aubergine, banana, beans with pods, broccoli, grapes, and rice.

None of the individual commodity long term exposure screening assessments performed in this quarter (for each of the pesticides found in this report) indicated any potential for adverse long term health effects based on the assessment of dietary intakes being below the ADI or other established long term health based reference values.

Full details of suppliers and retailers of the food sampled, and full analytical results, are available on [data.gov.uk](#) as ODS (Open Document Spreadsheet) files. We hope this data format is useful for people wanting to look at the individual results in more detail.

These samples were collected after the UK had left the EU, so this is the first report where we have reported the results for samples collected in Great Britain (GB) separately from those collected in Northern Ireland (NI). Samples collected in GB are subject to GB MRLs. GB MRLs are set by inclusion in a new statutory Register, implemented and updated by means of a database¹. For samples collected in NI, certain aspects of EU food law, including compliance with EU set MRLs, continue to apply under the terms of the Northern

¹ [GB MRL Register \(pesticides.gov.uk\)](#)

Ireland Protocol. In the detailed data files HSE are, for 2021 results, still separating out EU from non-EU origin foods in the results.

We asked suppliers and the authorities of the exporting countries for an explanation of our findings. Any responses we have received specifically for publication are available in [Section 2](#) sample details and supplier responses.

If you have any feedback or comments on the monitoring programme or the reports produced please send them to our secretariat at prif@hse.gov.uk.

Ann Davison
Chair of the Expert Committee on Pesticide Residues in Food

Consumer risk summary

1 sample each of aubergine, guar beans and broccoli resulted in information being sent to the Food Standards Agency (FSA) for further follow up.

HSE screen each residue detected for any consumer health issues to identify which need to be considered in more detail. We comment on any risks HSE considered in detail in our full report, and [HSE' s risk assessments](#) are also published.

HSE liaise with the Food Standards Agency (FSA) on consumer risk assessment and the FSA also take part in our meetings.

Headlines with links to detailed information

Table 1

Food	Number of pesticides sought	Samples tested	Detailed risk assessment presented?	MRL Non-Compliance*
All Foods tested				
Aubergine (GB)	374	18	Yes	1
Aubergine (NI)	374	4		0
Banana (GB)	392	22	Yes	0
Banana (NI)	392	4		0
Beans with pods (GB)	389	24	Yes	8
Beans with pods (NI)	389	3		0
Beef (GB)	115	18		0
Beef (NI)	38	18		0
Berries & small fruits (GB)	369	12		1
Berries & small fruits (NI)	369	6		0
Broccoli (GB)	371	19	Yes	1
Broccoli (NI)	371	6		0
Cheese (soft) (GB)	113	24		0
Cheese (soft) (NI)	38	19		0
Eggs (GB)	115	24		0
Eggs (NI)	115	3		0
Fish (white) (NI)	38	19		0
Grapes (GB)	395	17	Yes	0
Grapes (NI)	395	4		0

Food	Number of pesticides sought	Samples tested	Detailed risk assessment presented?	MRL Non-Compliance*
Melon (GB)	373	25		0
Melon (NI)	373	6		0
Milk (GB)	112	72		0
Milk (NI)	112	7		0
Mushroom (GB)	395	36		1
Mushroom (NI)	395	3		0
Peppers (GB)	394	30		0
Peppers (NI)	394	3		0
Potatoes (GB)	390	20		0
Potatoes (NI)	390	3		0
Raspberry (GB)	369	24		0
Raspberry (NI)	369	7		0
Rice (GB)	386	24	Yes	2
Rice (NI)	386	7	Yes	0

* Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act 2020. For samples collected in NI, under the Northern Ireland Protocol, certain aspects of EU food law, including compliance with EU MRLs apply.

Other issues

Suspected unauthorised uses

HSE passed details to their enforcement team of samples from GB that contained a residue which does not have a plant protection product (PPP) with that active authorised for use on that crop in GB.

- Mushrooms (3 samples) with a residue of mepiquat. HSE's enforcement concluded the residues were likely a result of residual levels in the wheat straw used in mushroom substrate. This carryover was taken into account when the MRL was set.

Organic samples with residues

HSE writes to the suppliers of samples of organic produce if they contain a pesticide residue. Defra's Organic Farming branch and the organic certification organisation are also informed.

- None reported

Further information

These findings are discussed in more detail in our [Quarterly Report](#)

Further information in an accessible format at [Pesticide Residues in Food Quarterly Data](#).

Includes:

- brand name, sampling point and origin information
- pesticides sought and residues found
- HSE detailed risk assessments

PRiF

September 2021

Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF)

The PRiF's role is to give Ministers, the Director of the Health and Safety Executive (HSE) and the Chief Executive of the Food Standards Agency (FSA) independent advice on the UK government's national rolling programme of surveys, in particular:

- the planning of surveillance programmes for pesticide residues in the UK food supply and the evaluation of the results;
- procedures for sampling, sample processing, new methods of analysis, the assessment of variability of pesticide residues in food and related issues.

The Expert Committee on Pesticide Residues in Food was established in 2011. Our members have a broad range of expertise relating to the food supply industry. The main function of the Committee is to oversee Government's £2 million pesticide residues surveillance programme. Previously this work was carried out by the Pesticide Residues Committee.

Our Chair, Ann Davison has worked in consumer affairs for most of her career, running consumer organisations and networks. The committee also includes members with expertise in food science, food production and supply as well as two public interest experts.

Information on the membership of the PRiF is also available on the PRiF's website:

[Expert Committee on Pesticide Residues in Food](#)

UK National Monitoring Programmes

HSE, working under Defra, and the Scottish and Welsh governments authority, has official responsibility to organise a monitoring programme of GB food for pesticide residues. Similarly, HSE working under the Department of Agriculture, Environment and Rural affairs authority has official responsibility to organise a monitoring programme of NI food for pesticide residues, including participating in the EU multi-annual control programme.

The programmes are made up of a risk-based rolling programme of surveys and statutory programmes required by GB or EU law. It is a surveillance programme, which is designed based upon evidence gathered in the previous year, including previous results, PRiF advice and border control information. It is not an enforcement programme and its design is generally not adjusted during the year. HSE is also responsible for considering the safety of people who eat the food (in co-operation with the Food Standards Agency if necessary) and following up adverse or unexpected results. They are also responsible for determining whether food is compliant with the law, specifically whether any pesticide residue found is within the Maximum Residue Level.

Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in food which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not authorised for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory

analysis. This provides a mechanism for statutory controls on pesticides in food which is put into circulation and for monitoring the correct use of these chemicals.

HSE assessment of risk

HSE conducts a screening assessment of all the residues we find in the PRiF programme. If screening identifies any dietary intakes exceeding the relevant health based reference values, then we present more detailed risk assessments, to consider whether there are any implications for health. Detailed risk assessments, where needed, are presented in [Section 3](#). If we understand that a pesticide residue has a risk of genotoxicity (has potential to cause damage to genetic material), we will include this in the commentary.

Pesticide dietary intakes are assessed using models that combine data on the levels of residues in food with food dietary consumption values. If intakes are within the health based reference values, then taking account of the precautions built into the model assessments we conclude that an effect on health is not anticipated. If dietary intakes exceed the reference values this does not automatically mean there are expected adverse health effects. However, this acts as a 'trigger' for HSE to consider these cases more thoroughly.

HSE conducts both short term (acute) and long term (chronic) assessments based on the residues found in the PRiF. Each of these is tailored accordingly. Further information on the nature of HSE's assessments and approach is provided in the bullet points below, and in more detail, with reference to international assessment contexts in [section 3](#) and on HSE's website (<https://www.hse.gov.uk/pesticides/pesticides-registration/data-requirements-handbook/consumer-exposure.htm>)

- For acute assessment we use short term estimation values that use the highest residue found in a commodity and short term consumption values for calculating short term dietary intakes. These are then compared to the ARfD, a suitable health based reference value for effects that could be caused by a single day or one-off consumption of a higher than usual residue. For acute assessment we consider the variation in residues that could occur within a residue sample, and a variability (multiplication) factor is included for that purpose, in order to address exposure to a higher than usual residue in a single item, such as a single apple or potato.
- For chronic assessment we use long term estimation values (based on median residues and long term consumption values for calculating long term dietary intakes) for each commodity survey and compare to the ADI, a suitable health based reference value for life-time. The issue is more fully considered in regulatory contexts pre-authorisation and at the time of MRL review. Then the issue is considered across all commodities (so more precautionary) by pesticide levels determined in GAP compliant trials, intended to address highest likely residues that might arise following pesticide use according to label recommendations.
- For fruit and vegetables that have peel or skin that might not be consumed we present alternative risk assessments for 'without peel -flesh only' where peel versus pulp residue distribution data are available. As standard, we present a 'worst case' assessment for when all of the peel is consumed with the fruit.

- We calculate dietary intakes for different consumer groups, from infants, toddlers and children of varying age, to adults, elderly, and vegetarians, to take account of people with low bodyweights and varying dietary habits. As such the assessments we perform are protective for all consumers.
- For multiple residues, we consider the 'cocktail effect'- the possible implications to health of more than one pesticide being found in samples. We currently focus in detail on selected groups that we think are a priority to consider, based on toxicity considerations and prevalence.

Table 2: 2021 Survey Design

Fruit and vegetables

Food	Sampling points	Sampled during	Reporting
Asparagus (GB)	Retail Outlets and Supply chain	Quarters 2 and 4	Quarters 2 and 4
Asparagus (NI)	Retail Outlets	Quarters 2 and 4	Quarters 2 and 4
Aubergine (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Aubergine (NI)	Retail Outlets	Quarterly	Quarterly
Banana (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Banana (NI)	Retail Outlets	Quarterly	Quarterly
Beans with pods (GB)	Retail Outlets and Supply chain	Quarterly	Rolling and Quarterly
Beans with pods (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Berries and small fruit (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Berries and small fruit (NI)	Retail Outlets	Quarterly	Quarterly
Broccoli (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Broccoli (NI)	Retail Outlets	Quarterly	Quarterly
Grapefruit (GB)	Retail Outlets and Supply chain	Quarter 2 - 4	Quarter 2 - 4
Grapefruit (NI)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4
Grapes (GB)	Retail Outlets and Supply chain	Quarterly	Rolling and Quarterly
Grapes (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Melon (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Melon (NI)	Retail Outlets	Quarterly	Quarterly
Mushrooms (GB)	Retail Outlets	Quarterly	Quarterly
Mushrooms (NI)	Retail Outlets	Quarterly	Quarterly
Peppers (GB)	Retail Outlets and Supply chain	Quarterly	Quarterly
Peppers (NI)	Retail Outlets	Quarterly	Quarterly
Potatoes (GB)	Supply chain	Quarterly	Rolling and Quarterly
Potatoes (NI)	Retail Outlets	Quarterly	Rolling and Quarterly
Raspberry (GB)	Retail Outlets	Quarterly	Quarterly
Raspberry (NI)	Retail Outlets	Quarterly	Quarterly
Spring green and kale (GB)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4
Spring green and kale (NI)	Retail Outlets	Quarter 2 - 4	Quarter 2 - 4

Animal products

Food	Sampling points	Sampled during	Reporting
Beef (GB)	Retail Outlets	Quarterly	Quarterly
Beef (NI)	Retail Outlets	Quarterly	Quarterly
Cheese (soft) (GB)	Retail Outlets	Quarterly	Quarters 1, 2 and 4
Cheese (soft) (NI)	Retail Outlets	Quarterly	Quarterly
Eggs (GB)	Retail Outlets	Quarterly	Quarterly
Eggs (NI)	Retail Outlets	Quarterly	Quarterly
Fish (white) (NI)	Retail Outlets	Quarterly	Quarterly
Milk (GB)	Retail Outlets	Quarterly	Quarterly
Milk (NI)	Retail Outlets	Quarterly	Quarterly

Cereal products

Food	Sampling points	Sampled during	Reporting
Bread (GB)	Retail Outlets	Quarters 2 - 4	Quarter 3 and 4
Bread (NI)	Retail Outlets	Quarters 2 - 4	Quarter 3 and 4
Rice (GB)	Retail Outlets	Quarter 1 and 3	Quarter 1 and 3
Rice (NI)	Retail Outlets	Quarter 1 and 3	Quarter 1 and 3
Wheat (GB)	Retail Outlets	Quarterly	Quarter 2 and 4
Wheat (NI)	Retail Outlets	Quarterly	Quarter 2 and 4

Miscellaneous products

Food	Sampling points	Sampled during	Reporting
Edible seeds (GB)	Retail Outlets	Quarter 2 and 3	Quarter 3
Infant food (cereal based) (GB)	Retail Outlets	Quarter 3	Quarter 3
Infant food (cereal based) (NI)	Retail Outlets	Quarter 3	Quarter 3
Nuts (GB)	Retail Outlets	Quarters 3 and 4	Quarter 4
Olive Oil (GB)	Retail Outlets	Quarterly	Quarters 2 - 4
Olive oil (NI)	Retail Outlets	Quarterly	Quarters 2 - 4
Processed peppers (GB)	Retail Outlets	Quarter 3	Quarter 4
Soya milk (GB)	Retail Outlets	Quarter 3	Quarter 3
Soya products (GB)	Retail Outlets	Quarter 3	Quarter 3

Sampling points

- Retail outlets: samples bought by market research contractor shoppers.
- Supply Chain: samples taken by inspectors from the Animal and Plant Health Agency from a range of points in the supply chain (wholesalers, retail depots, ports and import points).

Reporting

- Results for certain higher-priority foods are produced, followed up and published more frequently at [Data.gov.uk](https://data.gov.uk)
- All results are published in the quarterly report. Some surveys are included in every quarter, some are every other quarter and some in just one quarter.

Table 3: Summary of Results

Food Type	Analysed	With residues at or below the MRL ²	With residues above the MRL ³	With residues of unauthorised pesticides	With multiple residues	Organic samples tested	Organic samples with residues
Aubergine (GB)	18	12	1	0	9	1	0
Aubergine (NI)	4	4	0	0	3	0	0
Banana (GB)	22	15	0	0	14	7	0
Banana (NI)	4	3	0	0	3	1	0
Beans with Pods (GB)	24	11	8	0	12	0	0
Beans with pods (NI)	3	1	0	0	1	0	0
Beef (GB)	18	0	0	0	0	2	0
Beef (NI)	18	1	0	0	0	0	0
Berries and small fruits (GB)	12	6	1	0	5	2	0
Berries and small fruits (NI)	6	6	0	0	5	0	0

². In analytical terms this is a reportable value between LOD and the MRL

³ Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act 2020. For samples collected in NI, under the Northern Ireland Protocol, certain aspects of EU food law, including compliance with EU MRLs apply.

Food Type	Analysed	With residues at or below the MRL ²	With residues above the MRL ³	With residues of unauthorised pesticides	With multiple residues	Organic samples tested	Organic samples with residues
Broccoli (GB)	19	12	1	0	8	1	0
Broccoli (NI)	6	4	0	0	2	1	0
Cheese (soft) (GB)	24	5	0	0	0	0	0
Cheese (soft) (NI)	19	0	0	0	0	0	0
Eggs (GB)	24	0	0	0	0	7	0
Eggs (NI)	3	0	0	0	0	0	0
Fish (white) (NI)	19	4	0	0	1	0	0
Grapes (GB)	17	17	0	0	16	0	0
Grapes (NI)	4	3	0	0	3	1	0
Melon (GB)	25	21	0	0	14	0	0
Melon (NI)	6	5	0	0	3	0	0
Milk (GB)	72	0	0	0	0	22	0
Milk (NI)	7	0	0	0	0	0	0
Mushroom (GB)	36	15	1	3	9	16	0
Mushroom (NI)	3	2	0	0	1	0	0
Peppers (GB)	30	20	0	0	16	4	0

Food Type	Analysed	With residues at or below the MRL ²	With residues above the MRL ³	With residues of unauthorised pesticides	With multiple residues	Organic samples tested	Organic samples with residues
Peppers (NI)	3	2	0	0	1	0	0
Potatoes (GB)	20	13	0	0	3	0	0
Potatoes (NI)	3	1	0	0	0	0	0
Raspberry (GB)	24	18	0	0	17	1	0
Raspberry (NI)	7	5	1	0	6	0	0
Rice (GB)	24	11	2	0	10	5	0
Rice (NI)	7	3	0	0	2	0	0

Table 4: Summary of samples sent to FSA for follow up

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)
3505/2021	22/02/2021	Guar Beans	India	Quality Veg Suppliers Ltd	Units 14/16, St James's Market, Essex Street, Bradford BD4 7PN	None stated	Indo British Imports 0-1 1st Floor, Export Building, ADCM Veg Market, Vashi, Navi Mumbai 400 705, India	monocrotophos 0.005 (MRL = 0.01*) omethoate 0.1 (MRL = 0.01*) profenofos 0.02 (MRL = 0.01*) cypermethrin (sum) 0.03 (MRL = 0.7) dimethoate 0.04 (MRL = 0.01*)
0093/2021	08/03/2021	Aubergine	The Netherlands	Iceland Foods	Barnfield Retail Park, Swindon SN2 2DJ	Iceland	Iceland Foods Ltd, Second Avenue, Deeside Industrial Park, Deeside, Flintshire, CH5 2NW	chlorate 1.6 (MRL = 0.4)

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)
0050/2021	25/01/2021	Tenderstem Broccoli	Kenya	Sainsbury's	Nottingham Road, Mansfield NG18 1BW	by Sainsbury's	Sainsbury's Supermarkets Ltd 33 Holborn, London EC1N 2HT	dimethoate 0.04 (MRL = 0.02) chlorantraniliprole 0.02 (MRL = 1) chlorpyrifos 0.06 (MRL = 0.01*) spinosad (sum) 0.01 (MRL = 2) omethoate 0.01 (MRL = 0.01*) lambda-cyhalothrin 0.01 (MRL = 0.1) indoxacarb 0.01 (MRL = 0.3)

Table 5: Summary of MRL Exceedances

Sample ID	Food Type	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg) ⁺	MRL exceedance after allowing for measurement uncertainty
Aubergine (GB)						
0093/2021	Aubergine	The Netherlands	chlorate	1.6	0.4	Yes
Beans with pods (GB)						
3504/2021	Speciality Beans	Kenya	bupirimate	0.08	0.05*	No
3505/2021	Speciality Beans	India	dimethoate	0.04	0.01*	Yes
			omethoate	0.1	0.01*	Yes
			profenofos	0.02	0.01*	Yes
3810/2021	Speciality Beans	India	diafenthiuron	1.1	0.01	Yes
3829/2021	Speciality Beans	India	profenofos	0.05	0.01*	Yes
3830/2021	Speciality Beans	India	hexaconazole	0.02	0.01*	No
3832/2021	Speciality Beans	India	hexaconazole	0.03	0.01*	Yes
3869/2021	Speciality Beans	Pakistan	profenofos	0.7	0.01*	Yes
3872/2021	Speciality Beans	Pakistan	profenofos	0.03	0.01*	Yes

Berries and small fruits (GB)

3924/2021	Fresh: Blackberries	Mexico	thiamethoxam	0.02	0.01*	Yes
-----------	---------------------	--------	--------------	------	-------	-----

Broccoli (GB)

0050/2021	Fresh	Kenya	chlorpyrifos	0.06	0.01*	Yes
			dimethoate	0.04	0.02	No

Mushroom (GB)

1208/2021	Chestnut	Ireland	BAC (sum)	0.2	0.1	No
-----------	----------	---------	-----------	-----	-----	----

Raspberry (NI)

0019/2021	Frozen	UK	dithiocarbamates	0.08	0.05*	No
-----------	--------	----	------------------	------	-------	----

Rice (GB)

0144/2021	Basmati	UK	buprofezin	0.02	0.01*	No
			thiamethoxam	0.09	0.01*	Yes
			tricyclazole	0.1	0.01*	Yes
2347/2021	Basmati	UK	tricyclazole	0.02	0.01*	Yes

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. Either insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop permitted. However, they may be permitted elsewhere.

+ Samples collected in GB must comply with GB set MRLs unless the goods are qualifying Northern Ireland goods and are subject to unfettered access under the terms of the UK Internal Market Act 2020. For samples collected in NI, under the Northern Ireland Protocol, certain aspects of EU food law, including compliance with EU MRLs apply.

Section 1: findings by food

Aubergine (GB)

Samples tested

18 samples were tested for up to 374 pesticide residues

- 3 samples were imported from outside the EU
- 15 samples came from the EU

Pesticide residues detected from those sought

- 5 samples contained no residues from those sought
- 13 samples contained residues above the reporting limit
- 1 sample contained a residue above the MRL
- 1 sample was labelled as organic. It did not contain any residues from those sought

Risk assessments

A sample of aubergine contained a residue of chlorate at 1.6 mg/kg where the effect on health needed to be considered in more detail. Based on the HSE Chemicals Regulation Division's risk assessment (please see [Section 3](#)) of the residue detected we consider a short term effect on health to be unlikely.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to '[how HSE perform the risk assessments](#)' for further details.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We considered a residue of chlorate at 1.6 mg/kg, and above the MRL, in more detail. HSE's assessment of risk concluded that an effect on health would be unlikely. This finding is unusually high. We are testing aubergine for the whole year so we will be able to determine if it was an outlier.

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in aubergine. Details are available in [Table 5](#).

HSE have passed details of the following sample to FSA for further consideration. Further details are in [Table 4](#)

- One sample from The Netherlands containing chlorate at 1.6 mg/kg.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Aubergine (NI)

Samples tested

4 samples were tested for up to 374 pesticide residues

- 4 samples came from the EU

Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Banana (GB)

Samples tested

22 samples were tested for up to 392 pesticide residues

Banana (eating)

- 19 samples were imported from outside the EU

Plantain

- 3 samples were imported from outside the EU

Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- 15 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 7 samples were labelled as organic. None contained residues from those sought

Risk assessments

A sample of banana contained a residue of chlorpyrifos at 0.004 mg/kg where the effect on health needed to be considered in more detail.

Based on the HSE assessment of risk for this sample containing chlorpyrifos, if the bananas are consumed without the peel a short term effect on health is not expected.

HSE always undertake assessments that consider both when the peel is not eaten, as per the MRL assessment, and one where it is assumed that the peel is eaten. These assessments are detailed in [section 3](#) and should be consulted for the full assessment of risk.

Assuming bananas are eaten whole, including all of the peel, this sample containing a residue of chlorpyrifos at 0.004 mg/kg, we consider a short term effect on health to be unlikely. However, if the peel is not consumed then only 2% of the residue remains and based on this lower intake a short term effect on health is not expected.

As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on this residue of chlorpyrifos (0.004 mg/kg) in banana.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to '[how HSE perform the risk assessments](#)' for further details.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

The potential short term effect of one residue needed to be considered in more detail, (residue of chlorpyrifos at 0.004 mg/kg). Based on HSE's assessment of short term risk it was concluded that assuming bananas are eaten whole, including all of the peel, we consider an effect on health to be unlikely. However, if the peel is not consumed then only 2% of the residue remains and based on this lower intake an effect on health is not expected. We also concluded any risk of adverse health effects due to genotoxicity to be low.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Banana (NI)

Samples tested

4 samples were tested for up to 392 pesticide residues

Bananas (eating)

- 4 samples were imported from outside the EU

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 3 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Beans with pods (GB)

Samples tested

24 samples were tested for up to 389 pesticide residues

Dwarf Beans

- 4 samples were imported from outside the EU

Fine Beans

- 2 samples were imported from outside the EU

Green Beans

- 5 samples were imported from outside the EU

Runner Beans

- 1 sample was imported from outside the EU

Speciality Beans

- 12 samples were imported from outside the EU

Pesticide residues detected from those sought

- 5 samples contained no residues from those sought
- 19 samples contained residues above the reporting limit
- 8 samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

Following screening assessment, there were four pesticides where the effect on health needed to be considered in more detail. These are outlined in the bullet points below. For each of these, please see [section 3](#) for the full details of the HSE assessment of the risks.

- Residues of chlorpyrifos were found at up to 0.01 mg/kg in Guar beans. Based on the low short term intakes, HSE considers that a short term effect on health is not expected. As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the highest residue(s) found in this report.
- A sample of Guar beans contained a residue of diafenthiuron at 1.1 mg/kg. This is an old pesticide and modern reference values are not available. An ADI value set by Australia in 1993, was used as a surrogate reference value to assess the short term exposure. Although the ADI was exceeded, the assessment using the ADI is

considered to be precautionary for short term exposure. Based on HSE's assessment of the risk we conclude that an effect on health is unlikely.

- Dimethoate and omethoate are chemically related pesticides and for toxicology purposes are considered together. Omethoate is also the main metabolite of dimethoate. Residues of dimethoate and omethoate were found in beans with pods at levels up to 0.04 mg/kg and 0.1 mg/kg respectively. The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and omethoate, due to a lack of a fully supporting toxicological database. We think that at the anticipated highest exposures following consumption of the Guar beans sample containing the highest residues of dimethoate and omethoate, there is unlikely to be a risk of ill health effects based on short term toxicity. The EFSA Conclusion (2018) also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of dimethoate and omethoate are undesirable due to the uncertainty regarding genotoxicity at low doses. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the highest residue(s) found in this report.
- A sample of Guar beans contained a residue of monocrotophos at 0.005 mg/kg. Monocrotophos is an insecticide that has not been authorised for use in the EU since 2003. Although a low level residue, where a short term effect on health is not expected, there is uncertainty about the potential for monocrotophos to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of monocrotophos in food undesirable. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the level of 0.005 mg/kg in this beans with pods sample.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to ['how HSE perform the risk assessments'](#) for further details.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups, and may have similar toxicological effects. So the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined short term risk assessment of the relevant samples, as these contained different pesticides which might have a common effects; these pesticides are known to inhibit the enzyme acetylcholinesterase (please see the glossary on page 125). For these combined exposure assessments, we also refer to the above conclusions regarding genotoxicity (as these combinations also include pesticides which might be genotoxic).

- One sample contained omethoate and profenofos. HSE's combined risk assessment on this combination (see the detailed multiple exposure assessments in [section 3](#)) showed that the combined exposures are unlikely to inhibit acetylcholinesterase and a short term effect on health would not be expected.
- One sample contained omethoate and chlorpyrifos. HSE's combined risk assessment on this combination (see the detailed multiple exposure assessments in [section 3](#)) showed that the combined exposures are unlikely to inhibit acetylcholinesterase, and overall, a short term effect on health would not be expected.
- One sample contained dimethoate, omethoate, monocrotophos and profenofos. HSE's combined risk assessment on this combination (see the detailed multiple exposure assessments in [section 3](#)) showed that a short term effect on health would be unlikely. It is not anticipated that the monocrotophos and profenofos residues will contribute significantly to the overall combined intake when compared to the single substance assessment for dimethoate and its metabolite omethoate.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We needed to consider four residues in more detail and following HSE's assessment of risk, concluded the likelihood of adverse health effects to be low or unlikely.

There are a small number of examples of older pesticides (no longer approved in the UK) that might be genotoxic, where modern data to further investigate the genotoxic potential is not expected to be made available. It is likely that these will only be found in imported foods. For many of these old pesticides, the toxicological reference doses are low. We have decided to test for them in some surveys using lower reporting limits to ensure that these residues are found even at very low levels, as we know they are of particular interest to consumers.

We have surveyed beans with pods for a number of years and publish the results of this sampling throughout the year as part of our rolling reporting programme as we know that a high percentage of the samples are imported and may contain residues of these older pesticides.

Residues measured above the MRL

The laboratory detected 6 residues above the MRL in beans with pods. Details are available in [Table 5](#).

HSE have passed details of the following sample to FSA for further consideration. Further details are in [Table 4](#)

- 1 sample from India containing dimethoate at 0.04 mg/kg and omethoate at 0.1 mg/kg

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Beans with pods (NI)

Samples tested

3 samples were tested for up to 389 pesticide residues

Fine Beans

- 1 sample was imported from outside the EU

Green Beans

- 1 sample was imported from outside the EU

Runner Beans

- 1 sample was imported from outside the EU

Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 1 sample contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in the sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)

- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Beef (GB)

Samples tested

18 samples were tested for up to 115 pesticide residues

- 15 samples came from the UK
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the beef was produced. It may be where the beef was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 18 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Beef (NI)

Samples tested

18 samples were tested for up to 38 pesticide residues

- 15 samples came from the UK
- 3 samples came from the EU

The country of origin of samples may not be the same as the country where the beef was produced. It may be where the beef was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 17 samples contained no residues from those sought
- 1 sample contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

The residue detected by the laboratory would not be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

None of the samples contained more than one residues, so we did not carry out a combined risk assessment.

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Berries and small fruits (GB)

Samples tested

12 samples were tested for up to 369 pesticide residues

Fresh: Blackberries

- 2 samples were imported from outside the EU

Fresh: Blueberries

- 7 samples were imported from outside the EU
- 1 sample came from the EU

Frozen: Blackberries

- 1 sample came from the UK

Frozen: Blueberries

- 1 sample came from the EU

The country of origin of samples may not be the same as the country where the berries were produced. It may be where the berries were processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 5 samples contained no residues from those sought
- 7 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 2 samples were labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in blackberries. Details are available in [Table 5](#).

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Berries and small fruits (NI)

Samples tested

6 samples were tested for up to 369 pesticide residues

Fresh: Blueberries

- 4 samples were imported from outside the EU

Frozen: Blueberries

- 1 sample came from the UK
- 1 sample came from the EU

The country of origin of samples may not be the same as the country where the berries were produced. It may be where the berries were processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)

- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Broccoli (GB)

Samples tested

19 samples were tested for up to 371 pesticide residues

Fresh

- 2 samples were imported from outside the EU
- 15 samples came from the EU

Frozen

- 1 sample came from the UK
- 1 sample came from the EU

Pesticide residues detected from those sought

- 6 samples contained no residues from those sought
- 13 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

Risk assessments

Following screening, one tenderstem broccoli sample from Kenya contained residues of chlorpyrifos and dimethoate (and its metabolite omethoate) where the effect on health needed to be considered in more detail. The risk assessment for these pesticides is outlined in the bullet points below. Please see [Section 3](#) for the full details of the HSE assessment of the risks.

- Residues of chlorpyrifos were found at 0.06 mg/kg in tenderstem broccoli. As outlined in HSE's full risk assessment ([section 3](#)), EFSA's 2019 statement on the human health assessment of chlorpyrifos indicated that no toxicological reference values could be determined for chlorpyrifos. See [Section 3](#) for more detail.
- Regarding the short term exposure assessment, although we cannot conclude with certainty whether or not presence of food residues of chlorpyrifos at this level would have any effect on health after eating large portions (97.5th percentile consumption) of broccoli, HSE has stated a number of reasons why the detailed assessment they have performed is especially precautionary. Please refer to the full assessment of risk in [section 3](#) for further details.
- The EFSA 2019 statement on the human health assessment of chlorpyrifos also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). Regarding genotoxicity, we conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. However, we consider any such risks of adverse health effects are low due to the limited levels of exposure anticipated based on the chlorpyrifos residue found in broccoli in this report.
- Dimethoate and omethoate are chemically related pesticides and for toxicology purposes are considered together. Omethoate is also the main metabolite of

dimethoate. Residues of dimethoate and omethoate were found in broccoli at levels up to 0.04 mg/kg and 0.01 mg/kg respectively. The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and omethoate, due to a lack of a fully supporting toxicological database. We think that at the anticipated highest exposures following consumption of broccoli containing these residues of dimethoate and omethoate, there is unlikely to be a risk of ill health effects based on short term toxicity.

- The EFSA Conclusion (2018) for dimethoate also includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of dimethoate and omethoate are undesirable due to the uncertainty regarding genotoxicity at low doses. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the dimethoate and omethoate residues found in broccoli in this report.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

In HSE’s long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. One of these samples contained residues from pesticides which belong to similar chemical groups, and may have similar toxicological effects. So the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample which contained chlorpyrifos and dimethoate (and its metabolite omethoate). HSE’s combined risk assessment on this combination concluded that the presence of chlorpyrifos in the sample does not significantly contribute to the overall combined risk of an effect on acetylcholinesterase (please see the glossary on page 125) when compared to that identified for dimethoate and omethoate. Please refer to the conclusions of the single substance assessments above and in [section 3](#) for chlorpyrifos and dimethoate (and omethoate). For full details of HSE’s assessment please refer to [section 3](#).

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We needed to consider the risk assessment for a sample of tenderstem broccoli containing residues of chlorpyrifos and dimethoate (and its metabolite omethoate) in more detail. Please see above and [Section 3](#) for more detail.

It is very unusual to find any exceedances of the MRL in broccoli surveys (just one exceedance prior to 2021 since 2015). Chlorpyrifos was not found in recent surveys (2015 and 2018). We will continue to test this food as part of the GB co-ordinated programme.

Residues measured above the MRL

The laboratory detected 2 residues above the MRL in broccoli. Details are available in [Table 5](#).

HSE have passed details of the following sample to FSA for further consideration. Further details are in [Table 4](#)

- 1 sample from Kenya containing chlorpyrifos at 0.06 mg/kg and dimethoate at 0.04 mg/kg

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Broccoli (NI)

Samples tested

6 samples were tested for up to 371 pesticide residues

Fresh

- 3 samples came from the EU

Frozen

- 1 sample came from the UK
- 2 samples came from the EU

Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 4 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Cheese (soft) (GB)

Samples tested

24 samples were tested for up to 113 pesticide residues

Brie

- 2 samples came from the EU

Camembert

- 1 sample came from the UK
- 6 samples came from the EU

Cottage Cheese

- 3 samples came from the UK

Cream Cheese

- 1 sample came from the UK

Feta

- 2 samples came from the EU

Mozzarella

- 1 sample came from the UK
- 2 samples came from the EU

Ricotta

- 4 samples came from the EU

Soft Cheese

- 1 sample came from the UK
- 1 sample came from the EU

The country of origin of samples may not be the same as the country where the cheese was produced. It may be where the cheese was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 19 samples contained no residues from those sought
- 5 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessment

None of the samples contained more than one residue, so we did not carry out a combined risk assessment.

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Cheese (soft) (NI)

Samples tested

19 samples were tested for up to 38 pesticide residues

Brie

- 3 samples came from the EU

Camembert

- 1 sample came from the UK
- 6 samples came from the EU

Cream Cheese

- 1 sample was imported from outside the EU

Mozzarella

- 4 samples came from the EU

Other

- 1 sample came from the EU

Ricotta

- 1 sample came from the EU

Soft Cheese

- 2 samples came from the EU

The country of origin of samples may not be the same as the country where the cheese was produced. It may be where the cheese was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 19 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)

- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Eggs (GB)

Samples tested

24 samples were tested for up to 115 pesticide residues

Duck

- 2 samples came from the UK

Hens

- 22 samples came from the UK

Pesticide residues detected from those sought

- 24 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- 7 samples were labelled as organic. None contained residues from those sought

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Eggs (NI)

Samples tested

3 samples were tested for up to 115 pesticide residues

Hens

- 3 samples came from the UK

Pesticide residues detected from those sought

- 3 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Fish (white) (NI)

Samples tested

19 samples were tested for up to 38 pesticide residues

Basa

- 1 sample was imported from outside the EU

Cod

- 6 samples were imported from outside the EU

Haddock

- 5 samples were imported from outside the EU

Pollock

- 3 samples were imported from outside the EU

Sea bass

- 2 samples came from the EU

Sea bream

- 1 sample was imported from outside the EU

Sole

- 1 sample came from the UK

Where no sea area information is available, the country of origin on the packaging does not necessarily indicate where the fish was caught or farmed. It could be where it was landed or processed or where it was packed for retail sale.

Pesticide residues detected from those sought

- 15 samples contained no residues from those sought
- 4 samples contained residues above the reporting limit
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in this sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Grapes (GB)

Samples tested

17 samples were tested for up to 395 pesticide residues

- 17 samples were imported from outside the EU

Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

A sample of grapes contained a residue of ethephon at 0.9 mg/kg where the effect on health needed to be considered in more detail. Based on the HSE Chemicals Regulation Division's risk assessment (please see [Section 3](#)) of the residue detected we consider a short term effect on health to be unlikely.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to '[how HSE perform the risk assessments](#)' for further details.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We needed to consider one sample in more detail (with a residue of ethephon at 0.9 mg/kg). We consider that an effect on health would be unlikely.

We have surveyed grapes for a number of years and publish the results of this sampling throughout the year as part of our rolling reporting programme as we know that a high percentage of the samples are imported.

Further Information

- [Summary table of results](#)
- [Survey Design](#)

- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Grapes (NI)

Samples tested

4 samples were tested for up to 395 pesticide residues

- 4 samples were imported from outside the EU

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 3 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

We have surveyed grapes for a number of years and publish the results of this sampling throughout the year as part of our rolling reporting programme as we know that a high percentage of the samples are imported.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Melon (GB)

Samples tested

25 samples were tested for up to 373 pesticide residues

Cantaloupe

- 5 samples were imported from outside the EU

Galia

- 4 samples were imported from outside the EU

Honeydew

- 4 samples were imported from outside the EU

Piel De Sapo

- 2 samples were imported from outside the EU

Prepared Fresh Melon (except watermelon)

- 2 samples came from the UK

Prepared Fresh Watermelon

- 4 samples came from the UK

Watermelon

- 3 samples were imported from outside the EU

Yellow

- 1 sample was imported from outside the EU

The country of origin of the samples may not be the same as the country where the melon was produced. It may be where the melon was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 4 samples contained no residues from those sought
- 21 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant samples. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Melon (NI)

Samples tested

6 samples were tested for up to 373 pesticide residues

Cantaloupe

- 1 sample was imported from outside the EU

Honeydew

- 3 samples were imported from outside the EU

Prepared Fresh Watermelon

- 2 samples came from the UK

The country of origin of the samples may not be the same as the country where the melon was produced. It may be where the melon was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 5 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Milk (GB)

Samples tested

72 samples were tested for up to 112 pesticide residues

Cows milk

- 63 samples came from the UK

Goats milk

- 9 samples came from the UK

Pesticide residues detected from those sought

- 72 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- 22 samples were labelled as organic. None contained residues from those sought

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Milk (NI)

Samples tested

7 samples were tested for up to 112 pesticide residues

Cows milk

- 7 samples came from the UK

Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Additional comments by the PRiF

The laboratory did not detect any residues at or above the reporting limit.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Mushroom (GB)

Samples tested

36 samples were tested for up to 395 pesticide residues

Button

- 14 samples came from the UK
- 6 samples came from the EU

Chestnut

- 11 samples came from the UK
- 3 samples came from the EU

Oyster

- 1 sample came from the EU

Portabello

- 1 sample came from the UK

The country of origin of the samples may not be the same as the country where the mushroom was produced. It may be where the mushroom was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 20 samples contained no residues from those sought
- 16 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- 16 samples were labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We note the findings of mepiquat in three samples of mushrooms most likely arose from straw treated with this pesticide. We understand that the labels of the pesticides used to treat wheat contain information not to use the treated wheat for compost. We have asked HSE to follow this up with the industry.

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in chestnut mushrooms. Details are available in [Table 5](#).

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Mushroom (NI)

Samples tested

3 samples were tested for up to 395 pesticide residues

Button

- 3 samples came from the EU

The country of origin of the samples may not be the same as the country where the mushroom was produced. It may be where the mushroom was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 2 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Peppers (GB)

Samples tested

30 samples were tested for up to 394 pesticide residues

Fresh

- 5 samples were imported from outside the EU
- 25 samples came from the EU

The country of origin of samples may not be the same as the country where the peppers were produced. It may be where the peppers were processed, where they were packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 10 samples contained no residues from those sought
- 20 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information

- Pesticides sought and residues found

Peppers (NI)

Samples tested

3 samples were tested for up to 394 pesticide residues

Fresh

- 1 sample was imported from outside the EU
- 2 samples came from the EU

The country of origin of samples may not be the same as the country where the peppers were produced. It may be where the peppers were processed, where they were packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 2 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

One sample contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information

- Pesticides sought and residues found

Potatoes (GB)

Samples tested

20 samples were tested for up to 390 pesticide residues

- 18 samples came from the UK
- 2 samples were imported from outside the EU

Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- 13 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Potatoes (NI)

Samples tested

3 samples were tested for up to 390 pesticide residues

- 3 samples came from the UK

Pesticide residues detected from those sought

- 2 samples contained no residues from those sought
- 1 sample contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

None of the samples contained more than one residue, therefore we did not carry out a combined risk assessment.

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Raspberry (GB)

Samples tested

24 samples were tested for up to 369 pesticide residues

Fresh

- 4 samples were imported from outside the EU
- 8 samples came from the EU

Frozen

- 5 samples came from the UK
- 7 samples were imported from outside the EU

The country of origin of the samples may not be the same as the country where the raspberry was produced. It may be where the raspberry was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 6 samples contained no residues from those sought
- 18 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- 1 sample was labelled as organic. None contained residues from those sought

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Further Information

- [Summary table of results](#)

- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Raspberry (NI)

Samples tested

7 samples were tested for up to 369 pesticide residues

Fresh

- 1 sample came from the EU

Frozen

- 3 samples came from the UK
- 3 samples were imported from outside the EU

The country of origin of the samples may not be the same as the country where the raspberry was produced. It may be where the raspberry was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 1 sample contained no residues from those sought
- 6 samples contained residues above the reporting limit
- 1 sample contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

None of the residues detected by the laboratory would be expected to have an effect on health.

In HSE's long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. We do not expect these residues to have an effect on health, either separately or in combination. The pesticide residues found in each sample do not include more than one of the pesticides from the groups that the Health and Safety Executive (HSE) consider separately.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We would not expect any of the residues detected to have an effect on health.

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in raspberries. Details are available in [Table 5](#).

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Rice (GB)

Samples tested

24 samples were tested for up to 386 pesticide residues

Basmati

- 13 samples came from the UK
- 2 samples were imported from outside the EU
- 1 sample came from the EU

Brown

- 2 samples came from the UK
- 1 sample came from the EU

Other

- 4 samples came from the UK
- 1 sample came from the EU

The country of origin of the samples may not be the same as the country where the rice was produced. It may be where the rice was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 13 samples contained residues above the reporting limit
- 2 samples contained residues above the MRL
- 5 samples were labelled as organic. None contained residues from those sought

Risk assessments

Following screening assessment, there were two pesticides where the effect on health needed to be considered in more detail. These are outlined in the bullet points below. For each of these, please see [Section 3](#) for the full details of the HSE assessment of the risks.

- Residues of chlorpyrifos were found at 0.004 mg/kg in rice. Based on the low short term intakes, HSE considers that a short term effect on health is not expected. As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the residue level found.
- Residues of tricyclazole were found at up to 0.1 mg/kg in rice. Based on HSE's assessment of the risk we conclude that a short term effect on health is not expected. Tricyclazole is a fungicide that has not been an approved substance in the UK or the EU. There is uncertainty about the potential for tricyclazole to cause

genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of tricyclazole in food undesirable. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the levels of up to 0.1 mg/kg in rice.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

In HSE’s long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Some samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We needed to consider two residues in more detail and following HSE’s assessment of risk, we concluded that short term effects on health would be unexpected. We also concluded any risk of adverse health effects due to genotoxicity to be low.

The MRL for tricyclazole was lowered on 30 June 2017. However transitional provisions applied, meaning that products that were imported or placed on the market before this date were subject to the previous MRL. Members feel that the change in MRL was sufficiently long enough ago that exceedances of the MRL should no longer be seen. We have asked HSE for the issue to be raised again with the rice industry.

Residues measured above the MRL

The laboratory detected 3 residues above the MRL in Basmati rice. Details are available in [Table 5](#).

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Rice (NI)

Samples tested

7 samples were tested for up to 386 pesticide residues

Basmati

- 5 samples came from the UK

Other

- 1 sample came from the UK

White

- 1 sample came from the UK

The country of origin of the samples may not be the same as the country where the rice was produced. It may be where the rice was processed, where it was packed for consumer purchase or the address of the brand owner.

Pesticide residues detected from those sought

- 4 samples contained no residues from those sought
- 3 samples contained residues above the reporting limit
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Risk assessments

Following screening assessment, there were three pesticides where the effect on health needed to be considered in more detail. These are outlined in the bullet points below. For each of these, please see [Section 3](#) for the full details of the HSE assessment of the risks.

- Residues of chlorpyrifos were found at 0.005 mg/kg in rice. Based on the low short term intakes, HSE considers that a short term effect on health is not expected. As outlined in HSE's full risk assessment ([section 3](#)), EFSA issued a 2019 statement on the human health assessment of chlorpyrifos which includes a consideration of the potential for genotoxicity (whether damage to genetic material can occur). We conclude that on a precautionary basis any findings of chlorpyrifos are undesirable due to the uncertainty regarding genotoxicity. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the residue level found.
- Residues of tricyclazole were found at 0.01 mg/kg in rice. Based on HSE's assessment of the risk we conclude that a short term effect on health is not expected. Tricyclazole is a fungicide that has not been an approved substance in the UK or the EU. There is uncertainty about the potential for tricyclazole to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of tricyclazole in food undesirable. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the levels of in rice.

- A sample of rice contained a residue of carbofuran at 0.001 mg/kg. Carbofuran is a fungicide that has not been an approved substance in the UK or the EU. Although a low level residue, where a short term effect on health is not expected, there is uncertainty about the potential for carbofuran to cause genetic damage at low doses. Therefore, on a precautionary basis we consider any findings of carbofuran in food undesirable. However, we consider any risks of adverse health effects are low due to the limited levels of exposure anticipated based on the level of 0.001 mg/kg in this rice sample.

Other risk assessment screening work undertaken did not indicate any other expectation of effects on health. Please refer to [‘how HSE perform the risk assessments’](#) for further details.

In HSE’s long term exposure screening assessment for these Quarter 1 2021 samples, NI and GB samples were combined.

Combined risk assessments

Two samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant sample. We would not expect any of these combinations to have an effect on health.

Further information of how HSE consider combined risk assessments is in [Section 3: HSE assessment of risk](#).

Additional comments by the PRiF

We needed to consider three residues in more detail and following HSE’s assessment of risk, we concluded that short term effects on health would be unexpected. We also concluded any risk of adverse health effects due to genotoxicity to be low.

The MRL for tricyclazole was lowered on 30 June 2017. However transitional provisions applied, meaning that products that were imported or placed on the market before this date were subject to the previous MRL. Members feel that the change in MRL was sufficiently long enough ago that exceedances of the MRL should no longer be seen. We have asked HSE for the issue to be raised again with the rice industry.

Further Information

- [Summary table of results](#)
- [Survey Design](#)
- [Glossary](#)
- [HSE Assessment of Risk](#)
- Detailed reference information in an accessible format at [Pesticide Residues in Food Quarterly Data](#), including
 - Brand name, sampling point and origin information
 - Pesticides sought and residues found

Section 2: Sample details and supplier responses

Sample details

The sample details are published on [Pesticide Residues in Food Quarterly Data](#) as a dataset in ODS format.

About sample information

The following information is available on each sample collected this quarter:

- Date and place of collection
- Description (e.g. 'runner bean', organic milk);
- Country of origin or manufacture;
- Brand name and packer/manufacturer; and
- Residues detected (results shown in green indicate residues above the MRL).
- Where the brand name of a sample is given the produce involved may have been on sale in other retail premises at the same time.

The description and country of origin are taken from labelling on the food or at the point of sale. The country of origin of processed food may not be the country where the unprocessed produce was produced. This is true even of food that has undergone minimal processing, such as meat that has been butchered or frozen vegetables.

Samples with residues above the MRL are in bold, green text.

Some brand name details have been withheld – these will be published once enquiries are complete.

The Government's 'brand naming' policy

The Government has decided that brand name information should be published as part of the Government food chemical surveillance programme. Brand names have been published for most pesticide residue surveys since 1998. Certain samples are excluded from the release of brand name information. These include samples taken as part of any pesticide residues enforcement programme and those taken as part of surveys to study individual people/farms. This policy was reviewed in 2000/1, when Ministers agreed to its continuation.

Where we find residues above an MRL or the presence of non-authorized plant protection product, brand owners/retailers/ growers are notified of the result in advance of publication of reports and given four weeks to comment.

Interpreting brand name information

There is no ready definition of what constitutes a brand in all cases. For clearly branded produce like breakfast cereals or biscuits the "brand owner" is shown. In the case of "own brand" goods this may be one of the multiple retailers. For fruit and vegetables, the retailer is generally shown. For meat, milk and most other animal products the retailer is also

generally shown. Finally, for all commodities the country of origin is shown where this was displayed either on the produce or in the store.

Our programme aims to take samples of produce in approximate proportion to the market share of the main retailers. This has been done to ensure we obtain an accurate representation of a sector (e.g. fruit and vegetables).

Individual programmes are not capable of generating statistically valid information on residues in particular crops from particular retailers. This would require the collection of a much larger number of samples: either substantially increasing costs or greatly reducing the range of different foods sampled in any one year. Therefore, results from an individual survey cannot be taken as a fair representation of the residues status of any particular brand.

However, we do collect samples from a variety of outlets in a range of locations, over a period of years. Successive programmes should therefore help generate information on the typical residues profile of particular types of produce and on major trends in the incidence and levels of pesticides. It should be noted that this quarterly report is not intended to give a comprehensive comparison with previous surveys of the same commodities.

A particular issue arises in relation to the country of origin of fruit and vegetables. The origins included in the reports are those recorded either on the produce or in the store. However, it is not uncommon for mixing to occur on shop shelves. We have responded by increasing the proportion of pre-packed goods sampled. However, pre-packed samples are not available for some produce in some stores and it could also introduce bias to surveys if loose produce were not sampled. Loose produce is therefore sampled but the origin of the sample should be interpreted with a degree of caution.

Action taken by HSE

HSE wrote to:

- The suppliers of all samples containing residues above the MRL
- The authorities of the exporting countries of all samples containing residues above the MRL
- The suppliers of GB and NI samples that contained residues of actives which do not have a plant protection product authorised for the crop detected in.
- The Organics branch of Defra about samples that were labelled as organic and contained any residues of pesticides.
- The suppliers and certification organisation of all organic samples containing any residues of pesticides.

Recipients of the letters are given 4 weeks to provide a statement for inclusion in the report. The Expert Committee on Pesticide Residues in Food reviews any replies received.

Supplier responses

Rice sample 0144/2021: Response from Veetee

Despite rigorous control checks (pesticide testing) in country of origin, it was difficult to get 100% compliance in previous years. Veetee has now changed to contract farming so that farmers work with approved pesticides, and we can eliminate any irregularity. Veetee are still making all efforts to ensure all food coming into UK is fully compliant with current regulation.

Section 3: HSE assessment of risk

The surveillance programme is designed to enable the regulatory authorities to check that pesticides are being found at levels, as expected, under the MRLs. This confirms that the regulatory processes are working correctly, and as part of this, that pesticide users are complying with any specified conditions that were part of the authorisation. In addition, this work checks that dietary intakes of residues are within acceptable limits. This may be more challenging when pesticide residues are found in food products that have not been grown in the UK or EU, notably when older pesticides have been used. One of the roles of the PRiF, using the work of HSE, is to call out any pesticide residue which is higher than expected and explain more about any risks to consumers from this.

This section details how risks from dietary intakes are assessed.

When assessments are carried out

HSE performs screening assessment for each residue and commodity combination to identify residue levels that would lead to intakes above the relevant health based reference doses (these are also sometimes referred to as toxicological reference values “TRVs”). Further information on this screening approach is available on request from HSE. We then present detailed assessments for every case where the actual residue level found could lead to an intake by any group above the reference dose.

Assessing Dietary intakes

Assessing the acceptability of dietary intakes is complicated. HSE carries out consumer risk assessments for both short-term (peak) and long-term intakes. These assessments use information on food consumption collected in UK dietary surveys in conjunction with the residue levels we find. Occasionally, HSE uses additional pesticide specific information on the losses of residues that occur during preparation and/or cooking of food.

How the assessment is carried out

Short term risk assessment

HSE calculate short-term intakes (also called NESTIs) using consumption data for high-level (97.5th percentile) consumers, based on single-day consumption values and the highest residue found in a food commodity. The residue found is multiplied by a variability factor to take account of the fact that residues may vary between individual items that make up the sample analysed. This is why in some of our detailed risk assessments we refer to some of the general variability factors (of 5, 7 and 10) that are applied in short term risk assessments. Sometimes, regulatory assessment of data for a pesticide can support an alternative specific value of the variability factor, and where justified, HSE will apply these to the risk assessment and explain this. The estimated intake is compared to the Acute Reference Dose (ARfD). This is done for ten consumer groups; adults, infants, toddlers, 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, vegetarians, elderly living in residential homes and elderly living in their own homes.

Long term risk assessment

HSE also calculate long-term intakes (NEDIs) for high-level (97.5th percentile) consumers, but in this case the consumption data are high-level long-term values rather than peak single-day events. Similarly, the residue values used reflect long-term average levels (we use the median value across each commodity type) rather than occasional high values. Again, these estimates are made for the ten consumer groups. In this case the estimated intake is compared to the Acceptable Daily Intake (ADI).

Where do we get Reference Doses from?

The reference doses (ADI, ARfD or other suitable health based reference values such as Tolerable Daily Intake (TDI, sometimes used instead of an ADI)) are set following regulatory assessment. In the UK, these values will be reviewed prior to establishment by the Expert Committee on Pesticides (ECP). We currently use reference values from a range of respected sources, including the EU and values set by EFSA. Up to 2019, the UK was part of a harmonised approach to the assessment of pesticide substances in the EU, and we have taken part in the peer review of previously established values. PRiF will also use values from other respected international sources, such as those established by The Joint FAO/WHO Meeting on Pesticide Residues (JMPR, which evaluates and publishes residues and toxicological evaluations of pesticides) and levels set by regulatory authorities in other countries. For a small number of pesticides, the reference doses used have been determined by HSE (e.g. prothiofos, tecnazene). These have not been independently peer-reviewed and should therefore be regarded as provisional.

Further explanation of the models we use and application to the PRiF work

We use Deterministic models

The assessments we use are 'deterministic' which means we use a defined level of input (such as a median or highest residue and 97.5th percentile dietary consumption values). More information on the deterministic intake assessments is available on HSE's website: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links. See below for an explanation of probabilistic models, where the inputs into the assessments can be varied and more realistic assessments reflecting a range of possible scenarios can be modelled.

Detailed Risk Assessment work is carried out before pesticides are authorised

The fundamental full complement of risk assessment work for pesticides is done at the pre-authorisation stage considering the residues data packages when trials reflecting the label uses of the pesticides are assessed. These trials profile the highest likely residues that might arise when the pesticides are used as intended (crops and permitted doses of use). MRLs are set on the basis of these data sets, and the post-approval monitoring work then serves as a check for whether residues found are in line with this prior expectation. The risk assessments supporting the MRLs assess the highest residue observed in each crop (or animal product) for short term assessment. For the long term MRL assessments, the median residues, across all crops and animal products are taken together to assess combined intakes over the long term as 'total dietary intakes', taking account of all possible food exposures. Authorisation for pesticide products can only be granted where these assessments of dietary intake do not exceed the health based reference values.

MRL considerations

Although MRLs are not safety levels, an MRL would not be established if the residue concentrations measured in the supervised trials used to support the MRL would give rise to health concerns. In most cases residues present at the MRL result in intakes below the ARfD and the ADI. So even if the MRL is exceeded this does not always lead to an intake above the ARfD or ADI.

What happens if we find a pesticide intake above an ADI or an ARfD?

In addition, an estimated intake in the monitoring work that exceeds the ADI or ARfD does not automatically result in concerns for consumer health, because a protective approach is used in setting the ADI and ARfD. In the unusual circumstance of an intake exceeding the ADI or ARfD, HSE undertake an evaluation of the toxicological data, and we present details of this assessment.

When we present the outcomes of risk assessments, we provide a conclusion on the possible impact on human health based on the degree of concern following the HSE assessment of risk. These conclusions keep to the following order of increasing severity:

Effects on health are unexpected (toxicological reference values not exceeded) < unlikely risk (of effects on health) < low risk < higher risk (exposures are undesirable⁴). Most detailed consumer intake assessments that we present with the PRiF reports are for short-term exposure rather than chronic exposure. This is because in most cases the monitoring data show the majority of samples tested contain residues below the reporting limit and so chronic exposure would not present a concern. The reporting limits are set at suitable low levels based on analytical laboratory procedures so that dietary intakes are typically far below the health based reference values.

Monitoring data presented in the PRiF quarterly reports provides a “snapshot view” of the residues found in specific foods in a survey usually conducted over a 12 week period and limited to around 30 samples for most commodities. It is important not to use isolated findings of higher than expected residues in small surveys to make judgements on long term effects over a lifetime exposure.

For PRiF work, long-term exposure assessments use median residue levels, rather than the highest residues found. For quarterly assessment (data obtained over three months only) we currently only assess long term dietary assessment commodity by commodity and not as total dietary intakes across commodities. Even where a number of samples in a PRiF commodity survey do contain the same pesticide, it is very rare that the ‘screening assessment’ we undertake leads to the need for a more detailed assessment to be presented (only where the dietary intakes exceed the ADI or TDI).

We do not see a high number of PRiF samples for any pesticide that exceed an MRL. This means that, over time, median residue levels found in PRiF monitoring don’t often exceed the median residues used in the trials assessed at the time of MRL setting and so don’t often require consideration of long term effects. When HSE does need to assess long term effects, it is likely to overestimate exposure to an assessed pesticide residue in a single

⁴ Furthermore, PRiF will always conclude that on a precautionary basis any findings of genotoxic substances in food are undesirable (please see the explanation regarding genotoxicity in the below section on ‘Implications for health’).

food item. For a single commodity risk assessment, we assume high level (97.5th percentile) consumption at the median residue level in that food for each day of lifetime. Although the HSE long term assessments by their quarterly nature are indicative only, the assessment we perform is conservative. Furthermore, alternative published assessments (for example those considering trends and large bodies of data) are available which further consider the long term exposure to pesticides⁵. Some pesticides contribute more significantly to long term dietary intakes across commodities based on their toxicology and prevalence (such as chlorpyrifos, cyfluthrin, deltamethrin, diazinon, dieldrin, dimethoate, dithiocarbamates, fenamiphos, fipronil, imazalil, lambda-cyhalothrin, omethoate and pirimiphos-methyl), based on the chronic exposure assessments presented in EFSA, 2020⁶).

We will continue to have focus on residues of consumer relevance, when they are found, in both UK and NI produced and imported produce. As pesticide use changes, including when there are impacts of regulatory action on pesticides, the profiles of residues in the monitoring can change over time. For example, chlorpropham, which was previously found at levels well above the reporting limits in potatoes, is no longer permitted for use in the UK and EU, and this pesticide will no longer have dietary intakes which take up a substantial portion of its health based reference values.

Implications for health

Where intakes exceed a reference dose, it is necessary for the underlying toxicological studies (animal studies) to be considered to enable the significance for the consumer of such an exceedance to be understood. Toxicological studies supplied by the registrants in the regulatory data packages are conducted using different doses to determine the nature of any ill health effects as well as the levels at which such effects can be expected to occur.

Toxicological studies that we refer to and use in the HSE risk assessments are conducted using test animals to identify the highest experimental dose that causes no detectable adverse effects (the NOAEL). Where there is more than one relevant toxicological study, the lowest appropriate NOAEL for the most sensitive adverse effect is typically used. There is some uncertainty in extrapolating between animals and people and it is therefore important to use a 'safety factor' to account for sources of variation. This safety factor is incorporated (by dividing the NOAEL by the safety factor) in deriving a reference dose, either an ADI or an ARfD, to which consumer intakes are compared. A safety factor therefore extrapolates from the animal testing to the general population. Factors in the order of x100 are commonly used, x 10 for animal to man, and x10 for within human population differences in sensitivity. However, toxicologists may propose different values (e.g. from 5 to 1000) based on scientific reasoning in accordance with study designs and the quality of the data that has been generated from the studies.

⁵ Total diet studies (e.g. those performed by US FDA [US FDA total diet study program](#), various Nougadère A et al., publications in Environment International journal on TDS in France); see also EFSA evaluations of chronic exposures to pesticides (2020 and 2021 examples included in the list of EFSA publications on cumulative exposure to pesticides outlined in this section (under Multiple residues)). annual EU monitoring data published on the EFSA website: 2018 report (published in 2020) noted below at <https://doi.org/10.2903/j.efsa.2020.6057>.

⁶ EFSA (European Food Safety Authority), Medina-Pastor P and Triacchini G, 2020. The 2018 European Union report on pesticide residues in food. EFSA Journal 2020;18(4):6057, 103 pp. <https://doi.org/10.2903/j.efsa.2020.6057>

In order to ensure exposures to pesticides do not pose unacceptable risk to humans a wide range of investigations are performed. Most of these are performed on experimental animals because the only end-points that can be examined in human volunteers are those involving observation or blood and urine sampling. Human volunteer studies involving pesticides are not generated in current regulatory work. There is debate at the international level as to whether human studies that have been generated should be used for risk assessment purposes. The UK policy is not to use these data in pre-authorisation assessments which support the registration of a pesticide; the JMPR chose to apply judgement in the appropriate use of these data if available. The HSE risk assessments will usually refer to test animal species, such as dog, rat, and rabbit. All toxicological work is undertaken based on principles of minimising animal distress. Where scientifically valid human data are available the risk assessments will refer to these as they reduce the uncertainty in the assessment. Therefore, human data is only referred to in more limited circumstances.

Acute (short term) toxicology is not a concern for all pesticides, as some are not acutely toxic. In terms of the pesticides that have been found in fruit and vegetables through the surveillance programme an acute risk assessment would not be necessary on the following examples: maleic hydrazide, diphenylamine, kresoxim-methyl, and quintozene.

During regulatory assessment, careful consideration is given to any pesticides that may exhibit any potential to be genotoxic (able to damage genetic material) in live animals. In the PRiF programme we note residue types that have been shown in the toxicological data sets to have genotoxic potential or those where data are suggestive of genotoxicity but not certain. There is a small number of cases of older pesticides, likely found only in imported foods, that might be genotoxic. These are examples where modern data to investigate the true genotoxic potential are not expected to be made available. In such situations, we might conclude on a precautionary basis that any findings of these pesticides are undesirable due to the uncertainty regarding genotoxicity, and at low residue levels any risks of adverse health effects are low due to the limited levels of exposure anticipated. PRiF uses low reporting limits for these pesticides to detect these residues even at very low levels, as we know they are of particular interest to consumers.

Consumption data and refining the risk assessment

Consumption values

As the surveillance programme monitors residues in all types of food, from raw commodities (e.g. potatoes) to processed (e.g. wine), dried (e.g. dried fruit) and composite foods (e.g. fruit bread), consumer risk assessments are specifically tailored to address processed and mixed food products. Sometimes this can be affected by availability of consumption data. For example, for pâté, we assess this using consumption data for liver (all types of liver), and for fish pâté we use consumption data for fish (all sources and types of fish). However, we use specific consumption data where FSA have provided data to us (e.g. data on orange juice, dried grapes, and bread). Consumption data are available for most raw commodities, but where data are limited then we will suggest using alternative data. This may involve considering other commodities (e.g. using potato data as 'surrogate' for sweet potato), or alternative sources of consumption data such as EU PRIMo or JMPR consumption and dietary assessment models, to consider items that do not currently feature in UK data sets. Where alternative data are used in our screening and written assessments we explain this in our presentation of the risk assessment work for

each quarterly report (for examples, please see the bullet points before the table of detailed risk assessments in [section 3](#)).

Fruit and vegetables with removable peel

For fruit and vegetables that have peel or skin that might not be consumed we present alternative risk assessments for 'without peel -flesh only' where peel versus pulp residue distribution data are available. As standard, we present an assessment for 'all of the peel' consumed. It is not expected that consumers will always eat peel, so these standard assessments are considered to be highly cautious and not necessarily realistic. Further data are being generated to better understand whether some people do eat the peel of these, and if so how much of the peel they tend to eat.

Dithiocarbamate Residues

Dithiocarbamate residues are determined as carbon disulphide which is a common product from different dithiocarbamate pesticides. For the risk assessment we take a precautionary approach. For short term assessment the worst case dithiocarbamate residue is calculated by assuming the residue is derived from thiram (a molecular weight conversion is applied to estimate the level of residue based on thiram) and this is compared to the ARfD for thiram. Where it can be confirmed that a specific dithiocarbamate was applied the equivalent residue of the specific active substance is estimated and the intake compared to the appropriate reference dose. We only present a detailed risk assessment when dietary intake exceeds either the thiram or other suitable reference dose.

The analysis of dithiocarbamates is further complicated by an expectation that some types of crops, such as members of *Brassicaceae* (e.g. watercress) and *Caricaceae* (e.g. papaya) might contain natural sources of sulphur compounds that could be also determined as carbon disulphide during analysis in the laboratory. The PRiF will consider and explain in the report whether residues reported as dithiocarbamates could be from natural sources or whether they have arisen as a result of fungicide treatment.

Probabilistic Modelling

The standard 'deterministic' calculations of consumer exposure used in regulatory assessment and the HSE risk assessments for PRiF work use realistic consumption data and residue levels. However, they tend to overestimate intakes in most circumstances. This is due to the assumptions used; fruit and vegetables would contain high levels of residue in an individual unit and that these would be consumed by high-level consumers. They do not take into account the possible range of residue levels and consumption distributions that may occur in reality. These possible combinations of residues and consumption levels can be taken into account using modelling/simulation techniques to produce probability distributions of residue intake levels to indicate the range of consumer intakes, presented as a probabilistic assessment of consumer exposure. These techniques are not yet routinely used to estimate dietary intakes of pesticide residues.

Multiple residues and other developments in risk assessment for pesticides

The risk assessment process is not standing still. We are aware that some consumers are concerned by the 'cocktail effect'- the possible implications of residues of more than one

chemical occurring in, say, a single portion of fruit or vegetables or the interaction between mixtures of pesticides and veterinary medicines at residue levels.

Where more than one pesticide residue is found in a sample, we consider the need for further assessment. The question of which pesticides should be assessed together remains a challenge due to the complexity of the mixtures. In the PRiF work currently, we consider some combinations that we think are a priority (based on toxicological profile or prevalence of the co-occurring residues that are related to one another chemically). If more than one triazole, or more than one organophosphate/carbamate is found or the following combinations captan/folpet, BAC/DDAC, chlormequat/mepiquat, we will undertake an additional risk assessment. In a 'first step' screening assessment approach we will consider whether the sum of the dietary intakes of each pesticide taken together in that commodity (when expressed as a % of its own reference value) exceeds a total of 100. If this value is not exceeded, then we do not anticipate that there would be an effect on human health and the assessment is not considered further. If this value (of 100) is exceeded (in the initial screen) then we would present a more detailed risk assessment, including a table to show the dietary intakes of each of the pesticides within the group, in the report. Further information is available on: [The HSE Pesticide Website](#). Search for the Data Requirements Introduction and Index and follow the 'consumer exposure' links.

International research is aimed at improving the regulatory assessment of mixtures of residues to help understand whether there are any health implications from any observed combinations of pesticide residues in food. In our work, PRiF aims to keep our assessments at a high regulatory standard taking account of current knowledge in the field. We will look to adapt as new risk assessment approaches develop. We are keen to ensure our reports reflect consumer concerns.

The Food Standards Agency (FSA) asked the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) to assess these concerns. Their report "Risk Assessment of Mixtures of Pesticides" was published in 2002⁷.

The Committee concluded that the probability of any health hazard from exposures to mixtures is likely to be small. Nonetheless, it identified areas of uncertainty in the risk assessment process and made recommendations for further work. These fell under the broad headings of regulatory, surveillance, research and public information issues. An action plan to take forward the recommendations was published by the FSA. A number of research projects were commissioned by the FSA to help progress the action plan.

Further to the work done by COT in 2002, combined assessment methodology has been taken forward at the international level, especially the European Food Safety Authority (EFSA) to develop methodology.

Much of the existing recent work on cumulative exposures to pesticides uses probabilistic models and large EU wide monitoring data sets. Notable work includes the EFSA publications on cumulative exposure cited below. On the basis of the work to date, including consumer assessment case studies, EFSA concludes, with varying degrees of certainty for all the population groups assessed, that consumer risk from dietary

⁷ [Foods Standards Agency Risk Assessment of Mixtures of Pesticides \(COT Report, 2002\)](#)

cumulative exposure is below the thresholds established by EU risk managers. Further information can be obtained from EFSA's publications, news updates, and FAQs:

- EFSA Feb 2021 (Statement): Comparison of cumulative dietary exposure to pesticide residues for the reference periods 2014–2016 and 2016–2018 [link](#)
- EFSA Feb 2021 (Scientific Report) Cumulative dietary risk assessment of chronic acetylcholinesterase inhibition by residues of pesticides [link](#)
- EFSA April 2020: News Pesticides: first cumulative risk reports published This work is centred on two case studies (outlined below) [link](#)
- EFSA April 2020: Cumulative risk assessment of pesticides: FAQ [link](#)
- EFSA April 2020: Cumulative dietary risk characterisation of pesticides that have acute effects on the nervous system [link](#)
- EFSA April 2020: Cumulative dietary risk characterisation of pesticides that have chronic effects on the thyroid [link](#)
- EFSA news update (Jan 2016) Pesticides: breakthrough on cumulative risk assessment [link](#)
- EFSA Sept 2019: Establishment of (CAGs) cumulative assessment groups (effects on thyroid) [link](#)
- EFSA Sept 2019: Establishment of (CAGs) cumulative assessment groups (effects on the nervous system) [link](#)
- EFSA Jan 2014: Outcome of the public consultation on the Scientific Opinion on the identification of pesticides to be included in cumulative assessment groups (CAGs) on the basis of their toxicological profile [link](#)
- EFSA Dec 2013: Scientific Opinion on the relevance of dissimilar mode of action and its appropriate application for cumulative risk assessment of pesticides residues in food [link](#)
- EFSA Sept 2009: Scientific Opinion on Risk Assessment for a Selected Group of Pesticides from the Triazole Group to Test Possible Methodologies to Assess Cumulative Effects from Exposure through Food from these Pesticides on Human Health [link](#)
- EFSA May 2008: Opinion of the Scientific Panel on Plant Protection products and their Residues to evaluate the suitability of existing methodologies and, if appropriate, the identification of new approaches to assess cumulative and synergistic risks from pesticides to human health with a view to set MRLs for those pesticides in the frame of Regulation (EC) 396/2005 [link](#)

Further publications on topics related to consumer risk assessment that are under development are as follows:

- UK Committee on Carcinogenicity (2019) guidance note (COC Guidance Statement G09) on LTL exposure assessment. [COC 2019 LTL Guidance](#)
- EFSA: Update: use of the benchmark dose approach in risk assessment (2016) [BMDL link](#)
- WHO guidance on genotoxicity (2020). [EHC 240 \(updated 2nd Ed\) genotoxicity](#)

HSE (UK) is participating in a number of international initiatives related to residues and risk assessment (OECD Working group on residue definitions, and the ongoing JMPR programme of evaluation work/attendance at CCPR (CODEX) and participating in JMPR/CCPR discussions of a technical nature on general considerations for risk assessment.

Further advances in risk assessment methodology will be taken into account in developing the approach to risk assessments in the future.

Risk Assessment- dietary intake assessments

Screening assessments have been done for all pesticides to check that predicted intakes are within the relevant health based reference values. A short term (acute) exposure assessment is not done for pesticides which are not acutely toxic where it has been established that an ARfD is not required. EU toxicological endpoints can be found in the [EU Pesticide database](#).

[Toxicological reference values set by the JMPR \(The Joint FAO/WHO Meeting on Pesticide Residues\) can be found in individual pesticide evaluations at JMPR Evaluations \(an up to date index to pesticide evaluations is available in the latest report\).](#)

The screening assessment uses the internationally agreed approach to long-term (chronic) and short-term (acute) consumer exposure assessment with UK food consumption data as detailed within the UK NEDI and NESTI models which are available on the [HSE website](#).

For the Quarter 1 (2021) assessments, the following approaches have been taken to refine these assessments according to case-by-case issues and to ensure that appropriate consumption values are used for less frequently consumed commodities where available food consumption data may be limited:

- Data on aubergine were used, although there are a low number of consumers in several of the sub-groups for aubergine. However, use of these consumption data was considered reasonable after comparison with alternative data.
- Data on banana were used for plantain
- Data on beans with pods were used for all forms of green beans, including speciality beans
- Data on fish were used for all forms of white fish
- Data on meat (excluding poultry and offal) were used for all forms of beef
- Data on cheese were used for all forms of cheese
- Data on melon were used for all forms of melon and watermelon
- Data on both blackberries and raspberries were used for raspberries, blackberries and blueberries

Short term dietary risk assessment – single substance assessments where exceedance of the ARfD has been identified during screening

Aubergine

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group [†]	ARfD (mg/kg bw)	Source
Aubergine	Chlorate	1.6	0.016	0.040 (4-6 year old child)	0.036	EFSA, 2015

Comment on risk assessment

The intakes for 4-6 year old children exceeded the ARfD.

If 4-6 year old children ate large portions of aubergine containing chlorate at 1.6 mg/kg, their intake of chlorate could be 111 % of the Acute Reference Dose. This intake is 1.1 times higher than a dose which caused no observed adverse effect in a controlled clinical study with daily dosing for 12 weeks. The European Food Safety Authority used this valid human study as the basis of the ARfD; an uncertainty factor was not considered necessary to take into account uncertainties caused by possible differences in susceptibility between people. In this study, there were no changes in the extensive battery of biochemical and physiological parameters monitored in any of the people from which clinical observations were made at the tested dose.

Based on the residue intake representing a small (1.1 times) exceedance of the dose tested in the clinical trial, HSE concludes that an effect on health is unlikely.

Bananas

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group [†]	ARfD (mg/kg bw)	Source
Bananas	Chlorpyrifos	0.004	0.000048	0.00033 (infant) 0.00022 (toddler)	No toxicological reference values established	EU, 2019

				0.00017 (4-6 year old child)		
				0.00012 (7-10 year old child)		
				0.000069(11-14 year old child)		
				0.000059 (15-18 year old child)		
				0.000057(vegetarian)		
				0.00005 (elderly – residential)		
				0.000051 (adult)		
				0.000045 (elderly – own home)		

Comment on risk assessment

EFSA (2019)⁸ has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed (short term) indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. This view is further supported by the US EPA ARfD of 0.006 mg/kg bw (published in 2020). Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

⁸ [EFSA 2019 statement on human health- chlorpyrifos](#)

Banana flesh after peeling

The dietary intakes calculated for when the peel is removed prior to consumption, indicate that there are no exceedances of indicative toxicological reference value and, in this case, a short term effect on health is not expected.

The below risk assessment only applies if all of the peel is consumed. This is because only 2% of the residue of chlorpyrifos remains (EFSA, 2017) in the flesh when the fruit is peeled.

Whole banana, including all the peel

We consider that an effect on nervous system would be unlikely. Any effect on health would depend on a number of factors which would need to come together at the same time (the high residue (0.004 mg/kg of chlorpyrifos in bananas) found in the sample being consumed by the most critical consumer infants), high residue in single fruit item, peak consumption levels (97.5th percentile), and a large proportion of peel from the fruit being consumed.

The following risk assessment assuming all of the peel is consumed, is presented, although the PRiF consider this to be a 'worst case' form of assessment for the reasons explained above:

The intakes for infants exceeded the HSE proposed indicative toxicological reference value for short term exposure. Intakes for all other consumer groups were below the reference dose. If infants ate large portions of bananas (including the skin) containing chlorpyrifos at 0.004 mg/kg their intake could be 110 % of the above mentioned HSE proposed indicative toxicological reference value for short term exposure. Intakes are approximately 900 times lower than the lowest intake in repeat-dose animal studies which was reported to cause effects in a developmental neurotoxicity study where pregnant rats were dosed from day 6 of pregnancy through until the pups were 11 days old. The US EPA (2020) established an ARfD for chlorpyrifos of 0.006 mg/kg bw; this supports the view that the proposed indicative toxicological reference value from HSE is precautionary. These exposures are undesirable but are unlikely to cause any adverse effect.

Overall, HSE concludes that a short term effect on health would be unlikely in the case of consumption including all of the peel, and not expected in the case of peeling banana prior to consumption.

Please refer to the section below on 'Substances that might be genotoxic' for HSE's conclusions regarding potential genotoxicity.

Beans with pods

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group†	ARfD (mg/kg bw)	Source
Beans with pods (Guar beans)	Chlorpyrifos	0.01	0.000023	0.000050 (infant) 0.000050 (toddler) 0.000037 (4-6 year old child) 0.000028 (vegetarian) 0.000027 (15-18 year old child) 0.000023 (adult) 0.000022 (elderly – own home) 0.000020 (7-10 year old child) 0.000019 (11-14 year old child) 0.000011 (elderly – residential)	No toxicological reference values established	EU, 2019

Comment on risk assessment

EFSA (2019)⁹ has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

⁹ [EFSA 2019 statement on human health- chlorpyrifos](#)

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. This view is further supported by the US EPA ARfD of 0.006 mg/kg bw (published in 2020). Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

None of the intakes exceeded the HSE proposed indicative toxicological reference value for short term assessment. Based on the low intakes, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group†	ARfD (mg/kg bw)	Source
Beans with pods (Guar beans)	Diafenthuron	1.1	0.0025	0.0055 (infant) 0.0055 (toddler) 0.0041 (4-6 year old child) 0.0030 (vegetarian) 0.0030 (15-18 year old child)	No ARfD established (an ADI of 0.003 is available)	Australia, 1993

Comment on risk assessment

The pesticide is old, uses in the EU are no longer authorised and toxicological reference values set by the EU and the JMPR are not available. An ADI of 0.003 mg/kg bw/day was set by Australia in 1993. HSE has considered the summary toxicological

data available for this Australian assessment¹⁰. It is not possible to conclude based on the available data whether the substance causes effects following acute exposures that would necessitate setting an Acute Reference Dose. However, if an ARfD were needed then it is considered likely to be at a higher value than the ADI. The assessment based on the exceedance of the ADI is therefore considered precautionary.

The intakes for infants, toddler, 4 to 6 year old children, vegetarians and 15-18 year old children exceeded the ADI. The highest intake was for infants

If infants ate large portions of Guar beans containing diafenthiuron at 1.1 mg/kg, their intake of diafenthiuron could be 184% of the Acceptable Daily Intake (ADI), (used as a surrogate in the absence of an Acute Reference Dose).

This intake is 55 times lower than a dose which caused no observed adverse effect in a dog study, where animals were dosed daily over a one year period, that was the basis of the ADI. Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 55 still sufficient to make an effect on health unlikely. Furthermore, the ADI is expected to be precautionary for the assessment of short term effects, and the reduced bodyweight effects that were used for the establishment of the ADI are unlikely to apply to single exposures, at the level from this residue found in this sample of beans.

Based on this assessment, we conclude that a short term effect on health is unlikely.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group[†]	ARfD (mg/kg bw)	Source
Beans with pods (Guar beans)	Dimethoate and Omethoate	0.04 (D: dimethoate) and 0.1 (O: omethoate)	D: 0.000092	0.00020 (infant) 0.00020 (toddler) 0.00015 (4-6 year old child) 0.00011 (vegetarian)	Not established	EU, 2019

¹⁰ [National \(Australian\) Registration Authority document for diafenthiuron.](#)

			O: 0.00023	0.00011 (15-18 year old child) 0.000092 (adult) 0.000087 (elderly – own home) 0.000081 (7-10 year old child) 0.000078 (11-14 year old child) 0.000044 (elderly – residential) 0.00050 (infant) 0.00050 (toddler) 0.00037 (4-6 year old child) 0.00028 (vegetarian) 0.00027 (15-18 year old child) 0.00023 (adult) 0.00022 (elderly – own home) 0.00020 (7-10 year old child) 0.00020 (11-14 year old child) 0.00011 (elderly – residential)	Not established	EU, 2019
--	--	--	---------------	--	-----------------	----------

Comment on risk assessment

The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and its metabolite omethoate, due to a lack of a fully supporting toxicological database. Both dimethoate and omethoate are not approved in the EU and pesticide products containing dimethoate were withdrawn in the EU and UK in 2020.

For dimethoate, EFSA (2018) stated an indicative value for a hypothetical toxicological reference value for short term exposure of 0.0001 mg/kg bw/day. Using this indicative value, estimated dietary intakes of dimethoate for infants, toddlers, 4 to 6 year old children, vegetarians and 15 to 18 year old children exceeded this reference value. The intakes of omethoate for all consumer groups exceeded this hypothetical short term toxicological reference value for dimethoate. The highest intake was for infants and toddlers.

If infants and toddlers ate large portions of Guar beans (beans with pods) containing dimethoate at 0.04 mg/kg their intake could be 201 % of the above mentioned hypothetical toxicological reference value for short term exposure. If infants and toddlers ate large portions of Guar beans (beans with pods) containing omethoate at 0.1 mg/kg their intake could be 501 % of this hypothetical toxicological reference value for dimethoate. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September, 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary

These exposures are undesirable but it is not clear if they may cause any adverse effect. The estimated exposures are not expected to inhibit acetylcholinesterase¹¹, the basis of previous evaluations of the safety of dimethoate and omethoate. Based on this assessment, HSE concludes that a short term effect on health is unlikely after eating large portions (97.5th percentile consumption) of Guar beans (beans with pods) containing the levels found in this report.

Please refer to the section below on 'Substances that might be genotoxic' for HSE's conclusions regarding potential genotoxicity

¹¹ this enzyme, acetylcholinesterase, is included in the Glossary on page 123

Broccoli

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group†	ARfD (mg/kg bw)	Source
Broccoli	Chlorpyrifos	0.06	0.00077	0.0015 (4-6 year old child) 0.0014 (7-10 year old child) 0.0013 (toddler) 0.0012 (infant) 0.0010 (vegetarian) 0.00093 (11-14 year old child) 0.00084 (15-18 year old child) 0.00077 (adult) 0.00076 (elderly – own home) 0.00041 (elderly – residential)	No toxicological reference values established	EU, 2019

Comment on risk assessment

EFSA (2019)¹² has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

¹² [EFSA 2019 statement on human health- chlorpyrifos](#)

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. This view is further supported by the US EPA ARfD of 0.006 mg/kg bw (published in 2020). Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

The intakes for all of the consumer groups exceeded this HSE proposed indicative toxicological reference value. The highest intake was for 4-6 year olds.

If 4-6 year olds ate large portions of broccoli containing chlorpyrifos at 0.06 mg/kg their intake could be 500 % of the above mentioned HSE proposed indicative toxicological reference value for short term exposure. This intake is approximately 200 times lower than the lowest intake in repeat-dose animal studies at which effects were observed in a developmental neurotoxicity study where pregnant rats were dosed from day 6 of pregnancy through until the pups were 11 days old.

Toxicologists usually apply a factor of between 100 and 1000 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. However, given the nature of the findings, HSE took a precautionary approach and applied a factor of 1000. We consider this significant reduction in the uncertainty factor from 1000 to 200 undesirable. The developmental neurotoxicity study in which the effects on the brain measurements were observed reported no behavioural or developmental deficits, and there is an indication that the changes in brain measurement might be reversible. The interpretation of this study by regulatory assessors is uncertain, and, despite the precautionary nature of HSE's assessment, it is not possible to conclude on whether there might be any adverse short term health effects after eating large portions (97.5th percentile consumption) of broccoli containing the level found in this report.

Please refer to the section below on 'Substances that might be genotoxic' for HSE's conclusions regarding potential genotoxicity.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group†	ARfD (mg/kg bw)	Source
Broccoli	Dimethoate and Omethoate	0.04 (D: dimethoate) and 0.01 (O: omethoate)	D: 0.00051	0.00099 (4-6 year old child) 0.00090 (7-10 year old child) 0.00084 (toddler) 0.00082 (infant) 0.00067 (vegetarian) 0.00062 (11-14 year old child) 0.00056 (15-18 year old child) 0.00051 (adult) 0.00051 (elderly – own home) 0.00028 (elderly – residential)	Not established	EU, 2019
			O: 0.00013	0.00025 (4-6 year old child) 0.00022 (7-10 year old child) 0.00021 (toddler) 0.00021 (infant) 0.00017 (vegetarian)	Not established	EU, 2019

				0.00016 (11-14 year old child)		
				0.00014 (15-18 year old child)		
				0.00013 (adult)		
				0.00013 (elderly – own home)		
				0.000069 (elderly – residential)		

Comment on risk assessment

The EFSA Conclusion (2018) for dimethoate has indicated that no toxicological reference values could be determined for dimethoate and its metabolite omethoate, due to a lack of a fully supporting toxicological database. Both dimethoate and omethoate are not approved in the EU and pesticide products containing dimethoate were withdrawn in the EU and UK in 2020.

For dimethoate, EFSA (2018) stated an indicative value for a hypothetical toxicological reference value for short term exposure of 0.0001 mg/kg bw/day. Using this indicative value, estimated dietary intakes of dimethoate for all consumer groups exceeded this reference value. The intakes of omethoate for all consumer groups, except elderly in residential care, exceeded this hypothetical short term toxicological reference value for dimethoate. The highest intake was for 4-6 year old children.

If 4-6 year old children ate large portions of broccoli containing dimethoate at 0.04 mg/kg their intake could be 990 % of the above mentioned hypothetical toxicological reference value . If infants and toddlers ate large portions of broccoli containing omethoate at 0.01 mg/kg their intake could be 250 % of this hypothetical toxicological reference value for dimethoate. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September, 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary

These exposures are undesirable but it is not clear if they may cause any adverse effect. The estimated exposures are not expected to inhibit acetylcholinesterase¹³, the basis of previous evaluations of the safety of dimethoate and omethoate. Based

¹³ this enzyme, acetylcholinesterase, is included in the Glossary on page 123

on this assessment, HSE concludes that a short term effect on health is unlikely after eating large portions (97.5th percentile consumption) of broccoli containing the levels found in this report.

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

Grapes

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group [†]	ARfD (mg/kg bw)	Source
Grapes	Ethephon	0.9	0.018	0.055 (toddler)	0.05	EU, 2008

Comment on risk assessment

If toddlers ate large portions of grapes containing ethephon at 0.9 mg/kg, their intake of ethephon could be 110 % of the Acute Reference Dose. This intake is 109 times lower than a dose which caused no observed adverse effect in a 28 day oral dog study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. However, in this case the factor was larger (120) to ensure consistency with the findings of human volunteer studies. We consider the reduced factor of 109 (from 120) still sufficient to make an effect on health unlikely. More detail on the factors applied is on page 81 of this report.

Based on this assessment, we conclude that a short term effect on health is unlikely.

Rice

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group [†]	ARfD (mg/kg bw)	Source
Rice (Basmati)	Chlorpyrifos	0.005	0.00002	0.000063 (toddler) 0.000055 (7-10 year old child)	No toxicological reference values established	EU, 2019

				0.000055 (4-6 year old child)		
				0.000043 (15-18 year old child)		
				0.000041 (11-14 year old child)		
				0.000038 (vegetarian)		
				0.000030 (adult)		
				0.000028 (infant)		
				0.000020 (elderly – own home)		
				0.0000090 (elderly – residential)		

Comment on risk assessment

EFSA (2019)¹⁴ has indicated that no toxicological reference values could be determined for chlorpyrifos, due to concerns over genotoxicity. Additionally, EFSA raised concerns over neurological effects in the developing foetus and young child. Chlorpyrifos is not approved in the EU and UK and pesticide products containing chlorpyrifos were withdrawn in 2020.

HSE considers that for short-term risk assessment, an indicative toxicological reference value of 0.0003 mg/kg bw can be used based on the LOAEL set by EFSA for a developmental neurotoxicity study and applying a safety factor of 1000 to account for the severe nature of the findings (effects on brain measurements in a developmental neurotoxicity study). Toxicologists usually use safety factors of between 100 and a 1000 when a NOAEL cannot be determined within a study. The HSE proposed indicative toxicological reference value is conservative as it uses the highest uncertainty factor applied by toxicologists and is based on a LOAEL from a study with repeated dosing. This view is further supported by the US EPA ARfD of 0.006 mg/kg bw (published in 2020). Overall, the HSE approach is considered precautionary in protecting the nervous system in the developing foetus and child.

¹⁴ [EFSA 2019 statement on human health- chlorpyrifos](#)

None of the intakes exceeded the HSE proposed indicative toxicological reference value for short term assessment. Based on the low intakes, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day): Adult	Intake (mg/kg bw/day): Critical group[†]	ARfD (mg/kg bw)	Source
Rice (Basmati)	Tricyclazole	0.1	0.00061	0.0013 (toddler) 0.0011 (7-10 year old child) 0.0011 (4-6 year old child) 0.00085 (15-18 year old child) 0.00082 (11-14 year old child) 0.00075 (vegetarian) 0.00061 (adult) 0.00055 (infant) 0.00039 (elderly – own home) 0.00018 (elderly – residential)	No toxicological reference values established	EU, 2016

Comment on risk assessment

A formal risk assessment is not possible as toxicological reference values have not been established in the EU or by other international bodies (JMPR or regulatory authorities). An assessment was performed by EFSA (EFSA Conclusion, 2015¹⁵) prior to the non-renewal of tricyclazole in the EU (2016).

In 2015, EFSA concluded on NOAELs from acceptable toxicological studies, the most critical NOAEL suitable for short term exposure assessment being 5 mg/kg bw/day for maternal toxicity in a rat developmental toxicity. The highest intake for rice in the current assessment for the critical group toddler is more than 3,800 times lower than this NOAEL. In setting an ARfD, toxicologists usually apply a factor of 100 to the NOAEL dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. Therefore, HSE concludes that a short term effect on health is not expected.

Please refer to the section below on 'Substances that might be genotoxic' for HSE's conclusions regarding potential genotoxicity.

¹⁵ EFSA (European Food Safety Authority), 2015. Conclusion on the peer review of the pesticide risk assessment of the active substance tricyclazole. EFSA Journal 2015;13(2):4032, 65 pp. doi:10.2903/j.efsa.2015.4032 Available online: <http://www.efsa.europa.eu/efsajournal>

Short term dietary risk assessment – multiple assessments needed following screening assessment of samples

Samples which contain more than one pesticide from the groups we consider (samples containing more than one organophosphorus/carbamate or captan/folpet or DDAC/BAC or mepiquat/chlormequat or triazoles] and where a more detailed assessment was needed following screening.

Crop/Critical group	Pesticide	Residue mg/kg	Intake		ARfD	Source
			mg/kg bw	%ARfD		
Beans with pods (Guar beans)	Omethoate	0.002	0.000010	- } Total -	Not established 1	EU, 2019 JMPR, 2007
	Profenofos	0.05	0.00025	0.03 }		

Comment on risk assessment:

Omethoate the main metabolite of dimethoate is also assessed in the dimethoate EFSA Conclusion (2018). A toxicological reference value has not been established for either of dimethoate and omethoate, although the EFSA 2018 Conclusion for dimethoate indicated a hypothetical short term reference value of 0.0001 mg/kg bw/day for dimethoate. For further information, please see the above single substance assessments for beans with pods that cover omethoate. The above highest intake of omethoate of 0.000010 mg/kg bw/day is 10% of this hypothetical reference value for dimethoate. Whilst the short term toxicity of omethoate is anticipated to be higher than that for dimethoate, the difference in toxicity would still lead to the dietary intake of omethoate being below this hypothetical reference value for dimethoate. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary. The estimated exposures are unlikely to inhibit acetylcholinesterase¹⁶, the basis of previous evaluations of the safety of dimethoate and omethoate. Whilst profenofos can also affect acetyl cholinesterase at high doses, the highest intake based on the residue found is very low (0.03% of its own ARfD).

Based on the low intakes and this assessment, HSE concludes that a short term effect on health for the combined residues of omethoate and profenofos is not expected.

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

¹⁶ this enzyme, acetylcholinesterase, is included in the Glossary on page 123

Crop/Critical group	Pesticide	Residue mg/kg	Intake			ARfD	Source
			mg/kg bw	%ARfD			
Beans with pods (Guar beans)	Omethoate	0.003	0.000015	- }	Total	Not established	EU, 2019
	Chlorpyrifos	0.01	0.000050	- }	-		

Comment on risk assessment:

Omethoate the main metabolite of dimethoate is also assessed in the dimethoate EFSA Conclusion (2018). A toxicological reference value has not been established for either of dimethoate and omethoate, although the EFSA 2018 Conclusion for dimethoate indicated a hypothetical short term reference value of 0.0001 mg/kg bw/day for dimethoate. For further information, please see the above single substance assessments for beans with pods and broccoli that cover omethoate. The above highest intake of omethoate of 0.000015 mg/kg bw/day is 15% of this hypothetical reference value for dimethoate. Whilst the short term toxicity of omethoate is anticipated to be higher than that for dimethoate, the difference in toxicity would still lead to the dietary intake of omethoate being below this hypothetical reference value for dimethoate. This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September 2019) established an ARfD for dimethoate of 0.02 mg/kg bw; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary. The estimated exposures are unlikely to inhibit acetylcholinesterase¹⁷, the basis of previous evaluations of the safety of dimethoate and omethoate.

Chlorpyrifos can also affect acetyl cholinesterase and this was the basis for the previous ARfD of 0.005 mg/kg bw (EU, 2015). The HSE proposed indicative toxicological reference value for short term assessment (for further information, please see the above single substance assessments for banana, beans with pods, broccoli, and rice that cover chlorpyrifos), was based on a reported changes in in brain measurements in a developmental neurotoxicity study which were not seen in an equivalent study with dimethoate and where exposure to omethoate would be expected. Furthermore, the relationship between brain morphometry changes and acetyl cholinesterase inhibition is unclear. The known common effect from exposure to either omethoate or chlorpyrifos is therefore considered to be on acetyl cholinesterase. For chlorpyrifos, the highest intake based on the residue found in this sample is 0.00005 mg/kg bw, which is 1 % of the old ARfD established based on effects on acetyl cholinesterase.

Overall, HSE concludes that the presence of chlorpyrifos in the sample does not significantly contribute to the overall combined risk when compared to that identified for omethoate in the single substance risk assessments. Based on the low

¹⁷ this enzyme, acetylcholinesterase, is included in the Glossary on page 123

short term intakes, HSE concludes that a short term effect on health for the combined residues of omethoate and chlorpyrifos in this sample of Guar beans (beans with pods) is not expected.

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

Crop/Critical group	Pesticide	Residue mg/kg	Intake			ARfD	Source
			mg/kg bw	%ARfD			
Beans with pods (Guar beans)	Dimethoate	0.04	0.00020	- }	Total -	Not established	EU, 2019
	Omethoate	0.1	0.00050	- }		Not established	EU, 2019
	Monocrotophos	0.005	0.000025	1.3 }	0.002	JMPR, 1995	
	Profenofos	0.02	0.00010	0.01 }	1	JMPR, 2007	

Comment on risk assessment:

The presence of monocrotophos and profenofos in the sample does not significantly contribute to the overall combined intake when compared to dimethoate and omethoate. The estimated short term intakes of dimethoate and omethoate represent 200.5% and 501% of the hypothetical short term reference value for dimethoate proposed by EFSA (EFSA Conclusion for dimethoate, 2018). The overall risk is not expected to be different to the individual risk assessment for this sample presented for dimethoate and omethoate in Guar beans in the table above (see single substance assessment for dimethoate and its metabolite omethoate).

Please refer to the section below on ‘Substances that might be genotoxic’ for HSE’s conclusions regarding potential genotoxicity.

Crop/Critical group	Pesticide	Residue mg/kg	Intake			ARfD	Source
			mg/kg bw	%ARfD			
Broccoli	Chlorpyrifos	0.06	0.0015	- }	Total -	Not established	EU, 2019
	Dimethoate	0.04	0.00099	- }		Not established	EU, 2019
	Omethoate	0.01	0.00025	- }		Not established	EU, 2019

Comment on risk assessment:

If 4-6 year old children ate large portions of broccoli containing dimethoate at 0.04 mg/kg and omethoate at 0.01 mg/kg bw their intake would exceed the hypothetical toxicological reference value for short term exposure to dimethoate (see single substance risk assessments for beans with pods). This indicative toxicological reference value is a precautionary value intended to protect the nervous system in the developing foetus and child, which has been set well below intakes which caused no observed effects in animal studies. The JMPR (September, 2019) established an ARfD for dimethoate of 0.02 mg/kg bw based on; this supports the view that the proposed hypothetical reference value from the EFSA Conclusion is precautionary. The estimated exposures are unlikely to inhibit acetylcholinesterase, the basis of previous evaluations of the safety of dimethoate and omethoate.

Chlorpyrifos can also affect acetyl cholinesterase and this was the basis for the previous ARfD of 0.005 mg/kg bw (EU, 2015). The proposed indicative toxicological reference value for short term assessment (for further information, please see the above single substance assessments for banana, beans with pods, broccoli, and rice that cover chlorpyrifos), was based on reported changes in brain measurements in a developmental neurotoxicity study which were not seen in an equivalent study with dimethoate and where exposure to omethoate would be expected. Furthermore, the relationship between the reported brain morphometry changes and acetyl cholinesterase inhibition is unclear. The known common effect from exposure to either dimethoate, omethoate or chlorpyrifos is therefore considered to be inhibition of acetyl cholinesterase. For chlorpyrifos, the intake based on the residue found in this sample is below its old ARfD (EU, 2015) established based on effects on acetyl cholinesterase. Overall, HSE concludes that the presence of chlorpyrifos in the sample does not significantly contribute to the overall combined risk of an effect on acetyl cholinesterase when compared to that identified for dimethoate and omethoate in the single substance risk assessments. However, the potential adverse effects from exposure to chlorpyrifos on its own are discussed in the above single substancerisk assessment for the finding of chlorpyrifos in this broccoli sample and therefore it is not possible to conclude on whether there might be any adverse short term health effects after eating large portions (97.5th percentile consumption) of broccoli containing the highest levels found in this report.

Please refer to the section below on 'Substances that might be genotoxic' for HSE's conclusions regarding potential genotoxicity.

Long term dietary risk assessments needed following screening assessment of samples

As noted in [section 3](#) total long term dietary assessments across all commodities are not performed for these quarterly assessments. The issue is more fully considered in regulatory contexts pre-authorisation and at the time of MRL review. Then the issue is considered across all commodities (so more precautionary) by pesticide levels determined in GAP compliant trials, intended to address highest likely residues that might arise following pesticide use according to label recommendations

However, for the PRiF quarterly assessments, HSE do perform a screening exercise for all of the residues found for an individual commodity to see if the long term intakes (commodity by commodity) show any indication of exceedance of the ADI. If an exceedance was observed then HSE would consider further and we would present a more detailed risk assessment.

None of these individual commodity long term exposure screening assessments performed in this quarter (for each of the pesticides found in this report) indicated potential for adverse long term health effects. HSE assessed the dietary intakes to be below the ADI or other established long term health based reference value.

Substances that might be genotoxic (see explanation in the section on HSE's assessment of risk)

During regulatory assessment, careful consideration is given to any pesticides that may exhibit any potential to be genotoxic (able to damage genetic material) in live animals, so we need to consider the significance to the consumer when these residues are found. There are small number of examples of older pesticides that might be genotoxic, where modern data to investigate the true genotoxic potential is not expected to be made available. It is likely that these will only be found in imported foods. For many of these old pesticides, the toxicological reference doses are low and PRiF uses low reporting limits to ensure that these residues are found even at very low levels, as we know they are of particular interest to consumers. The evaluation of possible health implications for PRiF findings is complex as tests for genotoxicity are commonly performed at higher doses (orders of magnitude higher) than the dietary exposure levels that are assessed in PRiF reports. As such it is difficult to conclude specifically, and to extrapolate the findings in the laboratory to the context of findings in the PRiF monitoring and the presence of residues at low levels in foods. Where relevant some reassurance that any risks are likely to be small can be gained if increased cancer incidence, which may be due to gene mutations, does not occur in long term animal feeding studies, designed to detect such observations. Where relevant we will indicate this. Due to the uncertainty about the potential for genetic damage (genotoxicity) at low doses, PRiF will always conclude that on a precautionary basis any findings of genotoxic substances in food are undesirable.

Assessment of genotoxicity (Quarter 1 2021) and conclusions:

Residues found in this report that have genotoxic potential (concluded from laboratory studies on animals):
monocrotophos and omethoate

There is some evidence (*in vitro* and/or *in vivo*^[1]) that these pesticides can damage genetic material (are genotoxic). There is some reassurance that risks of developing ill health effects following single or repeat exposures are likely to be low, since they did not increase cancer incidence in studies with repeat daily doses over their life-span in rats or mice. The doses used in both the genotoxicity tests and the cancer studies were orders of magnitude higher than the exposures estimated in this assessment. It is not known if lower doses which are not toxic also have this effect.

Residues found in this report where toxicological data are suggestive of genotoxicity but not certain: dimethoate, chlorpyrifos, tricyclazole, and carbofuran

It is unclear whether these pesticides can damage genetic material (are genotoxic). There is some evidence from studies performed *in vitro* and/or *in vivo* that they may be genotoxic. Whilst there are negative results in the available *in vivo* studies, the currently recommended *in vivo* follow up studies, that may clarify the genotoxic potential of these pesticides, have not been performed. There is some reassurance that risks of developing ill health effects following single or repeat exposures are likely to be low, since they did not cause cancer in this or other long term studies with repeat daily doses in animals over their life-span. The doses used in these studies were orders of magnitude higher than the exposures estimated in this assessment. It is not known if lower doses which are not toxic also have this effect.

Conclusions: Overall, we conclude that on a precautionary basis any of these residue findings of these pesticides are undesirable due to the uncertainty regarding genotoxicity at low doses; however, we consider any risks of adverse health effects are low at the highest levels of exposure after eating large portions (97.5th percentile consumption) of the foods containing the levels of these pesticides found in this report.

^[1] *In vivo/in vitro* : see glossary

Section 4: issues arising in this report and updates on previous reports

Issues arising in this report

Chlorate

We have been testing a limited number of foods for chlorate since 2016. The pesticide sodium chlorate is a residual broad action weed killer that is not authorised for use in the EU. However, we are confident that the residues we are detecting come from use of chlorine-based disinfectants used to maintain microbiological safety (control microorganisms that cause food poisoning). Because these residues are unavoidable, and important for the maintaining of microbiological control vital for food safety, we are not treating these results as breaches of the MRL. **We are not advising that food companies change their existing practices as a result of our findings, but they should be aware about the ongoing discussion in this area.**

We are only part of the work going on across government and beyond to consider what to do about chlorate residues in food and water.

MRLs after 20 June 2020

After detailed discussion and consultation with stakeholders the EU agreed new MRLs for chlorate that came into force on 20 June 2020. All samples covered by this report were taken after 20 June 2020 when the new, higher MRLs were in place.

The new chlorate MRLs include a footnote referring specifically to taking account of the use of biocides during processing in addition to the MRLs for food as harvested or initially produced. The footnote exceptionally specifies that for considering compliance with chlorate MRLs, simple types of processing that do not affect the other residue levels, such as packing, washing, chopping and freezing can be taken into account.

The responsibility for providing evidence showing that residues from processing can be taken into account, lies with the food business operator, and so we will be interested to see such evidence where appropriate HSE will decide whether the footnote can be applied and if so this will be reflected in our reports.

The Food and Biocides Industry Group have produced more detailed information and guidance on this topic which is available on the Chilled Food Association's website at <https://www.chilledfood.org/fbig/>.

Best practice for use

The Food Standards Agency is working with the food industry to develop and promote best practice in the use of sanitisers. This is important because the presence of low-level residues of chlorate in food results from measures taken by the food and water industries to protect food safety by reducing microbiological contamination of food and drink (including drinking water, which is a significant source of chlorate in food). Chlorate itself is not used as a disinfectant, but chlorine-based sanitisers can contain small amounts of chlorate.

Drinking Water

Defra is also working on the EU recast of its Drinking Water Directive. Discussions are underway about the possible future monitoring of chlorate and the level to be achieved. In national legislation throughout the UK it is already a requirement to keep disinfection by-products as low as possible. This is usually achieved through management of disinfectant dosing and storage.

Advisory Committee on the Microbiological Safety of Food

Microbiological safety of food

The HSE are working with the Advisory Committee on the Microbiological Safety of Food to understand how changes to pesticide MRLs affect biocide use, microbiological food safety, and any change to the overall risk to consumers taking into account both chemical and microbiological safety.

Dietary intakes

Since 2018 the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) has been considering chlorate as part of its on-going work looking at the chemicals in the diet of infants and young children (up to 5 years). The European Food Safety Authority's 2015 opinion on chlorate¹⁸ establishes appropriate health-based guidance values for chlorate exposure to protect against acute and chronic risks to health.

DDT

The use of DDT is banned or heavily restricted in many countries. It isn't allowed for use on food crops anymore, but it is still used in some countries outside the EU as a public health insecticide. Residues of DDT take a long time to break down in the environment and can accumulate in fatty tissue which is a major reason that it has been banned in the EU and many other countries.

Due to the bans and restrictions on use, the levels in food have decreased substantially since the 1960s and 1970s. Even so, because it takes a long time to break down we do expect, and do see, occasional DDT residues in our monitoring results. Overall, the incidence and the size of residues have fallen steadily over time, which is what we would expect. In recent years none of our findings were unusual, unexpected or of concern. We can tell from the chemical form that we detect whether the residues we have found are from historic use (which is what we usually find). We explain this every time we publish DDT results to try to make it as clear as we can that the results show food producers are not using DDT today. However, there are occasional media stories about DDT and various links and associations, which do not make this distinction.

The residues we find nowadays are at levels that would not be expected to have any effect on health, either in the short term or in the long term, when checked against today's understanding of the effect of DDT on health. As a committee, we take care to ensure we look thoroughly at this, and the Food Standards Agency is also actively involved in our considerations.

¹⁸ [EFSA Journal 2015;13\(6\):4135 \[103 pp.\]](http://efsa.europa.eu/journal/2015/13(6):4135)
http://ec.europa.eu/food/plant/standing_committees/sc_phytopharmaceuticals/index_en.htm

Follow-up from Previous Reports

Quarter 3 2020

Beans with pods

Fluazifop-P: Sample number 4325/2020

We passed details of a sample of bobi beans from the UK that contained fluazifop-p (partial sum) to HSE. HSE have accessed the spray records and there was no evidence of use on the bobi beans or any neighbouring fields and the case was closed with no further action taken.

The brand name details have been included in this report.

Quarter 4 2020

Cauliflower

Fonicamid (sum): Sample number 4634/2020

We passed details of a sample of cauliflower from the UK that contained fonicamid (sum) to HSE. HSE enquiries are not yet complete and an update will appear in a future report.

Brand name details of samples where follow-up action is now complete

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)
4325/2020	11/08/2020	Bobo Beans	UK	S Thorogood & Sons	Stand 87-88A New Spitalfields Market, 17 Sherrin Road, London E10 5SQ	None stated	Wye Valley Produce Cobrey Farms, Coleraine Buildings, Coughton, Ross-on-Wye, Herefordshire HR9 5SG	tebuconazole 0.01 (MRL = 2) fluazifop-p (partial sum) 0.02 (MRL = 1.5)

In our next report:

In Quarter 2 of 2021 we will look at results for:

Samples collected in GB

- Asparagus
- Aubergine
- Banana
- Beans with pods
- Beef
- Berries and small fruits
- Broccoli
- Cheese (soft)
- Eggs
- Grapefruit
- Grapes
- Melon
- Milk
- Mushrooms
- Olive oil
- Pepper
- Potatoes
- Raspberry
- Spring greens and kale
- Wheat

Samples collected in NI

- Asparagus
- Aubergine
- Banana
- Beans with pods
- Beef
- Berries and small fruits
- Broccoli
- Cheese (soft)
- Eggs
- Fish (white)
- Grapefruit
- Grapes
- Melon
- Milk
- Mushrooms
- Olive oil
- Pepper
- Potatoes
- Raspberry
- Spring greens and kale
- Wheat

Section 5: background and reference

Reasons for pesticide residue testing

Food safety is important. Modern food production processes have given us plentiful supplies of a wide range of good quality affordable produce.

In the food industry of today the production environment can be managed from the preparation of seeds used for crops, through to growth, harvesting and storage of the produce.

One of the ways the food industry controls the environment in which foodstuffs are produced is by applying pesticides. They help farmers and growers maximise the production of food stuffs by, for example, preventing weeds inhibiting the growth of the crop, or insects destroying or infesting them. Pesticides can also be used to help protect seeds or prolong the life of crops after they have been harvested. Biological and physical (cultural) controls are also used to protect crops or as part of an integrated system.

As pesticides are used to control unwanted pests, weeds and diseases, they can potentially also harm people, wildlife and the environment. This is why the UK, in common with most other countries, imposes legally enforceable conditions as to how and when pesticides can be used. No pesticide can be supplied or used on a food or ornamental crops in the UK without Government authorisation. To obtain this authorisation the manufacturer of the pesticide must show that it does not present a concern for people's health or the environment. Naturally derived and synthetic pesticides are subject to the same regulation.

Once the authorisation has been granted Government authorities carry out follow up checks to ensure that the authorisation is providing the necessary degree of protection to users, consumers and the environment and that those who use pesticides are complying with conditions specified within it.

The Government authority responsible for checking pesticide residues in foodstuffs is the Health and Safety Executive. Defra's Expert Committee on Pesticide Residues in Food (PRiF) oversees and provides an independent check on this work. We know that the use of pesticides on crops may lead to traces (residues) of these chemicals in food and we expect to find these in our monitoring programme.

Detail of reporting practice

Results by food commodity

- We include information about the survey (for instance where samples came from) for each commodity
- Detailed tabulated results are at the back of this report - these tables are also available for download from our website
- We summarise our findings and any follow-up action taken.

Risk assessments – single residues

- All results are screened by HSE to check for intakes above the toxicological reference values, the Acute Reference Dose (ARfD) or the Acceptable Daily Intake (ADI). HSE assumes a relatively high level of intake and also assumes that most produce is eaten whole including peel/skin even when these are rarely consumed

- Where intakes above the toxicological reference values are identified, we consider a detailed risk assessment prepared by HSE (at section 3 of this report).
- Our observations and the follow-up action taken are summarised in the section for that food.

Risk assessments – multiple combined residues

- Residues of more than one pesticide from the same category/class of particular categories of pesticides, which have a similar toxicological mode of action, are initially screened by HSE to check for intakes that might need further combined assessment.
- Where combined intakes exceed the initial screen 'trigger', we consider a detailed combined risk assessment prepared by HSE (at section 3 of this report). Further details on the approach are explained in Section 3.
- Our observations and any follow-up action taken are summarised in the section for that food commodity.

Risk assessment - conclusions

- Where, in the light of current knowledge and considering the usual level of scientific uncertainty the intake will not cause ill health the conclusion will say no effect on health is expected.
- Where, in the light of current knowledge the intake is not likely to cause ill health, the conclusion will be less definite and state that an effect on health is unlikely.
- Where scientific uncertainty is greater or if risk of adverse health effects could be higher more information is provided.

Residues in GB and NI produce of pesticides which do not have a PPP authorised for use on that crop in GB and NI.

- All residues found in foods produced in GB or NI are checked by HSE to make sure there is a PPP containing that pesticide authorised for use on that crop.
- Where there is no GB or NI authorisation is identified, details of the sample are referred to the Enforcement Section for follow up.
- Our observations and any follow-up action taken to date are summarised in the section for that food commodity. We may have to withhold details of samples while investigations are underway, in which case the details will be published in a later report.

Residues above the MRL, after taking into account measurement uncertainty

- Samples containing residues above the MRL are listed at Appendix B, and those which are clearly above the MRL after taking into account measurement uncertainty of plus or minus 50% are highlighted.
- Our observations and any follow-up action taken are summarised in the section for that food commodity.
- The results in our reports are rounded for publication but not adjusted for measurement uncertainty.
- We apply measurement uncertainty only to decide whether to highlight a result as over the MRL in the brand name annex. To do this we use the actual value reported by the laboratory before rounding. If after taking measurement uncertainty into account that value is found to be over the MRL the result will be highlighted in the brand name annex.

For example:

- The lab reports the results of duplicate analysis of a residue above an MRL at 0.023 mg/kg and 0.025 mg/kg giving an average value of 0.024mg/kg. For reporting purpose this value would be 0.02 mg/kg.

- If measurement uncertainty is then applied to the reported value of 0.02 mg/kg it could take the value to between 0.01 - 0.03 mg/kg. If the MRL is 0.01 mg/kg the lower value would be at the MRL and there is no exceedance.
- However, if measurement uncertainty is applied to the measured result, e.g. 0.024 mg/kg the value could then be in the range of 0.012 – 0.036 mg/kg. In this case the lower value is above the MRL and so will be treated as an exceedance.

Residues in organic food

- We monitor pesticide residues in all the GB and NI food supply, including organic food.
- We are not responsible for checking compliance with the rules associated with organic production. However, when we do detect residues in an organic food we explain whether or not those residues indicate a breach of the rules and inform Defra's Organic Farming Branch.

Brand Name Annex

- Full brand name details for samples included in this report are published in a brand name annex. Within this annex, samples with results of interest are highlighted.
- Brand name details are only published when enough follow-up work is completed for us to be reasonably sure whether a breach of the law or good practice has occurred.
- Therefore, sometimes brand name details are withheld pending completion of this work but are published in a later report.

Pesticides analysed as multi-component analytes and their reporting limits

Why some results cover more than one substance

Both the legal controls and our analytical tests are aimed at checking food for the presence of residues of specific pesticides. Residues are the chemical traces left behind after pesticides are used. In most cases the residue of a pesticide is measured by first identifying the pesticide and then measuring the quantity of that pesticide in the food itself. But for some pesticides the residue remaining in the food is known to be chemically different from the original pesticide and so the laboratory needs to look for more than one component. There are various reasons why this happens, for example:

- the animal or plant can change the pesticide into related chemicals
- the pesticide can change in the environment into related chemicals
- some pesticides are mixtures of chemicals, so the relevant components of the mixture need to be checked for
- in the laboratory sample preparation and/or analysis may change pesticides into related chemicals
- related chemicals may be pesticides in their own right

The MRL setting process takes account of all these issues. The EU may set a complex residue definition to ensure that the identity and quantity of the residue found is representative of the pesticide present. A complex residue definition may be set where it is necessary for safety reasons or to be able to accurately identify the pesticide residue present in the food. This definition usually includes the actual pesticide, plus other related chemicals. These residues are usually reported together as a "sum". Sometimes different foods need different definitions because different pesticide residues are known to occur in that food. For instance, plants and animals may metabolise a pesticide differently, which forms different residues.

The full definitions of pesticides that we have found in our surveys are described in the table below. If you would like more detail about a particular residue definition, please get in touch. You can email us at prif@hse.gov.uk and other contact details are on the back cover.

Where the detailed individual analysis results tell us something useful, we mention that in our conclusions.

How we calculate sums

Unless the definition says otherwise, the summed result is a simple addition. For individual components that are not detected that result is treated as a zero.

Where a residue definition says, “expressed as”, that means that the individual component results are adjusted by molecular weight before being added together. The residue definition is set this way so that the final calculated result for the whole definition is an expression of the level of the most toxic component, and so that value can be used directly in consumer risk assessment without further adjustment.

Complex residue definitions used in our reports

There are a large number of pesticides used and types of food in the world. So other complex residue definitions may apply to food/pesticide combinations not yet considered by PRiF. You can look up all the EU MRL definitions for pesticide residues at the European Commission’s pesticide database at [EU-Pesticide Database](#)

Short name we use in our reports	Legal residue definition – These definitions apply to all foods unless otherwise stated
2,4-D (sum)	2,4-D (sum of 2,4-D and its esters expressed as 2,4-D)
abamectin (sum)	Abamectin (sum of Avermectin B1a, AvermectinB1b and delta-8,9 isomer of Avermectin B1a)
aldicarb (sum)	Aldicarb (sum of Aldicarb, its sulfoxide and its sulfone, expressed as Aldicarb)
aldrin and dieldrin	Aldrin and Dieldrin (Aldrin and dieldrin combined expressed as dieldrin), aka dieldrin (sum)
Amitraz	Amitraz (amitraz including the metabolites containing the 2,4 - dimethylaniline moiety expressed as amitraz)
BAC (sum)	Benzalkonium chloride (mixture of alkylbenzyltrimethylammonium chlorides with alkyl chain lengths of C ₈ , C ₁₀ , C ₁₂ , C ₁₄ , C ₁₆ and C ₁₈)
benthiavalicarb (sum)	Benthiavalicarb (Benthiavalicarb-isopropyl (KIF-230 R-L) and its enantiomer (KIF-230 S-D) and diastereomers (KIF-230 R-L and KIF-230 S-D))
bixan (animal products)	Sum of bixafen and desmethyl bixafen expressed as bixafen This definition applies to animal products only
captan and folpet	Sum of captan and folpet aka captan/folpet This definition applies only to pome fruit (fruits such as apples and pears), strawberries, raspberries, currants, tomatoes and beans. For all other foods there are separate MRLs for captan only and for folpet only.

carbendazim (animal products)	Carbendazim and thiophanate-methyl, expressed as carbendazim
Carbendazim (sum)	Carbendazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim)
carbofuran (sum)	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)
chlordan (animal products)	Chlordane (sum of cis- and trans-isomers and oxychlordane expressed as chlordane) This definition applies to animal products only
chlordan (sum)	Chlordane (sum of cis- and trans- isomers) This definition applies to all foods except animal products
chlorpropham (potatoes)	Chlorpropham only This definition applies only to potatoes
chlorpropham (sum for animal products)	Chlorpropham and 4-hydroxychlorpropham-O-sulphonic acid (4-HSA), expressed as chlorpropham This definition applies only to animal products
chlorpropham (sum)	Chlorpropham (Chlorpropham and 3-chloroaniline, expressed as Chlorpropham) This definition applies to all foods except potatoes and animal products
DDAC (sum)	Didecyldimethylammonium chloride (mixture of alkyl-quaternary ammonium salts with alkyl chain lengths of C ₈ , C ₁₀ and C ₁₂)
DDT (sum)	DDT (sum of p,p'-DDT, o,p'-DDT, p-p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Dichlorprop	Sum of Dichlorprop, including dichlorprop-p and its conjugates, expressed as dichlorprop
dicofol (sum)	Dicofol (sum of p, p' and o,p' isomers)
Dimethenamid	Dimethenamid-p (Dimethenamid-p including other mixtures of constituent isomers (sum of isomers))
dimethoate (sum)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
disulfoton (sum)	Disulfoton (sum of disulfoton, disulfoton sulfoxide and disulfoton sulfone expressed as disulfoton)

dithiocarbamates	Dithiocarbamates are a group of pesticides that are chemically similar. Testing for them individually in routine analysis is not possible, so MRLs are set for a test for the group.
endosulfan (sum)	Endosulfan (sum of alpha- and beta-isomers and endosulfan-sulphate expressed as endosulfan)
fenamiphos (sum)	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
fenchlorphos (sum)	Fenchlorphos (sum of fenchlorphos and fenchlorphos oxon expressed as fenchlorphos)
fensulfothion (sum)	Fensulfothion (sum of fensulfothion, its oxygen analogue and their sulfones, expressed as fensulfothion).
fenthion (sum)	Fenthion (fenthion and its oxygen analogue, their sulfoxides and sulfone expressed as parent)
fenvalerate & esfenvalerate (all isomers)	Fenvalerate (any ratio of constituent isomers (RR, SS, RS & SR) including esfenvalerate)
fipronil (infant food)	Sum of fipronil and fipronil-desulfinyl, expressed as fipronil This definition applies to foods for babies only
fipronil (sum)	Fipronil (sum Fipronil and sulfone metabolite (MB46136) expressed as Fipronil) This definition applies to all foods except foods for babies
flonicamid (sum)	Flonicamid (sum of flonicamid, TNFG and TNFA) This definition applies to all food except animal products. The full definition must be sought. Residues found are usually of the metabolites.
fluazifop-p-butyl (sum)	Fluazifop-P-butyl (fluazifop acid (free and conjugate))
Fosetyl (sum)	Fosetyl-Al (sum of fosetyl, phosphonic acid and their salts, expressed as fosetyl)
haloxyfop (sum)	Haloxyfop including haloxyfop-R (Haloxyfop-R methyl ester, haloxyfop-R and conjugates of haloxyfop-R expressed as haloxyfop-R)
Heptachlor (infant food)	Sum of heptachlor and trans heptachlor epoxide This definition applies to foods for babies only
Heptachlor (sum)	Heptachlor (sum of heptachlor and heptachlor epoxide expressed as heptachlor) This definition applies to all foods except infant foods

hexachlorocyclohexane (sum)	Hexachlorocyclohexane (HCH), sum of isomers, except the gamma isomer This definition applies to all foods except animal products (For animal products the alpha and beta isomers have separate MRLs)
Malathion	Malathion (sum of malathion and malaoxon expressed as malathion)
MCPA (animal products)	[Residue definition, animal products] MCPA, MCPB and MCPA thioethyl expressed as MCPA This definition applies to animal products only
MCPA (sum)	MCPA and MCPB (MCPA, MCPB including their salts, esters and conjugates expressed as MCPA) This definition applies to all foods except animal products
Mepanipyrim (sum)	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine) expressed as mepanipyrim
methiocarb (sum)	Methiocarb (sum of methiocarb and methiocarb sulfoxide and sulfone, expressed as methiocarb)
methomyl (sum)	Sum of methomyl and thiodicarb expressed as methomyl
Oxydemeton-methyl (sum)	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)
parathion-methyl (sum)	Parathion-methyl (sum of Parathion-methyl and paraoxon-methyl expressed as Parathion-methyl)
Permethrin	Permethrin (sum of isomers)
phorate (sum)	Phorate (sum of phorate, its oxygen analogue and their sulfones expressed as phorate)
phosmet (sum)	Phosmet (phosmet and phosmet oxon expressed as phosmet) This definition applies to all foods except animal products
pirimicarb (sum)	Pirimicarb (sum of Pirimicarb and Desmethyl pirimicarb expressed as Pirimicarb) for certain animal products. Pirimicarb only for fruit and vegetables and some animal products.
Prothioconazole (sum)	Prothioconazole (sum of prothioconazole-desthio and its glucuronide conjugate, expressed as prothioconazoledesthio) This definition applies to animal products only
PTU & propineb	Sum of PTU and propineb This definition applies to food for babies only

quintozene (sum)	Quintozene (sum of quintozene and pentachloro-aniline expressed as quintozene)
Prochloraz (sum)	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)
Terbufos (sum)	Terbufos (sum of terbufos, its sulfoxide and sulfone This definition applies only to foods for babies)
thiamethoxam (sum)	Thiamethoxam (sum of thiamethoxam and clothianidin expressed as thiamethoxam) There are <u>also</u> separate clothianidin MRLs
tolyfluanid (sum)	Tolyfluanid (Sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)
triadimefon & triadimenol	Triadimefon and triademenol
vinclozolin (animal products)	Vinclozolin, iprodione, procymidone, sum of compounds and all metabolites containing the 3,5-dichloroaniline moiety expressed as 3,5-dichloroaniline This definition applies to animal products only
vinclozolin (sum)	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5-dichloroaniline moiety, expressed as vinclozolin) This definition applies to all foods except animal products

Glossary

This is a 'standard' glossary which defines the key terms used in the PRiF reports. Not all the terms listed here are used in this particular report.

97.5th percentile consumer: Please refer to glossary entry for 'High level consumer'.

Acceptable Daily Intake (ADI): This is the amount of a chemical which can be consumed every day for a lifetime in the practical certainty, on the basis of all known facts, that no harm will result. It is expressed in milligrams of the chemical per kilogram of body weight of the consumer. The starting point for the derivation of the ADI is usually the 'no observed adverse effect level' (NOAEL) that has been observed in animal studies for toxicity. This is then divided by an uncertainty factor (most often 100) to allow for the possibility that animals may be less sensitive than humans and also to account for possible variation in sensitivity between individuals. The studies from which NOAELs and hence ADIs are derived take into account any impurities in the pesticide active substance as manufactured, and also any toxic breakdown products of the pesticide.

Acetylcholine: Acetylcholine is a neurotransmitter, a chemical that carries signals through the nervous system. See *cholinergic*

Acetylcholinesterase: This is an enzyme which degrades acetylcholine and is involved in the regulation of nerve impulses. Inhibition of this enzyme can interfere with this nerve transmission function. This is a short-term effect of concern with organophosphate and carbamate pesticides at levels above the ARfD.

Acute Reference Dose (ARfD): The definition of the ARfD is similar to that of the ADI, but it relates to the amount of a chemical that can be taken in at one meal or on one day without appreciable health risk to the consumer. It is normally derived by applying an appropriate uncertainty factor to the lowest NOAEL in studies that assess acute toxicity or developmental toxicity.

As a matter of policy, the EU does not use NOAELs from tests that involve deliberate administration of pesticides to humans to determine ADIs and ARfDs. However, where such data have been ethically and scientifically derived some authorities, e.g. the World Health Organization, do consider such data. Where human data are used there is usually less uncertainty in the resulting reference value compared to extrapolating from animal tests to humans, and a lower uncertainty factor (most often 10) is used to account for the variation in sensitivity between individuals.

The initial risk assessments in PRiF reports use the agreed EU reference values. However, where intakes are above the EU value and a reference value based on acceptable human data is available a refined assessment, which is a more appropriate indicator of the risk, is also reported.

Analyte: This is the name for the substance that the PRiF surveys look for and measure if present; it could be a pesticide itself or a product from a pesticide when it is degraded, or metabolised.

COLEACP (Europe-Africa-Caribbean-Pacific Liaison Committee): It aims to promote the competitive export of fresh fruit, vegetables, flowers and ornamental plants from the ACP. Its specialised information and advisory services are open to all ACP companies in the horticultural

export sector and are financed by the European Commission. It has two overriding objectives to enable ACP companies to comply with European food safety and traceability requirements and to consolidate the position of small-scale producers in the ACP horticultural export sector.

Cholinergic: In relation to the animal nervous system, processes and structures are cholinergic if they release or use acetylcholine.

Cryogenic Milling: Processing of commodities at very low temperatures can be achieved by milling/grinding pre-frozen samples in the presence of dry ice, a procedure known as 'cryogenic milling'.

Extensions of Authorisations for Minor Use (EAMUs): Users and authorisation holders of agricultural Plant Protection Products (PPP) may apply to have the authorisation of specific PPP's extended to cover uses additional to those authorised and shown on the manufacturer's product label. For many reasons, label recommendations of authorised pesticides do not cover the control of every problem which may arise. This is particularly true for crops that are grown on a comparatively small scale in the UK as well as for pests and diseases that occur less often or which are new to the UK. As part of the process evidence on residues that would arise from the use is required, and consumer safety is evaluated and if necessary a specific MRL set. EAMU is pronounced "emu" these types of authorisations are also informally called "off labels".

Genotoxicity: Genotoxicity is the effect of substances (called genotoxins) which can alter or damage the genetic material (DNA, RNA or chromosomes) within a cell. Cells have the capacity to protect themselves from genotoxic effects by many repair processes and therefore many genotoxic events do not become evident as mutations. Where mutations occur, this can lead to cancer or effects that can be passed to unborn children (e.g. birth defects, inherited diseases).

Good Agricultural Practice in the Use of Pesticides (GAP): The nationally authorised safe uses of pesticides under conditions necessary for effective and reliable pest control (the way products should be used according to the statutory conditions of authorisation which are stated on the label). GAP encompasses a range of pesticide applications up to the highest authorised rates of use, applied in a manner which leaves a residue which is the smallest practicable. Authorised safe uses are determined at the national level and include nationally registered recommended uses, which take into account public and occupational health and environmental safety considerations. Actual conditions include any stage in the production, storage, transport, distribution and processing of food commodities and animal feed.

High-level Consumer: A term used in UK risk assessment calculations to describe the amount of food consumed by a person. In line with internationally agreed approaches, the PRiF uses the 97.5th percentile value, which is generally about three times the average amount consumed. This takes account of different eating patterns that may occur throughout the population.

Human Data: See under Acute Reference Dose

In vitro: a test performed *in vitro* "in the glass" means that it is performed outside of a living organism and usually involves isolated tissues, organs or cells.

In vivo: live animal studies

Import Tolerance: an MRL set for imported products where the use of the active substance in a plant protection product on a commodity is not authorised in the European Community (EC) or an existing EC MRL is not sufficient to meet the needs of international trade. All import tolerances are assessed for consumer safety.

Imported: The tables in the reports record whether the sample was of UK origin, or imported. This can mean different things depending on the commodity. See also 'Origin'. The PRiF report the country from where the produce has been imported only if this is clear from the packaging or labelling.

JMPR: Joint FAO/WHO Meeting on Pesticide Residues, which conducts scientific evaluations of pesticide residues in food.

LOD (Limit of Determination) and LOD MRLs: The Limit of Determination (LOD) is the lowest concentration of a pesticide residue or contaminant that can be routinely identified and quantitatively measured in a specified food, agricultural commodity or animal feed with an acceptable degree of certainty by the method of analysis.

LOD MRL (Maximum Residue Levels set at the LOD): These are marked by a '*'. For some pesticides and commodities insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).** Also, see under Reporting limit.

Lowest Observed Adverse Effect Level (LOAEL): The lowest concentration or amount of a substance, found by experiment or observation, which causes detectable adverse alteration of morphology, functional capacity, growth, development or life span of the target organism under defined conditions of exposure.

Off Label: See Extensions of Authorisations for Minor Use (EAMUs)

Maximum Residue Level (MRL): The maximum concentration of a pesticide residue (expressed as mg/kg) legally permitted in or on food commodities and animal feeds. MRLs are based on good agricultural practice data and residues in foods derived from commodities that comply with the respective MRLs are intended to be toxicologically acceptable.

MRLs are intended primarily as a check that GAP is being followed and to assist international trade in produce treated with pesticides. **MRLs are not in themselves 'safety limits'**, and exposure to residues in excess of an MRL does not automatically imply a hazard to health.

The MRLs applicable in the UK are now largely set under EC legislation.

Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in produce, which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not authorised for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory analysis. Thus, they provide a mechanism for statutory controls on pesticides in produce which is put into circulation and for monitoring correct use of these chemicals.

If no use of a pesticide on a crop is identified when MRLs are set the tolerance for that pesticide/crop combination is set at the limit of determination (effectively zero). Limit of determination MRL are marked by a '*'.

MRLs are established under the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuff) (England and Wales) Regulations 1999 (as amended), the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuff) (Scotland) Regulations 2000 and the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuff) Regulations (Northern Ireland) 2002. These Regulations list all statutory MRLs established under UK national or EC

procedures. Today, virtually all these MRLs are set under an ongoing EC programme and the Regulations are amended periodically as levels are set for increasing numbers of pesticides.

There are a number of pesticides which do not yet have statutory MRLs. In the absence of such MRLs we advise suppliers to adhere to any appropriate levels established by the Codex Alimentarius Commission (CAC) a United Nations body established to promote global trading standards. Codex MRLs are not statutory but have been risk-assessed when set and provide a suitable standard in the absence of a statutory MRL.

MRLs may be extended to composite and processed products but levels are not specifically laid down in legislation. They are derived by calculation on an individual basis.

Maximum Residue Levels set at the LOD (LOD MRL): See LOD MRL. For some pesticides and commodities, insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level, i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).**

MRL exceedances: When a residue is found at a level higher than that set for the MRL.

MRL Exceedances and Relationship with the Acceptable Daily Intake (ADI): Before permitting any use of a pesticide, a detailed assessment is made to ensure that residues in foods derived from commodities comply with MRLs and will not give rise to unacceptable risks to consumers. MRLs do take account of consumer safety aspects and, in effect, are set at levels below safety limits. However, MRLs must not be confused with safety limits, which are expressed in terms of the acceptable daily intake (ADI) of a particular pesticide residue from all sources. The ADI (expressed as mg/kg bw/day) is the amount of chemical that can be consumed every day of an individual's entire lifetime in the practical certainty, on the basis of all known facts, that no harm will result. See ADI for further information.

Whenever unexpectedly high or unusual residues occur during monitoring, the risk to consumers, from exposure to residues at the highest levels found, is assessed by comparison of predicted intakes with the ADI or ARfD as appropriate.

No MRL: For certain pesticides an MRL may not have been set.

Metabolite: A degradation or conversion product from a pesticide when it is metabolised.

Multiple Residues: In this report this term is used to describe when more than one pesticide is found in an individual food sample. It may have arisen because the crop was treated at different times with pesticides applied singularly, or when pesticides are applied as mixtures (several pesticides mixed in the spray tank at the same time) or the marketed pesticide product contains more than one pesticide or any combination of these three situations. Mixtures may be used in response to specific pest pressures and also as part of strategies to minimise pesticide resistance building up on pest populations.

NEDI: National Estimate of Daily Intake. An estimate of intake of pesticide in the diet over the long-term to compare to the ADI. The NEDI is based on median or mean residue levels and a high level consumption (97.5th percentile value) for the daily amounts of the food item consumed over the long term. For further details on the calculation of NEDIs please refer to section 3 of the data requirements handbook using the following link: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links.

NESTI: National Estimate of Short Term Intake. An estimate of peak intake of pesticide in the diet to compare to the ARfD. The NESTI is based on the highest residue found multiplied by a variability factor (see glossary description) and a high level consumption (97.5th percentile value) for the amount of the food item consumed over a single day. For further details on the calculation of NESTIs please refer to section 3 of the data requirements handbook using the following link: [The HSE Pesticide Website](#) then search for Consumer Exposure. Here you will find information and further links.

Neurotoxicity: Neurotoxicity is the effect of substances (called neurotoxins) which alter the normal working of an animal's nervous systems and/or damage the nervous tissue.

No Observed Adverse Effect Level (NOAEL): The greatest concentration or amount of a substance, found by experiment or observation, which causes no detectable adverse alteration of morphology, functional capacity, growth, development or life span of the target organism under defined conditions of exposure.

Off Label: See EAMUs

Origin: The brand name annex reports the origins of the samples tested. This can mean different things depending on the commodity. For example, butter is often labelled as 'UK origin'; however, the majority of it comes in bulk from New Zealand and is split into smaller blocks and packaged in the UK. Lettuce is a fresh produce and 'UK origin' usually means that it has been grown and packaged in the UK. Processed commodities such as cereal bars often contain multiple raw ingredients, each of which may come from a different source/origin. Therefore, the origin of the produce usually reflects the place where it was manufactured. The PRiF report the origin as stated on the packaging or labelling of the commodity concerned, unless other more accurate information is available to indicate that the origin is from elsewhere. Some products are listed as 'unknown origin' because the labelling does not give this information.

Parent: The chemical form of a pesticide as applied to plants, as opposed to metabolites and breakdown products.

Percentile: A percentile is a value that divides a sample of measurements at a specific point when they are listed in ascending order of magnitude. For example, the 97.5th percentile from a food consumption survey is a value that is equal to or more than 97.5% of the measurements and equal to or less than 2.5% of the measurements. So, in a sample of 40 daily food consumption values, the 97.5th percentile is equal to or more than 39 of the measurements. Such high percentile estimates of food consumption are used in risk assessments as they are more protective than using average consumption levels.

Permitted Level (PL): The permitted levels (expressed as mg/kg), in specific commodities, of some substances which can be classified as pesticides but are controlled under the Miscellaneous Food Additives Regulations 1995 (S.I. 1995 No. 3187).

Pesticide: A pesticide is any substance, preparation or organism prepared or used for destroying any pest. The majority of pesticides sought by the PRiF in its monitoring are those used to control pests in agricultural crops, although non-agricultural products may be included where there is a specific reason for doing so, e.g. where there are implications in terms of possible intakes of residues.

Probabilistic Modelling: The usual estimates of consumer exposure use single high values for both consumption amounts and residue levels. Whilst these are based on realistic UK dietary survey data and residue levels, they tend to overestimate most representative intakes. This is

because they do not take into account actual variations in both amounts consumed and residue levels. Probabilistic modelling is a technique that considers all the possible different combinations of consumption and residue levels. This provides information on the probability of particular intakes occurring.

Rapid Alert System for Food and Feed (RASFF): The European Commission's Rapid Alert System for Food and Feed (RASFF) allows member authorities (EU and EFTA member States) to quickly exchange information about measures taken when responding to risks detected in food or feed. This exchange of information helps authorities in countries inside the European single market to act more rapidly and in a coordinated way in response to a possible health threats caused by food or feed.

RASFFs notifications about pesticide residues are sent when a residue is over the MRL taking into account measurement uncertainty and a potential consumer risk has been identified. For pesticide residues in food traded in the single market this means when a risk assessment has identified that risk to people eating the food cannot be ruled out.

More information is available on the European Commission website at [RASFF - Food and Feed Safety Alerts](#).

Relationship between GAP and MRLs: The MRL can be defined as the maximum concentration of a pesticide residue (expressed as mg/kg) likely to **occur** in or on food commodities and animal feeds, after the use of the pesticide according to the GAP.

Reporting Limit: The reporting limit is the lowest level at which residues will be reported by a laboratory for a survey, as agreed in advance with the laboratory. It can be equal to or higher than the limit of quantification (sometimes also referred to as the limit of determination). The limit of quantification is the lowest concentration that has been validated to meet strict acceptance criteria, and may vary slightly from laboratory to laboratory depending on the equipment available and operating procedures used. The reporting limit should be at or below the MRL. For a small number of pesticides e.g. monocrotophos, we are looking for the pesticide below the LOD MRL because we are specifically interested in prevalence in food due to the nature of the pesticide. In such cases, tests are performed in the laboratory to support the lower reporting limits by validating the method at lower limits. **'None were detected above the Set RL':** This term is used in the Brand Name Annex, where no residues were found above their reporting limit.

Residue: Residues may be present in vegetable and animal products following the application(s) of a pesticide(s). They may not only include the pesticide that was applied but other degradation or reaction products and metabolites that may be of toxicological significance. The levels or amounts of residues present are expressed in milligrams of the chemical in a kilogram of crop/food/commodity (mg/kg), or parts per million.

Risk Assessment: A risk assessment is carried out when residues are found in foods to determine whether, at the levels found, they present a concern for consumer health or not. Consumer risk assessments are routinely conducted as part of the approval process for pesticides and are based on residue trials. Approval of a pesticide is only recommended when the consumer risk is acceptable.

Safety Factor: Values used in extrapolation from experimental studies in animals (usually 100) or humans (usually 10) to the population: for PRiF assessments this represents a value by which the NOAEL is divided to derive an ADI or ARfD. The value depends on the nature of the effect, the dose-response relationship, and the quality of the toxicological information available. The

use of such a factor accounts for possible differences in susceptibility between the animal species tested and humans, and for variation between different individuals in the population. The terms 'uncertainty factor' and 'assessment factor' are also sometimes used for this factor; the PRiF will use 'safety factor'.

Sample: The nature of all samples is as designated in the EC's 'sampling' Directive – 2002/63/EC. Examples are: apple – at least 10 apples weighing at least 1 kg; grapes – at least 5 bunches, weighing at least 2 kg.

Technical Exceedances: When an MRL has been set at the LOD because there have been no data to support a higher level. In the context of this report, 'technical exceedances' always relate to produce from third countries.

Variability Factor: A value that describes the variation in residue levels between the highest unit level and the average level in samples made up of many units. Internationally this is agreed to be the 97.5th percentile unit residue level divided by the average of the sum. The variability factor multiplied by the measured residue level from a composite sample (i.e. a sample made up by mixing several units before analysis) gives an estimate of the likely higher residue levels that may have occurred in individual units. These estimated higher levels are used in short-term risk assessments involving fruit and vegetables where consumers eat only a portion of a single item, e.g. melon, or a small number of units e.g. apples and potatoes.