

⠠	⠠	⠠	⠠	⠠	⠠	⠠	⠠
A	q	u	i	e	t	t	e
⠠	⠠	⠠	⠠	⠠	⠠		
#	2	.	5	m	g		
⠠	⠠	⠠	⠠	⠠	⠠	⠠	
T	a	b	l	e	t	s	

# AQUIETTE® 2.5 mg

## Oxybutynin hydrochloride Tablets

**Helps control Overactive Bladder Symptoms**  
 ✓urgent urination ✓frequent urination ✓accidental leaks

- When symptoms have been present for more than 1 month
- When bladder training alone for at least 6 weeks is not enough to control symptoms


28  
Tablets

28  
Tablets


scan me:  
  
**AQUIETTE®**  
 Oxybutynin hydrochloride  
 2.5 mg Tablets



**DIRECTIONS - Women aged between 18 - 65. Use this medicine in combination with bladder training. For oral use.**

How much?	How often?	After 6 weeks if symptoms improve	See your doctor
1 tablet 	2 to 3 times a day. Wait at least 6 hours between doses Use lowest effective dose	Take for another 6 weeks then stop taking Aquiette® and continue with bladder training alone	If symptoms get worse or do not get better after 6 weeks If you still need Aquiette after 12 weeks.

Do not take more medicine than the label tells you to.

BN  
 EXP  
  
 Batch Number  
 Expiry Date

**Read the enclosed leaflet before use.**  
**Aquiette® 2.5mg Tablets (Oxybutynin hydrochloride)**  
**Helps control overactive bladder symptoms:**  
 ✓ In women aged 18 to 65 years  
 ✓ When bladder training alone for at least 6 weeks is not enough to control symptoms  
 ✓ When symptoms have been present for at least one month

**Symptoms of overactive bladder:**  
 ✓ Sudden urgent need to pass water  
 ✓ Needing to go more than 8 times a day or more than once at night  
 ✓ Sometimes leading to accidental leaks  
 X Not suitable for stress incontinence symptoms alone (accidental leaks on coughing, sneezing or exercise).

**DO NOT TAKE and tell your pharmacist if you:**  
 X Have blood in your urine or cloudy urine  
 X Have difficulty or pain when you pass water  
 X Have fever or pain in the lower back at the side  
 X Have had repeated urinary tract infections in the last year  
 X Have a feeling of a bulge or lump in the vagina or vaginal problems of dryness, itching or burning  
 X Have period problems such as heavy periods, bleeding between periods or after the menopause  
 X Have a bladder problem caused by a nerve condition, or frequent urination due to diabetes, heart, or kidney disease  
 X Have muscle disease, glaucoma, stomach or bowel problems (such as constipation)  
 X Are pregnant or breast feeding  
**If you are taking any other medicines, speak to your pharmacist.**

**KEEP OUT OF THE SIGHT AND REACH OF CHILDREN**

**Warning:** This medicine may make you sleepy. If this happens, do not drive or use tools or machines.  
 Do not drink alcohol.

**Active ingredient:** 2.5 mg Oxybutynin hydrochloride  
**Also contains:** Lactose. Store below 25°C in a dry place See leaflet for further information.  
 MA holder: Maxwellia Ltd, Alderley Park, Alderley Edge, SK10 4TG, UK  
 PL number: 42807/0001



0 00000 00000 0 >



⠠	⠠	⠠	⠠	⠠	⠠	⠠	⠠
A	q	u	i	e	t	t	e
⠠	⠠	⠠	⠠	⠠	⠠		
#	2	.	5	m	g		
⠠	⠠	⠠	⠠	⠠	⠠	⠠	
T	a	b	l	e	t	s	

**KEEP OUT OF THE SIGHT AND REACH OF CHILDREN**

**Active ingredient:**  
2.5 mg Oxybutynin hydrochloride  
**Also contains:** Lactose.  
See leaflet for further information.

**Warning:** This medicine may make you sleepy. If this happens, do not drive or use tools or machines.  
Do not drink alcohol.

MA holder: Maxwellia Ltd, Alderley Park, Alderley Edge, SK10 4TG, UK  
Store below 25°C in a dry place  
PL number: 42807/0001



P 0 000004000000 0 >

# AQUIETTE® 2.5 mg Tablets

## Oxybutynin hydrochloride

**Helps control Overactive Bladder Symptoms**  
✓urgent urination ✓frequent urination ✓accidental leaks


- When symptoms have been present for more than 1 month
- When bladder training alone for at least 6 weeks is not enough to control symptoms

30  
Tablets

30  
Tablets



**DIRECTIONS - Women aged between 18 - 65. Use this medicine in combination with bladder training. For oral use.**

How much?	How often?	After 6 weeks if symptoms improve	See your doctor
1 tablet 	2 to 3 times a day. Wait at least 6 hours between doses Use lowest effective dose	Take for another 6 weeks then stop taking Aquiette® and continue with bladder training alone	If symptoms get worse or do not get better after 6 weeks If you still need Aquiette after 12 weeks.

**Do not take more medicine than the label tells you to.**

**Read the enclosed leaflet before use.**

**Aquiette® 2.5mg Tablets (Oxybutynin hydrochloride)**

**Helps control overactive bladder symptoms:**

- ✓ In women aged 18 to 65 years
- ✓ When bladder training alone for at least 6 weeks is not enough to control symptoms
- ✓ When symptoms have been present for at least one month

**Symptoms of overactive bladder:**

- ✓ Sudden urgent need to pass water
- ✓ Needing to go more than 8 times a day or more than once at night
- ✓ Sometimes leading to accidental leaks
- ✗ Not suitable for stress incontinence symptoms alone (accidental leaks on coughing, sneezing or exercise).

**DO NOT TAKE and tell your pharmacist if you:**

- ✗ Have blood in your urine or cloudy urine
- ✗ Have difficulty or pain when you pass water
- ✗ Have fever or pain in the lower back at the side
- ✗ Have had repeated urinary tract infections in the last year
- ✗ Have a feeling of a bulge or lump in the vagina or vaginal problems of dryness, itching or burning
- ✗ Have period problems such as heavy periods, bleeding between periods or after the menopause
- ✗ Have a bladder problem caused by a nerve condition, or frequent urination due to diabetes, heart, or kidney disease
- ✗ Have muscle disease, glaucoma, stomach or bowel problems (such as constipation)
- ✗ Are pregnant or breast feeding

**If you are taking any other medicines, speak to your pharmacist.**

BN  
EXP

Batch Number  
Expiry Date

