

Care Home outbreak testing for COVID-19 flowchart: staff and residents

Quick reference guide for the COVID-19 testing in Adult Social Care guidance.

Have there been 2 or more positive COVID-19 cases in a 14-day period?



If any staff member or resident becomes symptomatic, they should take an **LFD test** on day **0** and then on day **2**. If either of these are positive, follow the guidance in the **IPC - COVID-19 Supplement**.

Positive case If there is one positive case in the home, staff should conduct rapid response testing for 5 days. If there are further linked cases within the 5-day period, contact the HPT (or relevant authority) for advice.



Contact the HPT and start rapid response testing

Begin rapid response testing and contact the HPT (or relevant authority) if you suspect an outbreak.

An outbreak is 2 or hore positive (or clinically suspected) linked cases of COVID-19 associated with the same setting within a 14-day period.

Outbreak Testing (if a vise I)

Day 1: all staff and residents take PCR* and LFD tests.

Days 4 to 7: staff and residents take one PCR* and one LFD cone of these days.

Day 7 onwards: No further testing until outbreak recovery testing unless individuals become symptomatic.

2 Has anyone tested positive in the last 10 days of the outbreak?



Conduct outbreak recovery test in All staff and residents conduct a FNR* test.



Outbreak continues

Continue LFD testing any newly symptomatic individuals.

Were any new positives found?



Continue outbreak measures if new cases could be linked to the original outbreak. You can get advice from your HPT or relevant authority. After 10 days of no new positives, repeat step 2.

3 After outbreak recovery testing

Have there been 2 or more positive (or clinically suspected) linked cases in the home after outbreak recovery testing?



No further testing unless individuals become symptomatic.



This should be classed as a new potential outbreak and the care home should follow the process from **step 1**.



Scan QR Code for <u>COVID-19 testing</u> in Adult Social Care guidance