dstl

Depleted Uranium Survey Report: Kirkcudbright Training Area 2010 Part 2 Marine Environment

Aaron Scarlett

DSTL/TR52797 V1 31 March 2011 Environmental Sciences Department c/o INM Crescent Road Alverstoke Hampshire PO12 2DL © Crown copyright 2011 Dstl



Dstl is part of the Ministry of Defence

This document has been prepared for MOD and, unless indicated, may be used and circulated in accordance with the conditions of the Order under which it was supplied.

It may not be used or copied for any non-Governmental or commercial purpose without the written agreement of Dstl.

© Crown copyright 2011 Defence Science and Technology Laboratory UK

Approval for wider use or release must be sought from:

Intellectual Property Department Defence Science and Technology Laboratory Porton Down, Salisbury, Wiltshire, SP4 0JQ.

Authorisation			
	Name	Signature	Date
Chief Scientist	Dr Adrian Baker	Alfol ==>	6/4/11
Project Manager	Dee Emerson	10000	614/11
Technical Reviewer	Ron Brown	MBrown	6/4/11

Executive Summary

Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the extent of any radiological impact of the firings on the terrestrial and marine environments and any associated risk to humans.

This report presents the findings of the marine survey undertaken in the areas surrounding KTA during 2010; the terrestrial survey is reported separately in Part 1. The survey was undertaken to monitor the levels of any depleted uranium in the marine environment resulting from operations on the site and to identify the extent of any environmental transfer processes.

None of the samples analysed were radioactive within the meaning of the Radioactive Substances Act 1993, nor did they exceed a very small fraction of the Generalised Derived Limits advised by the Health Protection Agency (formerly the National Radiological Protection Board). In fact, the measured levels of uranium were found to be consistent with natural background levels.

The results of the 2010 survey are consistent with historical survey findings. There is no evidence to indicate that members of the public are exposed to a radiological hazard from the marine environment as a result of test firing DU ammunition at Kirkcudbright.

This page is intentionally blank

Table of Contents

Execu	ecutive Summary					
List o	ist of Figures					
List o	f Tables		6			
1	Introduc	tion	7			
2	Backgrou	ınd	7			
3	Depleted Uranium (DU)					
4	Different	iating DU from Natural Uranium	10			
5	5 Reference Levels					
6	Methodology					
7	Results and Interpretation					
8	8 Evaluation of Potential Exposure Pathways					
9	Conclusions					
10	List of R	eferences	22			
11	KTA Ma	rine Survey Results	24			
Histo	rical Data		27			
ANNI	EX A	Issues to be considered when interpreting or comparing uranium data	29			
ANNI	ANNEX B Change in the ²³⁸ U/ ²³⁴ U activity ratio of a medium containing natural uranium with the addition of depleted uranium		32			
ANNEX C Variability of uranium concentration and uranium isotopic ratios in man environmental samples		Variability of uranium concentration and uranium isotopic ratios in mari environmental samples	ine 33			
ANNI	EX D	Reference values for uranium in the Solway Firth and the UK	35			
Initia	l distribut	ion	37			

List of Figures

Figure 1. Number of DU projectiles fired at KTA between 1982 and 2010.	8
Figure 2. Approximate cumulative mass of DU projectiles fired at KTA between 1982 a 2010.	and 8
Figure 3. Kirkcudbright inter-tidal sampling locations 2010.	12
Figure 4. Approximate underwater sampling locations 2010 (showing line of fire for eabattery).	ich 15

List of Tables

Table 1. Mass compositions of uranium isotopes in natural and depleted uranium.	9
Table 2. Reference levels relevant to the Kirkcudbright marine survey.	11
Table 3. Summary of inter-tidal sample collection and measurements 2010.	14
Table 4. Summary of sample analyses - KTA 2010.	17
Table 5. Inter-tidal environmental gamma dose rates - KTA 2010.	24
Table 6. Inter-tidal sediment: alpha spectrometry analysis results showing total uranium an isotopic ratios - KTA 2010.	nd 24
Table 7. Seaweed samples: alpha spectrometry analysis results showing total uranium andisotopic ratios – KTA 2010.	25
Table 8. Biota samples: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2010.	25
Table 9. Underwater sediment: alpha spectrometry analysis results showing total uraniumand isotopic ratios – KTA 2010.	26
Table 10. Historical summary of total uranium and isotopic ratios for sediment and biota samples 1998 to 2010.	28

1 Introduction

- 1.1 Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the extent of any environmental impact of the firings on the terrestrial and marine environments and any associated radiological risk [1 to 13].
- 1.2 This report presents the findings of the marine survey undertaken in the areas surrounding KTA during 2010; the terrestrial survey is reported separately in Part 1 [14]. The survey was undertaken to monitor the levels of any uranium in the marine environment resulting from operations on the range and to identify the extent of any environmental transfer processes.
- 1.3 Since its inception in 1980, the monitoring programme has evolved to incorporate changes in best practice and increased knowledge of the local environment. The changes to sampling protocols over the years were fully explained in the marine report for 2005 [9]. Since 1996, the survey methodology has remained consistent and involves the annual sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood are also sampled.

2 Background

- 2.1 The KTA range is located on the coast of Dumfries and Galloway, near Castle Douglas. In April 2006, the range became part of the Defence Training Estate (DTE).
- 2.2 DU has been released into the environment at KTA as a consequence of the test firing of DU ammunition during design and accuracy assessment trials. DU projectiles are fired through soft vertical targets and continue their trajectory, coming to rest in the Solway Firth.
- 2.3 Testing of projectiles has historically taken place at five locations at KTA. Each battery location had a designated target and line of fire and hence a predictable area of impact ranging from several hundred metres to approximately 7 kilometres offshore. Although a small number of the DU rounds malfunctioned and impacted on land, the vast majority entered the Solway Firth.
- 2.4 The number of DU rounds fired each year at KTA from the five firing locations and the cumulative mass fired to date, are presented in Figures 1 and 2 respectively.



Figure 1. Number of DU projectiles fired at KTA between 1982 and 2010.



Figure 2. Approximate cumulative mass of DU projectiles fired at KTA between 1982 and 2010.

3 Depleted Uranium (DU)

- 3.1 Uranium is a naturally occurring radioactive material which exists as three isotopes: uranium-238 (²³⁸U), uranium-235 (²³⁵U) and uranium-234 (²³⁴U). The typical mass composition of these isotopes is shown in Table 1. In natural uranium, ²³⁸U and ²³⁵U are in approximate radioactive equilibrium with their daughter products¹. Together these isotopes emit a range of alpha and beta particles along with gamma radiation. Being a heavy metal, the chemical toxicity of uranium is approximately equal to that of lead.
- 3.2 Uranium in an 'enriched' form is used as fuel in nuclear reactors. The enrichment process increases the concentration of 235 U (above 0.72%) in comparison to the natural form. The by-product of this process is 'depleted' uranium (DU), which has a reduced concentration of 235 U. Uranium-234 is also removed in the depletion process, meaning that DU is consequently less radioactive than natural uranium (the specific alpha activity of the DU alloy used at Kirkcudbright is approximately 1.4 x 10⁷ milli becquerels per gram (mBq/g), compared to around 2.5 x 10⁷ mBq/g for natural uranium). The mass compositions of DU and natural uranium are presented in Table 1 below.

Form of Uranium	²³⁸ U	²³⁵ U	²³⁴ U
Natural uranium	99.274%	0.72%	0.00554%
The DU used at Kirkcudbright	99.8%	0.20%	0.0008%

Table 1. Typical mass composition of uranium isotopes in natural and depleted uranium.

3.3 As discussed in paragraph 3.1, 234 U normally exists in approximate equilibrium with 238 U in the natural environment. In comparison, DU exhibits a 238 U/ 234 U activity ratio of between 7:1 and 8:1, dependent on the degree of depletion achievable by different methods of processing. This distinction is important in differentiating DU contamination from naturally occurring uranium in the environment (see Section 4). For the remainder of this report, isotopic ratios will be stated in terms of activity rather than mass and as a single value representing the ratio of becquerels of 238 U to 1 becquerel of 234 U (i.e. a 238 U/ 234 U ratio of 7 rather than 7:1).

¹ A radioactive decay series occurs when a heavy radionuclide decays into successively lighter radionuclides known as daughter products. For example, ²³⁸U decays to ²³⁴Th, then ²³⁴mPa, then ²³⁴U and so on until a stable element is reached (²⁰⁶Pb).

4 Differentiating DU from Natural Uranium

- 4.1 The fundamental requirement of the DU environmental monitoring programme is to quantify the impact of DU firing. This is achieved partly by measuring the amount of total uranium in environmental materials and using this figure as an upper bound of DU contamination levels. However, as uranium is present at detectable levels in most environmental materials, this overestimates the risk. More sophisticated analyses involve the specific measurement of ²³⁸U and ²³⁴U isotopes (by activity and/or mass). Although isotope measurements are used in this survey, references to total uranium measurements are included for consistency with historic reports. The limitations of using total uranium concentrations are discussed further in Annex A.
- 4.2 A convenient fingerprint marker for DU contamination is the ${}^{238}U/{}^{234}U$ activity ratio. The DU fired at KTA has a ${}^{238}U/{}^{234}U$ activity ratio of approximately 7, whereas natural uranium in the environment has an activity ratio close to unity. Environmental samples are therefore analysed for isotopes of ${}^{238}U$ and ${}^{234}U$ to determine activity ratios and hence identify the origin of the uranium.
- 4.3 Substantial deposition of DU in the environment (in addition to an existing natural uranium background) is required before the ²³⁸U/²³⁴U activity ratio diverges significantly from its natural ratio. An illustration of the impact of DU contamination on the isotopic ratio is given in Annex B. For the ratio to approach 7 in an analytical sample, the mass of DU would have to be approximately one hundred times the mass of the uranium that is naturally present. Hence, the lower the natural uranium background, the lower the levels of DU contamination that may be detected by isotopic analysis.
- 4.4 Isotopic quantification is achieved by techniques such as alpha spectrometry and mass spectrometry. Alpha spectrometry can detect uranium to parts per billion, which is equivalent to mBq per kg, or to lower levels if count times are increased. Mass spectrometry is more sensitive, but the lower levels detectable are of no recognised health significance. Isotopic information can also be yielded from gamma spectrometry analyses, although limits of detection are not generally sufficient for measurement of environmental levels.

5 Reference Levels

- 5.1 The Depleted Uranium Firing Environmental Review Committee (DUFERC), on which the MOD is represented, has agreed investigation/action levels for levels of DU in the terrestrial environment. Reference levels for the marine environment are taken directly from legislation and guidance, as discussed below.
- 5.2 For uranium in sediment, the activity concentration can be compared to the Schedule 1 limit defined in the Radioactive Substances Act 1993 (RSA93) [15] and Generalised Derived Limits (GDLs) advised by the Health Protection Agency (formerly the National Radiological Protection Board) [16]. GDLs for uranium were last updated in 2000 and were referred to by the Royal Society in their studies of the potential health effects of using DU munitions [17]. They are based on a 1 milli-sievert (mSv) potential exposure to a member of the public: the dose limit set by the International Commission on Radiological Protection (ICRP) and incorporated into UK law [18].
- 5.3 Uranium levels in foodstuffs such as fish, crustaceans and molluscs can be compared against the relevant GDL (expressed as fresh mass of the edible fraction). For biota such as seaweed, the European Council Food Intervention Level (CFIL) for 'other foodstuff' is considered relevant [19].

Source	Reference Level	Activity concentration (mBq/g)
RSA 93	Schedule 1 limit - level at which regulatory control is required.	11,100
	Generalised Derived Limit: Marine sediment (dry weight)	100,000
NRPB (2000)	²³⁸ U in marine fish (fresh weight)	200
NKI B (2000)	²³⁸ U in crustaceans (fresh weight)	1,000
	²³⁸ U in molluscs (fresh weight)	1,000
CEC Regulation (Euratom) 2218/89	European Council Food Intervention Level (CFIL) for 'other foodstuff'	1,250

5.4 Reference levels relevant to the Kirkcudbright marine survey are provided in Table 2.

Table 2. Reference levels relevant to the Kirkcudbright marine survey.

5.5 In addition to these reference values, it is also possible to compare the results with those of similar surveys carried out in the UK. The Radioactivity in Food and the Environment (RIFE) [20] reports present the findings of an annual independent survey carried out by the Centre for Environment, Fisheries and Aquaculture Science (CEFAS) on behalf of the environment agencies and the Food Standards Agency. Where possible, monitoring results are compared against those of the RIFE reports and other relevant surveys such as

Reference 21. The latter includes monitoring data for Sandyhills Bay (also monitored by Dstl); which is sufficiently close to KTA to be similar in geology, but sufficiently far away to be considered unaffected by activities at the range.

6 Methodology

6.1 The survey methodology comprises the sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood are also sampled.

Inter-tidal zone sampling

6.2 Sampling and dose rate measurements were carried out at 11 locations as shown at Figure 3. These locations have been chosen to provide a suitable distribution of sampling points spanning the areas likely to be affected by the DU firing at KTA. It should be noted that sampling location 5 was re-positioned approximately 2 km to the North-East (Balcary Bay) in 2007 due to ongoing problems with safe access to the area.



Figure 3. Kirkcudbright inter-tidal sampling locations 2010.

6.3 The Ordnance Survey of Great Britain grid references for the sampling locations in 2010 are given below:

1.	South Carse	NX 99333 59800
2.	Sandyhills Bay	NX 89553 55298
3.	Port o'Warren Bay	NX 87892 53427
4.	Port Donnel	NX 84800 53400
5.	Balcarry Bay	NX 82182 49544
6.	Abbey Burn Foot	NX 74276 44391
7.	Mullock Bay	NX 70996 43765
8.	Lower Nunton Bay	NX 65868 48605
9.	Brighouse Bay	NX 63469 45708
10.	Carrick Point	NX 57459 50502
11.	Mossyard Bay	NX 55195 51777

- 6.4 Inter-tidal sediment was collected from each sampling location for analysis by alpha spectrometry. The sediment was collected as close to the low water mark as possible, giving due consideration for the safety of the survey team. Due to the abundance of available inter-tidal sediment, composite samples were collected by taking the top few millimetres of sediment from a number of locations until enough material was collected to fill a 0.8 litre container. Shells, stones and other unwanted debris were removed from samples and any excess water was drained off.
- 6.5 Seaweed samples were collected where available from sampling locations for analysis by alpha spectrometry. The most recent growth of seaweed was collected by cutting the top 5cm of fronds from a number of locations and combining the material to form a composite sample (approximately 1.8 litre sample volume). The collected samples, which consisted of a single species (*Fucus vesiculosus*), were washed in sea-water to remove any sediment.
- 6.6 Mollusc samples were collected from each sampling location where found in sufficient amounts. A composite sample was collected in a 1.8 litre sample container and washed in sea-water to remove any sediment. In 2010, the only mollusc species available for sampling was the common mussel (*Mytilus edulis*). The samples were washed a second time and boiled on the day of collection to prevent decomposition before analysis. The samples were placed in a re-sealable bag and frozen prior to transport to the laboratory for analysis by alpha spectrometry.
- 6.7 Environmental gamma dose rates were recorded at each sampling location using the Mini Instruments 6-80/81 fitted with an MC71 compensated Geiger-Muller tube. The instrument was allowed to record over a period of 300 seconds at three points along the low water mark and at one point at the high water mark (at one metre above the ground). The recorded count rates were converted to an environmental gamma dose rate using the methodology described in Reference 22.
- 6.8 A summary of samples collected and dose rate measurements taken from the inter-tidal zone is provided in Table 3.

Sample	Sample typ	pe and number	Environmental dose rates		
area number	Sediment	Seaweed	Mussel	LWM @ 1m height	HWM @ 1m height
1	~	×	×	~	\checkmark
2	\checkmark	\checkmark	×	✓	\checkmark
3	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
4	~	✓	✓	✓	✓
5	~	✓	✓	✓	✓
6	~	✓	×	✓	✓
7	~	✓	×	✓	✓
8	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
9	\checkmark	\checkmark	×	\checkmark	\checkmark
10	~	\checkmark	\checkmark	~	✓
11	\checkmark	\checkmark	×	\checkmark	~

Table 3. Summary of inter-tidal sample collection and measurements 2010.

Notes: LWM: low water mark, HWM: high water mark, **×**: sample not collected.

Underwater sediment sampling

6.9 Underwater sediment samples were collected from 25 locations by use of a scooping device with a rubberised bag which was lowered to the sea bed at each sampling point. The sample locations are shown in Figure 4 and the co-ordinates (latitude and longitude) are provided below:

1.	N 54 45 078	W 03 59 782	14. I	N 54 45 375	W 03 57 744
2.	N 54 45 600	W 03 59 808	15. I	N 54 45 673	W 03 57 125
3.	N 54 44 527	W 03 59 746	16. I	N 54 44 136	W 03 57 108
4.	N 54 45 569	W 04 02 577	17. I	N 54 44 575	W 03 57 081
5.	N 54 45 032	W 04 02 541	18. I	N 54 42 874	W 04 01 311
6.	N 54 44 479	W 04 02 522	19. I	N 54 42 750	W 04 01 290
7.	N 54 45 211	W 04 01 174	20. I	N 54 43 235	W 04 02 493
8.	N 54 45 083	W 04 01 243	21. I	N 54 42 732	W 04 02 476
9.	N 54 44 827	W 04 01 391	22. I	N 54 42 225	W 04 02 451
10.	N 54 44 684	W 04 01 477	23. I	N 54 43 292	W 04 00 243
11.	N 54 44 991	W 03 58 807	24. I	N 54 42 785	W 04 00 226
12.	N 54 44 856	W 03 58 758	25. I	N 54 42 258	W 04 00 192
13.	N 54 45 484	W 03 57 969			

6.10 The sampling locations may vary slightly due to the effect of underwater currents on the sampling rig which was dragged along the sea bed for distances ranging from 50 to 100

metres. The co-ordinates given in paragraph 6.9 should therefore be considered as the approximate centre point of sampling areas of approximately 100 metres in radius.

6.11 Sediment collected from the sea-bed was screened for the presence of DU fragments by monitoring using a Mini Monitor and 44B probe. A portion of the sediment (approximately 0.8 litres) was collected for analysis by alpha spectrometry.



Figure 4. Approximate underwater sampling locations 2010 (showing line of fire for each battery).

Locally sourced seafood

6.12 Locally sourced seafood is sampled as part of the Kirkcudbright marine survey to determine the extent of any potential transfer of uranium through the food chain. In 2010, a total of 3 lobsters (*Homarus gammarus*) and 3 crabs (*Cancer pagurus*) were obtained for analysis by alpha spectrometry. The samples were boiled on the day of collection before being frozen and transported to the laboratory.

Sample descriptors

- 6.13 Each sample was given a unique sample descriptor. For inter-tidal and underwater sediment, seaweed and mussel samples, the descriptor is followed by a number which relates to the specific sample point. For samples which do not come from a defined sample point (e.g. locally sourced lobster), no sample point is given. In this case, samples are given a one letter suffix to distinguish between them. Some examples of typical sample descriptors are provided below:
 - I/T 4: Inter-tidal sediment sample from sample point 4
 - U/W 8: Underwater sediment sample from sample point 8
 - Mussel 9: Mussel sample from sample point 9
 - Crab (a): Locally sourced crab sample (sample a)

Sample preparation and laboratory analysis

- 6.14 Sample preparation took place either at the time of collection or at the Dstl UKAS accredited radiochemistry laboratory. Samples were analysed by alpha spectrometry with uranium concentration reported in mBq/g (equivalent to Bq/kg) of dry weight. An outline of the analysis procedure is provided below.
- 6.15 Mussel, crab and lobster samples were boiled prior to analysis and any shells were removed. The brown and white crab meat was used for analysis, but the digestive tracts were discarded. Mussels were analysed as composite samples. The crab and lobster samples were analysed individually.
- 6.16 All samples were weighed before drying to remove moisture and then weighed again (with results being reported as dry weight). The samples were ashed to remove organic material and homogenised by hand. During this process the quantity of uranium in each of the samples does not alter significantly. The ashed samples were boiled in concentrated mineral acid (nitric acid and hydrochloric acid) to remove the 'loose' and leachable uranium from the sample. Recalcitrant matrices such as mineral grains are not broken down by the process and hence the natural uranium within them was not extracted. The samples were filtered to remove solids.
- 6.17 Uranium separation was carried out by extraction chromatography. Each eluted sample was electro-deposited onto a stainless steel planchette and the activity of each planchette was counted in a low background alpha spectrometer with a silicon surface barrier.

7 **Results and Interpretation**

7.1 A summary of the results for all marine samples collected in 2010 is given below in Table 4. The full marine monitoring results are provided in Tables 5 to 9 in Section 11. Historical monitoring results for the Kirkcudbright marine survey for the years 1998 to 2010 are presented in the Historical Data Section.

Sample Type		No. of samples	Total uranium concentration (mBq/g)					
	Number of Samples	possibly containing detectable DU ²	Mean	Standard deviation of the mean	Minimum	Maximum		
Inter-tidal Sediment	11	0	20.39	6.01	11.69	27.48		
Seaweed	10	0	16.54	5.75	7.60	23.00		
Mussels	5	0	8.06	1.11	6.94	9.75		
Lobsters	3	0	0.24	0.14	0.09	0.34		
Crabs	3	0	0.81	0.58	0.48	1.49		
Underwater Sediment	25	8	23.05	3.34	18.39	32.14		

Table 4. Summary of sample analyses - KTA 2010.

Inter-tidal zone and biota sampling

- 7.2 Environmental gamma dose rate measurements for the inter-tidal monitoring locations are provided in Table 5. Measurements ranged from 0.072 to 0.103 micro grays per hour (μ Gy/h). These results are consistent with those recorded in previous surveys [1-13], indicating that they are due to natural background radiation. Measurements recorded over salt marsh at Kirkcudbright and reported in the most recent RIFE report [20] are consistent with these measurements (average of 0.082 μ Gy/h), although it should be noted that no specific measurements of the inter-tidal area were carried out as part of the RIFE survey.
- 7.3 Alpha spectrometry analysis results for inter-tidal sediment samples are provided in Table 6. No sample was radioactive within the meaning of RSA93, nor exceeded 0.03% of the GDL for marine sediment. The level of total uranium ranged from 11.69 ± 1.90 to

 $^{^{2}}$ It is assumed that an underwater sediment sample may possibly contain DU if it has a 238 U/ 234 U isotopic ratio greater than 0.8 at the 95% confidence level (i.e. after subtraction of the measurement uncertainty). However, ratios of around 1.0 were recorded even before DU firings began; further information is provided in Annexes C and D. For inter-tidal sediment and biota, samples are assumed to contain a degree of depletion if the isotopic ratio exceeds 1.0 (following subtraction of measurement uncertainty).

 $27.48 \pm 3.46 \text{ mBq/g}$, which is consistent with the typical background levels reported in the literature for the UK³ and with the results of previous surveys. The results also compare favourably with the level of ²³⁸U found at Sandyhills Bay in an independent study (14 mBq/g dry weight) [21] and those values reported by CEFAS in 2007 [see Annex D]; both of which can be seen as good indicators of background uranium levels.

- 7.4 Following subtraction of the associated uncertainty values, the isotopic ratios of all intertidal sediment samples are found to be below a value of 1.0, indicating that any uranium present is of natural origin.
- 7.5 Analysis results for the 10 inter-tidal seaweed samples (*F. vesiculosus*) are provided in Table 7. The levels of total uranium detected ranged from 7.60 ± 1.10 to 23.00 ± 2.51 mBq/g. Although the maximum values lie slightly outside the typical UK range reported in the literature (3.8 to 18.6 mBq/g, see Annex D), it is emphasised that the levels represent less than 2 % of the CFIL for 'other foodstuffs' and that the isotopic ratios are indicative of natural uranium.
- 7.6 Alpha spectrometry analysis results for biota samples collected in 2010 are presented in Table 8. It should be noted that GDLs and biota results reported by CEFAS [20] are expressed as activity per *fresh* mass. In contrast, Dstl results are expressed as activity per *dry* mass; resulting in higher activity concentrations. In order to directly compare the results, it is necessary to convert the reported result to fresh mass activity concentrations using the recorded wet:dry ratio (the preparatory drying stage leads to a reduction in mass of approximately 70%). The biota results are discussed further in the following paragraphs.
- 7.7 A total of five mussel (*M. edulis*) samples were analysed in 2010. The total uranium content ranged from 6.94 ± 0.70 to 9.75 ± 0.96 mBq/g (dry weight). Even before converting these values to the lower fresh weight equivalent, they represent less than 1% of the GDL for molluscs (1000 mBq/g). Although the results are slightly higher than those reported from Sandyhills Bay in an independent study [21], they are consistent with the results of previous surveys [1 to 13]. Furthermore, isotopic ratios suggest that any uranium present is natural in origin.
- 7.8 Trace levels of uranium were detected in the three crab (*C. pagurus*) and lobster (*H. gammarus*) samples analysed in 2010. These results are consistent with those which have been recorded previously [1 to 13] and represent less than 1% of the GDL for crustaceans (1000 mBq/g). Where it has been possible to calculate an isotopic ratio from the low levels present, this has indicated that the uranium present is naturally occurring.

Underwater sediment sampling

7.9 A total of 25 underwater sediment samples were collected in 2010. Screening of these samples with a Mini Monitor and 44B probe did not indicate the presence of any gross contamination or fragments of DU. Alpha spectrometry analysis results are shown in

³ Minerals containing uranium are widely distributed throughout the Earth's crust and the concentrations of natural uranium found locally can vary significantly due to the nature of the underlying geology. Consequently, there is no definitive reference level for uranium in soils and sediments, although there is broad agreement in the typical range of values published in the literature. Typical values in UK coastal sediment range from 3.6 to 32.3 mBq/g (dry weight) [see Annex D], but values exceeding 100 times this typical range can be found locally.

Table 9. No sample was radioactive within the meaning of RSA93 nor did it exceed 0.04% of the GDL for marine sediment. The measured levels of 238 U (9.34 ± 1.78 to 16.10 ± 2.84 mBq/g) lie within the typical UK coastal sediment range of 3.6 to 32.3 mBq/g (see Annex D). The levels of total uranium present are consistent with those found during previous surveys.

7.10 Eight of the twenty-five underwater sediment samples exhibited isotopic ratios which were greater than 0.8 at a confidence level of 95% or higher (maximum value 1.25 ± 0.35). Whilst this can be seen as an indication of low level DU contamination in marine sediments (see Annex C), it should be noted that ratios of around 1.0 have been recorded in sediments around KTA even before DU munitions testing began. In any case, the radiological implications of these results are insignificant as the levels of 238 U represent less than 0.04% of the relevant GDL.

8 Evaluation of Potential Exposure Pathways

8.1 Any contamination of the marine environment with DU could result in three potential exposure pathways for humans, as described below:

- External radiation exposure from contaminated sediment or seaweed;
- Inhalation of re-suspended DU contamination; and
- Ingestion of seaweed or seafood products contaminated with DU.

External radiation exposure

8.2 The measured radiation levels on the inter-tidal zone are consistent with natural background levels. Furthermore, only background levels of uranium isotopes have been identified by alpha spectrometry of inter-tidal sediment samples. The measurements carried out are sensitive enough to detect radiation at levels far below anything which could be considered as a health risk; it is therefore concluded that there is no external dose risk associated with the firing of DU munitions at Kirkcudbright.

Inhalation of re-suspended DU

8.3 DU which has been deposited on sediment may become re-suspended in the air especially if it is attached to items which are subject to disturbance (e.g. fisherman's netting). Once the DU has been re-suspended in the air, it is then free to be inhaled by persons in close proximity. However, the levels of uranium identified in this report are consistent with those expected due to natural background radioactivity; the amount of ²³⁸U representing a very small fraction of relevant GDLs. It is concluded that the risk from potential inhalation of re-suspended DU is indistinguishable from the risk due to natural background exposure.

Ingestion of DU contaminated foodstuffs

- 8.4 The levels of uranium isotopes found in biota samples were consistent with those expected due to natural background radioactivity and represented a very small fraction of the relevant GDLs. This indicates that, in terms of potential DU contamination, there is no risk associated with the consumption of food stuffs in the Kirkcudbright area.
- 8.5 Although the inadvertent consumption of seawater by members of the public is possible, it was recommended by SEPA in 2001 to discontinue seawater sampling. The basis of this recommendation was that the immense dilution of the Solway would never give rise to detectable levels of DU or to any significant radiation exposures.

9 Conclusions

- 9.1 The 2010 annual Kirkcudbright marine monitoring programme was undertaken to assess the levels of any DU in the environment resulting from firing of DU munitions. The monitoring programme involved the sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood were also sampled.
- 9.2 The findings of the survey indicate that the levels of uranium present in the marine environment are consistent with natural background levels. None of the samples were radioactive within the meaning of the Radioactive Substances Act 1993; nor did they exceed a very small fraction of the Generalised Derived Limits advised by the Health Protection Agency.
- 9.3 The results of the 2010 survey are consistent with historical survey findings. There is no evidence to indicate that members of the public are exposed to any radiological hazard from the marine environment as a result of the test firing of DU ammunition at Kirkcudbright.

10 List of References

- 1. Marine Environmental Survey Report DTEO Kirkcudbright 1996 DRPS 10/97.
- 2. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area DRPS 9/99.
- 3. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area DRPS DERA/CBD/DRPS/17/00.
- 4. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2000 Dstl/CR12021.
- 5. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2001 Dstl/CR12350.
- 6. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2002 Dstl/CR113980 (2005).
- 7. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2003 Dstl/CR 13981 (2005).
- 8. Toqué, C. (2006) Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2004 Dstl/CR18154 V1.
- 9. Toqué, C. (2007) Marine Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area 2005 Dstl/CR19579.
- 10. Toqué, C. (2007) Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area 2006 Dstl/CR26985 V1.
- 11. Toqué, C. (2008) Marine Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area 2007 Dstl/CR30510.
- 12. Scarlett, A. (2010) Depleted Uranium Survey Report: Kirkcudbright Training Area 2008. Parts 1 & 2 Terrestrial and Marine Environments DSTL/TR37166 and DSTL/TR37610.
- 13. Scarlett, A. (2010) Depleted Uranium Survey Report: Kirkcudbright Training Area 2009. Parts 1 & 2 Terrestrial and Marine Environments DSTL/TR40942 and DSTL/TR40950.
- 14. Scarlett, A. (2011) Depleted Uranium Survey Report: Kirkcudbright Training Area 2010. Part 1 Terrestrial Environment. DSTL/TR51735.
- 15. Radioactive Substances Act 1993. HMSO (ISBN 0-10-54 1293-7).

- 16. Generalised Derived Limits for Radioisotopes of Polonium, Lead, Radium and Uranium. Documents of the NRPB, Vol. 11, No 2 (2000).
- 17. The Health Hazards of Depleted Uranium Munitions (Part II), (ISBN 0854035745) The Royal Society (2002).
- 18. Ionising Radiations Regulations 1999 Statutory Instrument 1999 No 3232.
- 19. CEC Council Regulation (Euratom) No 2218/89 amending Regulation (Euratom) No 3954/87 laying down maximum permitted levels of radioactive contamination of foodstuffs and feeding stuff following a nuclear accident or any other case of radiological emergency. Official Journal of the European Community. L211/1 (1989).
- 20. Radioactivity in Food and the Environment 15 (2010) Centre for Environment, Fisheries and Aquaculture Science.
- 21. McDonald, P. Cook, G.T. and Baxter, M.S. (1991) Natural and Artificial Radioactivity in Coastal Regions of UK. in Radionuclides in the Study of Marine Processes (Ed. Kershaw P.J. and Woodhead D.S.), Elsevier Applied Science, London.
- 22. Her Majesty's Inspectorate of Pollution (1995) Technical Guidance Note (Monitoring) Routine Measurement of Gamma Ray Air Kerma Rate in the Environment. HMSO ISBN 0 11 753132 4.

Station number	Location	Mean dose rate (µGy/h) (n = 4)		
1	South Carse	0.085		
2	Sandyhills Bay	0.084		
3	Port o'Warren Bay	0.098		
4	Port Donnel	0.085		
5	Balcarry Bay	0.103		
6	Abbey Burn Foot	0.095		
7	Mullock Bay	0.098		
8	Lower Nunton Bay	0.081		
9	Brighouse Bay	0.074		
10	Carrick Point	0.072		
11	Mossyard Bay	0.081		

11 KTA Marine Survey Results

Table 5. Inter-tidal environmental gamma dose rates - KTA 2010.

Sample	Wet	Dry	Ashed	Measured	Measured Activity of Dry Sample (mBq/g)				
Descriptor	weight (g)	weight (g)	weight (g)	²³⁸ U	²³⁵ U	²³⁴ U	Total U	$^{238}U/^{234}U$	
IT1	116.8	70.0	65.3	12.14 ± 2.17	< 0.51	11.88 ± 2.14	24.14 ± 3.06	1.02 ± 0.26	
IT2	108.6	74.5	71.2	12.13 ± 2.21	< 0.51	12.63 ± 2.27	24.89 ± 3.17	0.96 ± 0.25	
IT3	127.7	87.9	84.6	10.19 ± 1.93	< 0.48	8.79 ± 1.75	19.33 ± 2.62	1.16 ± 0.32	
IT4	118.0	71.6	67.7	12.33 ± 2.41	< 0.62	13.65 ± 2.59	26.55 ± 3.56	0.90 ± 0.25	
IT5	120.2	74.0	69.1	13.34 ± 2.37	0.50 ± 0.35	11.75 ± 2.16	25.60 ± 3.22	1.14 ± 0.29	
IT6	132.2	88.0	83.2	7.11 ± 1.55	< 0.52	$6.60 \hspace{0.1in} \pm \hspace{0.1in} 1.48$	14.00 ± 2.17	1.08 ± 0.34	
IT7	129.4	75.2	71.0	14.61 ± 2.58	< 0.54	12.39 ± 2.29	27.48 ± 3.46	1.18 ± 0.30	
IT8	118.1	86.1	81.5	10.71 ± 2.10	< 0.55	11.52 ± 2.21	22.54 ± 3.06	0.93 ± 0.26	
IT9	146.3	113.2	110.3	6.41 ± 1.43	< 0.51	5.45 ± 1.30	11.98 ± 1.94	$1.18\pm\ 0.38$	
IT10	149.6	106.3	101.1	5.58 ± 1.31	< 0.50	5.93 ± 1.36	11.69 ± 1.90	0.94 ± 0.31	
IT11	161.9	108.5	103.4	7.95 ± 1.79	< 0.65	7.95 ± 1.79	16.05 ± 2.54	1.00 ± 0.32	

Table 6. Inter-tidal sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2010.

Sample	Wet	Dry	Ashed	Measure	Measured Activity of Dry Sample (mBq/g)					
Descriptor	weight	weight	weight	238 _{1 1}	235 _{1 1}	234 _{I I}	Total U	$238_{11}/234_{11}$		
-	(g)	(g)	(g)	U	U	U	Total U	070		
F. vesic 2	175.0	34.0	9.8	3.64 ± 0.76	< 0.22	3.91 ± 0.80	7.60 ± 1.10	0.93 ± 0.27		
F. vesic 3	109.2	41.7	8.9	10.46 ± 1.65	$0.20~\pm~0.14$	12.34 ± 1.89	23.00 ± 2.51	0.85 ± 0.19		
F. vesic 4	70.6	34.0	7.1	8.07 ± 1.30	$0.30~\pm~0.16$	8.58 ± 1.37	16.95 ± 1.90	0.94 ± 0.21		
F. vesic 5	61.9	29.8	6.6	9.91 ± 1.52	$0.48~\pm~0.21$	10.89 ± 1.64	21.29 ± 2.25	0.91 ± 0.20		
F. vesic 6	152.6	41.0	6.4	6.26 ± 0.92	0.32 ± 0.13	7.59 ± 1.08	14.17 ± 1.43	0.82 ± 0.17		
F. vesic 7	39.2	15.2	2.7	8.96 ±1.32	0.46 ± 0.18	10.09 ± 1.46	19.50 ± 1.97	0.89 ± 0.18		
F. vesic 8	56.8	19.0	5.7	5.81 ± 1.10	< 0.25	5.46 ± 1.05	11.41 ± 1.52	1.06 ± 0.29		
F. vesic 9	98.4	26.9	5.2	10.41 ± 1.48	0.50 ± 0.18	11.44 ± 1.60	22.34 ± 2.19	0.91 ± 0.18		
F. vesic 10	49.6	17.7	2.9	4.11 ± 0.65	0.14 ± 0.08	4.39 ± 0.68	8.64 ± 0.94	0.94 ± 0.21		
F. vesic 11	118.6	27.2	6.3	9.02 ± 1.33	0.30 ± 0.15	$11.\overline{23 \pm 1.60}$	$20.\overline{55 \pm 2.09}$	0.80 ± 0.16		

Table 7. Seaweed samples: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2010.

Sample	Wet	Dry	Ashed	Measure	d Activity of	Dry Sample	e (mBq/g)	Defined
Descriptor	weight	weight	weight	238 _{1 1}	235 _{1 1}	234 1 I	Total U	$\frac{\text{Ratio OI}}{238}$
-	(g)	(g)	(g)	U	U	U	Total U	0/ 0
M. edulis 3	78.6	18.9	2.0	3.18 ± 0.48	$0.10~\pm~0.06$	4.05 ± 0.59	7.33 ± 0.76	0.79 ± 0.17
M. edulis 4	78.8	18.8	1.5	3.62 ± 0.54	0.14 ± 0.07	4.04 ± 0.59	7.80 ± 0.80	0.89 ± 0.19
M. edulis 5	89.6	21.7	2.2	4.58 ± 0.65	$0.18~\pm~0.08$	5.00 ± 0.70	9.75 ± 0.96	0.92 ± 0.18
M. edulis 8	119.1	30.1	2.4	3.91 ± 0.56	0.17 ± 0.07	4.38 ± 0.61	8.46 ± 0.83	0.89 ± 0.18
M. edulis 10	99.8	25.0	2.5	3.23 ± 0.48	0.14 ± 0.07	3.57 ± 0.52	6.94 ± 0.70	0.90 ± 0.19
C.pagarus(a)	48.9	14.6	2.7	0.17 ± 0.10	< 0.11	0.28 ± 0.12	0.48 ± 0.16	0.61 ± 0.43
<i>C.pagarus</i> (b)	66.2	20.4	3.3	0.19 ± 0.10	< 0.11	0.26 ± 0.12	0.48 ± 0.16	0.74 ± 0.51
<i>C.pagarus</i> (c)	61.2	16.3	3.3	0.75 ± 0.22	< 0.12	0.71 ± 0.21	$1.49 \hspace{0.1in} \pm \hspace{0.1in} 0.31$	$1.07 \ \pm 0.45$
H.gammarus (a)	68.9	22.3	1.0	< 0.06	< 0.06	< 0.06	0.09 ± 0.05	N/A
<i>H.gammarus</i> (b)	71.5	21.5	1.1	0.17 ± 0.08	< 0.07	0.15 ± 0.07	0.34 ± 0.11	1.13 ± 0.76
<i>H gammarus</i> (c)	80.3	25.0	1.2	0.14 ± 0.07	< 0.10	0.15 ± 0.07	0.31 ± 0.10	0.93 ± 0.64

Table 8. Biota samples: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2010.

Sampla	Wet	Dry	Ashed	Measure	Measured Activity of Dry Sample (mBq/g)				
descriptor	weight	weight	weight	238 _{1 1}	235 _{T T}	234 _{I I}	Total II	$238_{11}/234_{11}$	
descriptor	(g)	(g)	(g)	U	U	U	Total U	070	
UW1	193.9	133.6	127.8	10.84 ± 2.07	0.63 ± 0.40	8.70 ± 1.78	20.17 ± 2.76	1.25 ± 0.35	
UW2	130.5	91.5	87.0	9.68 ± 1.97	< 0.59	9.38 ± 1.93	19.36 ± 2.77	1.03 ± 0.30	
UW3	171.6	112.5	90.4	9.53 ± 1.71	< 0.39	8.70 ± 1.61	18.39 ± 2.35	1.10 ± 0.28	
UW4	88.6	62.1	57.7	16.10 ± 2.84	0.82 ± 0.49	15.23 ± 2.73	32.14 ± 3.97	1.06 ± 0.27	
UW5	176.5	120.0	113.2	12.77 ± 2.62	< 0.75	11.13 ± 2.39	24.60 ± 3.58	1.15 ± 0.34	
UW6	235.5	170.6	163.3	13.17 ± 2.40	$0.63~\pm~\pm 0.41$	12.00 ± 2.25	25.80 ± 3.31	1.10 ± 0.29	
UW7	125.6	88.0	83.1	10.02 ± 1.99	< 0.58	$9.90 \hspace{0.2cm} \pm \hspace{0.2cm} 1.97$	20.32 ± 2.82	1.01 ± 0.28	
UW8	198.4	136.8	129.9	9.98 ± 2.01	< 0.60	$9.99 \hspace{0.1in} \pm \hspace{0.1in} 2.02$	20.11 ± 2.86	1.00 ± 0.28	
UW9	198.4	140.9	135.2	11.24 ± 2.52	< 0.85	12.14 ± 2.65	24.09 ± 3.70	0.93 ± 0.29	
UW10	97.4	69.9	66.1	12.11 ± 2.33	< 0.61	10.34 ± 2.09	22.77 ± 3.15	1.17 ± 0.33	
UW11	127.7	87.8	83.6	11.11 ± 2.06	0.81 ± 0.44	9.57 ± 1.86	21.49 ± 2.81	1.16 ± 0.31	
UW12	138.4	99.3	94.1	12.27 ± 2.34	0.62 ± 0.42	12.87 ± 2.42	25.76 ± 3.40	0.95 ± 0.26	
UW13	125.7	85.8	82.0	11.62 ± 2.24	< 0.58	12.91 ± 2.41	24.92 ± 3.31	0.90 ± 0.24	
UW14	102.2	71.3	68.2	11.02 ± 2.13	< 0.54	12.40 ± 2.31	23.55 ± 3.14	0.89 ± 0.24	
UW15	102.8	71.5	68.5	$10.55 ~\pm~ 2.01$	< 0.51	9.31 ± 1.84	20.03 ± 2.73	1.13 ± 0.31	
UW16	152.4	110.9	107.2	9.52 ± 1.90	< 0.54	$10.43~\pm~2.02$	$20.24 ~\pm~ 2.79$	$0.91 \hspace{0.2cm} \pm \hspace{0.2cm} 0.25$	
UW17	146.0	103.2	98.6	9.34 ± 1.78	< 0.45	$11.34 ~\pm~ 2.04$	$20.78 ~\pm~ 2.71$	$0.82 \ \pm \ 0.22$	
UW18	123.5	84.4	79.2	10.57 ± 2.01	< 0.52	9.67 ± 1.89	20.66 ± 2.78	1.09 ± 0.30	
UW19	98.6	68.3	65.6	14.48 ± 2.55	< 0.57	11.62 ± 2.19	26.59 ± 3.38	1.25 ± 0.32	
UW20	148.7	102.3	98.1	13.23 ± 2.36	< 0.54	12.69 ± 2.29	26.36 ± 3.30	1.04 ± 0.26	
UW21	161.0	112.2	104.2	10.48 ± 1.85	0.57 ± 0.34	9.82 ± 1.77	20.87 ± 2.59	1.07 ± 0.27	
UW22	148.3	101.3	96.8	14.37 ± 2.59	< 0.58	14.60 ± 2.62	29.46 ± 3.70	0.98 ± 0.25	
UW23	135.4	94.0	90.0	11.89 ± 2.10	< 0.49	11.30 ± 2.03	23.65 ± 2.93	1.05 ± 0.26	
UW24	152.6	101.9	96.1	$10.67 ~\pm~ 1.94$	< 0.46	$10.54~\pm~1.93$	21.51 ± 2.75	1.01 ± 0.26	
UW25	74.2	49.1	46.4	10.37 ± 1.85	< 0.40	12.04 ± 2.06	22.57 ± 2.77	0.86 ± 0.21	

Table 9. Underwater sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2010.

Note (for tables 6 to 9): Activity results have been rounded to two decimal places. All uncertainties are stated at a 95% confidence level. Limits of Detection (LoD) are calculated by a 'modified Currie formula' at 95% (Hurtgen C., Jerome S. & Woods M. (2000) 'Revisiting Currie – how low can you go?' *Applied radiation and Isotopes* 53 pp 45-50). The total activity is calculated from the sum of the actual activities for each isotope, regardless of the quoted LoD. Therefore, in Tables 6 to 9, where activities are reported as less than LoD for any of the uranium isotopes, the total uranium value may not be equal to the sum of the individual isotopic values.

Historical Data

To allow historical comparison, data from KTA marine surveys from 1998 to 2010 are presented in Table 10. This includes inter-tidal sediment, underwater and biota samples

Sample Type	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Inter-tidal sediment	12.2- 31.8 (10) (0.6-1.2)	11.0-32.6 (10) (0.9-1.1)	2.2- 44.7 (10) (0.8-1.2)	10.7-31.1 (9) (0.9-1.4)	12.0- 31.9 (10) (0.9-1.2)	9.8-714.4 (10) (0.8-1.3)	4.2-35.0 (11) (0.9-1.3)	9.8-24.5 (11) (0.8-1.3)	7.5-29.9 (11) (0.8-1.3)	6.9-27.6 (11) (0.9-1.1)	11.3–35.8 (11) (0.8-1.3)	8.5-26.2 (11) (0.7-1.2)	11.7-27.5 (11) (0.9-1.2)
Seaweed	5.0-19.6 (9) (0.8-1.2)	7.8-14.1 (9) (0.8-1.3)	1.3 -5.1 (9) (0.9 -1.1)	6.8-12.1 (7) (0.8-1.3)	0.2-16.7 (9) (0.7-1.2)	7.2-34.2 (9) (0.7-0.9)	9.2-22.6 (9) (0.8-1.0)	4.9-18.2 (10) (0.8-1.2)	6.9-16.0 (9) (0.8-1.0)	10.7-25.6 (10) (0.5-1.1)	5.3-12.3 (10) (0.8-1.0)	9.1-33.6 (10) (0.4-1.0)	7.6-23.0 (10) (0.8-1.1)
Mussels	7.6-12.2 (5) (0.8-1.1)	6.6-11.1 (5) (0.8-1.1)	3.6 -7.2 (5) (0.8-0.9)	2.9-11.6 (7) (0.3-0.9)	6.4-10.5 (5) (0.9-1.0)	5.6-7.4 (5) (0.7-0.9)	4.0-7.8 (4) (0.8-0.9)	5.1-10.7 (4) (0.9-1.0)	2.6-12.6 (7) (0.9-1.1)	4.8-12.0 (5) (0.6-0.9)	5.7-9.6 (5) (0.8-1.0)	6.2-13.7 (5) (0.7-0.9)	6.9-9.8 (5) (0.8-0.9)
Whelks	2.5-3.0 (3) (0.8-1.0)	0.2-0.4 (4) (0.3-1.3)	0.6-1.3 (5) (0.5-1.6)	2.0-3.7 (8) (0.5-1.4)	1.1-39.4 (8) (0.9±0.5)	×	×	0.7-2.9 (6) (0.8-1.6)	0.1-0.1 (4) (n/a)	0.4-1.2 (8) (0.9-1.1)	0.8-1.4 (6) (0.8-1.4)	×	×
Scallops	2.8-3.2 (3) (0.8-1.1)	0.7-1.1 (4) (1.3-1.5)		2.8-13.3 (8) (0.7-1.6)	×	1.5-33.0 (6) (n/a)	2.8-3.1 (2) (0.9-9.9)	2.8-13.3 (3) (n/a)	0.1-0.2 (4) (n/a)	×	×	×	×
Crabs	×	0.3-0.4 (2) (1.5-1.7)	0.2-1.4 (8) (0.1-3.7)	1.2-2.3 (2) (0.9-1.0)	$\begin{array}{c} 0.8-23.8 \\ (4) \\ (0.9, 0.5^{a}) \end{array}$	0.1-2.3 (3) (<i>n/a</i>)	×	0.4-1.6 (6) (n/a)	0.1-0.6 (5) (n/a)	0.4-0.8 (3) (1.6±0.8)	0.2-0.5 (5) (0.3-0.4)	0.3-0.6 (3) (1.2)	0.5-1.5 (3) (0.6-1.1)
Lobsters	0.2 ± 0.1 (1) (<i>n/a</i>)	0.3-0.3 (2) (0.1-0.5)		2.4-3.2 (4) (0.7-1.6)	(0.5 0.5) 1.2-1.8 (2) (n/a)	0.3-3.4 (3) (<i>n/a</i>)	$0.4 \pm 0.1 \\ (3) \\ (1.0 \pm 0.4)$	0.4-0.9 (4) (<i>n/a</i>)	0.1-0.1 (2) (<i>n/a</i>)	0.1-0.2 (4) (<i>n/a</i>)	0.1-0.3 (3) (<i>n/a</i>)	0.3-0.5 (2) (n/a)	0.1-0.3 (3) (0.9-1.1)
Underwater sediment	21.1- 28.0 (6) (0.9-1.1)	15.4-32.1 (6) (0.5-1.1)	0.2-12.8 (6) (0.8-1.3)	19.3-31.0 (34) (0.8-1.3)	15.2- 30.4 (33) (0.9-1.2)	20.2-30.9 (36) (0.9-1.2)	17.0-33.2 (64) (0.8-1.2)	16.3-24.3 (18) (0.9-1.3)	16.6-26.6 (25) (0.8-1.3)	19.5-42.3 (25) (0.8-1.3)	23-31.9 (25) (0.9-1.4)	17-28.4 (25) (0.9-1.6)	18.4-32.1 (25) (0.8-1.3)

Table 10. Historical summary of total uranium and isotopic ratios for sediment and biota samples 1998 to 2010.

Notes: Values provided on the top row show the range of total uranium values in mBq/g (e.g. 1.5 - 3.2). The value in bold shows the total number of samples collected (e.g. (12)). The values in the bottom row show the range of 238 U/ 234 U isotopic ratios (e.g. (0.8 - 1.1)). Measurement uncertainties can be found in the annual reports.

ANNEX A Issues to be considered when interpreting or comparing uranium data

- A.1 There are a number of issues that may give rise to uncertainties when interpreting or comparing uranium data as shown below. Further information is available from Reference [A1].
 - Analytical technique;
 - Statistical variation;
 - Spatial variability;
 - Temporal variability; and
 - Species variation (for plant and animal samples).

Analytical approaches

- A.2 Sediment sample results may be reported as either dry weight or wet weight depending on whether the masses of the samples were obtained prior to, or after, drying. This will have implications for comparison of results between the surveys at Kirkcudbright, which are reported as dry weight and other UK uranium in sediment data, which may be reported as wet weight. Samples reported as dry weight will appear to have concentrations of uranium approximately 20% higher than those reported as wet weight (although this will depend on the moisture content).
- A.3 For analysis techniques such as inductively coupled plasma mass spectrometry (ICP-MS) or alpha spectrometry the uranium present in a sample may be extracted into solution by either leaching the soil samples, or totally dissolving them. Total dissolution will give rise to higher uranium results, because the analysis will include all uranium, including that which is contained within the mineral grains, whereas leached samples will only contain uranium that is either easily dissolved, or is fixed to the surfaces of mineral grains. This limitation is acceptable as the primary purpose of the environmental survey is to assess any levels of DU in addition to naturally occurring uranium. However, this limitation causes a conservative bias on the ²³⁸U/²³⁴U isotopic ratios for mineral samples and other recalcitrant matrices. Total sample analysis techniques such as gamma spectrometry will give results similar to those for total dissolution. Given the differences between the results for total analysis and leached analysis, care should be taken when comparing sets of data to ensure that either the same approach has been used, or differences are appropriately discussed.
- A.4 Uranium concentrations in seaweed may be affected by contamination of surfaces with sediment particles. Preparation of seaweed for analysis may or may not involve a washing stage; hence it is important to be aware of the preparation approaches that have been applied when comparing the results of different seaweed analyses.
- A.5 Seaweed and marine biota sample results may be reported as either dry weight or wet weight depending on whether the masses of the samples were obtained prior to, or after,

drying. For marine biota, samples reported as dry weight will appear to have concentrations of uranium higher than those reported as wet weight. The relative increase in uranium concentration will depend on the moisture content, and can be as much as 300% for biota with high water content such as molluscs.

Statistical variations

A.6 There is a statistical uncertainty associated with the result of a radiochemical analysis (identified in the counting statistics for the technique). However, this uncertainty is likely to be small in comparison with the overall variation in activity between environmental samples.

Spatial variations

- A.7 DU contamination will not be uniformly distributed within a sampling area, but will depend on factors such as water flow, tidal movements and sediment drift. Repeat sampling and analysis of sediments from within an area may therefore give rise to some degree of variation.
- A.8 In addition to DU contamination due to firing at the range, there may be variations in uranium concentrations due to local anthropogenic or natural discharges. For example, natural uranium concentrations may be enhanced by the local application of phosphate based fertiliser to agricultural land. Veins rich in uranium minerals occur naturally along the coast of the Solway Firth, such as uraninite found at Needle's Eye, Caulkerbush, approximately 24 km away from KTA on the north coast of the estuary. These features are thought to be present across the region [A2, A3], although this has not been studied specifically.

Temporal variations

- A.9 There will be natural temporal variations in the uranium concentration and the abundance of the various isotopes in the samples, due to seasonal variations in rainfall. Rainfall can impact on the dissolution of uranium and its migration in surface waters.
- A.10 The activities of samples from any particular sampling site may vary from year to year. This may relate to temporal changes in uranium concentration, but will also be affected by spatial variation (see above).
- A.11 Marine plant uptake of radionuclides is affected by the period in the plant growing cycle. This phenomenon also occurs in animal uptake of radionuclides within their life cycle.

Species variations

- A.12 Plant uptake of radionuclides is affected by the substrate characteristics (uranium concentration and speciation as well as other soil physico-chemical characteristics) and varies with plant species.
- A.13 Animal uptake of uranium is affected by their life habits, feeding patterns, physiology and the uranium concentration in their foodstuffs and the environment.

Annex A References

- [A1] International Commission on Radiation Units and Measurements Report 75 (2006) Sampling for Radionuclides in the Environment in *Journal of the ICRU Volume 6 No 1*.
- [A2] Basham, I R; Milodowski, A E; Hyslop, E K; Pearce, J M. (1989) The location of uranium in source rocks and sites of secondary deposition at the Needle's Eye natural analogue site, D&G, British Geological Survey Technical Report WE/89/56.
- [A3] Milodowski, A E. *et al.* (1990) Uranium-mineralised micro-organisms associated with uraniferous hydrocarbons in southwest Scotland. Nature, 347, 465.

Mass proportion of	Activity c	oncentration (mBq/kg)**	Ratio of total activity natural	²³⁸ U/ ²³⁴ U activity ratio
DU added*	U-238	U-235	U-234	activity	
0	$3.7 \text{ x} 10^4$	$1.7 \text{ x} 10^3$	3.8 x10 ⁴	1.0	1.0
1	$7.4 \text{ x} 10^4$	$2.2 \text{ x} 10^3$	$4.3 ext{ x10}^4$	1.6	1.7
2	1.1 x10 ⁵	$2.7 \text{ x} 10^3$	$4.8 \text{ x} 10^4$	2.1	2.3
3	1.5 x10 ⁵	$3.2 ext{ x10}^3$	$5.3 \text{ x} 10^4$	2.6	2.8
4	1.9 x10 ⁵	$3.7 ext{ x10}^3$	5.8 x10 ⁴	3.2	3.2
5	$2.2 \text{ x} 10^5$	$4.1 \text{ x} 10^3$	$6.3 ext{ x10}^4$	3.8	3.6
6	2.6 x10 ⁵	$4.6 ext{ x10}^3$	6.8 x10 ⁴	4.3	3.9
7	$3.0 ext{ x10}^{5}$	$5.1 \text{ x} 10^3$	$7.2 \text{ x} 10^4$	4.9	4.1
8	3.4 x10 ⁵	$5.6 ext{ x10}^3$	7.7 x10 ⁴	5.4	4.3
9	3.7 x10 ⁵	$6.0 ext{ x10}^3$	$8.2 \text{ x} 10^4$	6.0	4.5
10	4.1 x10 ⁵	$6.5 ext{ x10}^3$	8.7 x10 ⁴	6.5	4.7
20	7.8 x10 ⁵	1.1 x10 ⁴	1.4 x10 ⁵	12.0	5.8
60	2.3 x10 ⁶	$3.1 \text{ x} 10^4$	3.3×10^5	34.1	6.9
80	3.0 x10 ⁶	$4.0 \text{ x} 10^4$	4.3 x10 ⁵	45.1	7.1
100	3.8 x10 ⁶	$5.0 \text{ x} 10^4$	5.2 x10 ⁵	56.1	7.2
200	7.5 x10 ⁶	9.8 x10 ⁴	1.0 x10 ⁶	111.0	7.4
600	$2.2 \text{ x} 10^7$	2.9 x10 ⁵	3.0 x10 ⁶	332.0	7.6
800	3.0 x10 ⁷	39 x10 ⁵	3.9 x10 ⁶	442.0	7.6
1000	$3.73 \text{ x}10^7$	$4.8 ext{ x10}^{5}$	$4.9 ext{ x10}^{6}$	552.0	7.6

ANNEX B Change in the ²³⁸U/²³⁴U activity ratio of a medium containing natural uranium with the addition of depleted uranium

Table reproduced from Volume 2 - Appendices, WS Atkins Environmental Assessment on DU Firings.

*

The value represents the additional mass of depleted uranium added (all radionuclides) relative to the original mass of natural uranium present (3 mg U/kg soil).

** Table assumes 3 mg U/kg of natural uranium present in soil in following proportion: ²³⁸U (2.978 mg /kg); ²³⁵U (0.022 mg /kg); ²³⁴U (2e-04 mg /kg), prior to addition of DU.

ANNEX C Variability of uranium concentration and uranium isotopic ratios in marine environmental samples

- C.1 There are few specific examples that demonstrate the variability of uranium concentrations and isotopic ratios within marine environmental samples. In order to understand this variability, it is useful to consider the flux of uranium between the various components of the marine environment.
- C.2 Uranium occurs naturally in seawater and its concentration generally varies in proportion to salinity. It is present in open seawater at an average concentration of 82.5 ± 5 Bq/m³, with a ²³⁸U/²³⁴U activity ratio of 0.88 ± 0.03 at a salinity of 35 %. The isotopic ratio of seawater is below unity as ²³⁴U is preferentially mobilised from the soil during chemical weathering, thus enhancing its presence in seawater [C1 & C2]. The physical mixing of low uranium river water¹ and high uranium sea water in estuary environments generally results in a dilution of uranium in the estuarine waters. As a result, uranium concentrations in estuarine water are mostly lower than open ocean values and increase linearly with salinity [C2].
- C.3 Concentrations of uranium in marine sediment are variable (from 32.5 to 1,625 mBq/g dry weight) and vary primarily as a function of the geology of the area.

Sample Type	Uranium Concentration	Typical ²³⁸ U/ ²³⁴ U ratio
Sea water	82.5 Bq/m ³	0.88
Estuarine water	$< 82.5 \text{ Bq/m}^{3}$	<0.88
Marine sediment (²³⁸ U only)	32.5-1,625 mBq/g	0.81

C.4 Levels of uranium also vary depending on the type of sediment present, as the physical and chemical characteristics of the sediment determine the amount of uranium which is concentrated from marine waters [C2 & C3]. Both low oxygenation² and low salinity³ provide favourable conditions for uranium scavenging from the water column by a variety of processes. These include the precipitation of uranium rich colloids into the low salinity zone and the reduction of uranium into insoluble forms [C3 & C4].

¹ Concentrations of uranium in rivers vary considerably, with carbonate and dissolved solids concentrations, with an average of 7.5-15 Bq/m³, and an isotopic ²³⁸U/²³⁴U activity ratio of 0.77-0.83 as ²³⁴U is preferentially mobilised during chemical weathering.

² Low oxygenation is found with increasing depth and increasing organic content.

³ Salinity is dependent on river rate of flow, proximity to river outflow and depth of water.

Annex C References

- [C1] Murphy, W M. Shock, E L. (1999) Environmental Aqueous Geochemistry of Actinides, in Uranium: Mineralogy, Geochemistry and the Environment, edited by Burns, P and Finch, R. Reviews in Mineralogy, Vol 38, The Mineralogical Society of America.
- [C2] Cochran, J K. (1982) The Oceanic Chemistry of the U- and the Th-series Nuclides, in Uranium Series Disequilibrium: Applications to Environmental Problems, edited by Ivanovich, M and Harmon RS. Clarendon Press, Oxford.
- [C3] Van den Berg, C M G. Huang, Z Q. (1984) Analytica Chemica Acta, 164, 209-222.
- [C4] Ragnarsdottir, V. Charlet, L. (2000) Uranium Behaviour in Natural Environments in: Environmental Mineralogy: Microbial Interactions, Anthropogenic Influences, Contaminated Land and Waste Management, Mineralogical Society Series Vol. 9, Mineralogical Society of Great Britain & Ireland.

ANNEX D Reference values for uranium in the Solway Firth and the UK

- D.1 For UK coastal sediment, ²³⁸U concentrations generally range from 3.6 to 32.3 mBq/g of dry weight [D1]. Values reported for inter-tidal sediment uranium concentrations in the Solway Firth vary with location and the specific uranium rich geology of the area.
- D.2 Values for ²³⁸U, ²³⁵U and ²³⁴U for sediment collected at Kirkcudbright in 2007 were reported by CEFAS as 10, 0.92 and 13 mBq/g of dry weight respectively [D2].
- D.3 Values reported for 238 U concentrations at Sandyhills Bay, in southwest Scotland, were lower than worldwide average values for seawater and sediment reported in ANNEX C. Seawater and sediment from Sandyhills Bay had 238 U concentrations of 40 ± 3 Bq/m and 14 ± 0.4 mBq/g dry weight, respectively [D3]. No other uranium isotopes were measured, so no indication of isotopic ratio could be provided.
- D.4 However, semi-quantitative analysis of shore sediment samples obtained from the mudflats offshore from the uraninite vein, located approximately 24 km east from KTA at Needle's Eye, yielded uranium values of up to several hundred parts per million. These activities were measured both around open, oxygenated root channels and in near surface peat material. This analysis has suggested that uranium disperses seawards and in ground waters, and is retarded in this by organic material in the sediment [D4].

Sample	Location	Activity per fresh weight (mBq/g)	Activity ²³⁸ U per dry weight (mBq/g)
Seaweed	[D3] Sandyhills Bay [D3] UK		3.8 ± 0.1 3.8 to 18.6
Mussel	[D3] Sandyhills Bay [D3] UK		1.1 ± 0.1 1.01 to 37.1
Mollusc	[D3] Sandyhills Bay(winkle)[D3] UK[D2] UK (mollusc & winkle)[D2] Parton (winkle)	0.89 1.3	2.72 ± 0.01 1.36 to 18.9
Crab	[D2] UK [D2] Parton	0.046 0.052	
Lobster	[D2] UK [D2] Parton	0.035 0.028	

D.5 Literature values reported for ²³⁸U in seaweed and marine biota samples for the Solway Firth area are shown in the table above; also reported are estimated values of ²³⁸U from natural sources in aquatic foodstuff for the UK given by CEFAS [D2, D3]. Analysis results from Parton (near Whitehaven) are also included, although it should be noted that the uranium inventory at Parton is dominated by historical anthropogenic input of natural uranium from a local phosphate processing plant.

Annex D References

[D1]	Ragnarsdottir, V. Charlet, L. (2000) Uranium Behaviour in Natural Environments in: Environmental Mineralogy: Microbial Interactions, Anthropogenic Influences, Contaminated Land and Waste Management, Mineralogical Society Series Vol. 9, Mineralogical Society of Great Britain & Ireland.
[D2]	Radioactivity in Food and the Environment 13 (2007) Centre for Environment, Fisheries and Aquaculture Science ISSN 1365-6414.
[D3]	McDonald, P. Cook, G T. Baxter, M S. (1991) Natural and Artificial Radioactivity in Coastal Regions of UK. In: Radionuclides in the Study of Marine Processes, edited by Kershaw, P J and Woodhead, D S. Elsevier Applied Science, London.
[D4]	Milodowski, A E. et al. (1990) Uranium-mineralised Micro-organisms Associated with Uraniferous Hydrocarbons in Southwest Scotland. Nature, 347, 465.

Initial distribution

CESO(Army) SO2 RAD (2 hard copies plus electronic copy) House of Commons Library (Electronic copy) Head of Radioactive Waste - Environment Group, Scottish Government (8 copies, of which 6 copies are to be placed in the Scottish Government Library) Aaron Scarlett Dee Emerson Knowledge Services (Electronic copy)

This page is intentionally blank.

Report Documentation Form

A copy of this form is to be completed by the principal author for all Dstl reports. When complete, it is to be bound as the last numbered pages of the finished report.

1. Originators Report Number incl. Ve	ersion No	DSTL/TR52797 V1			
2. Report Protective Markings and an	y other markin	gs e.g. Caveats, Descript	ors, Privacy markings		
UNCLASSIFIED	UNCLASSIFIED				
3. Title of Report					
Depleted Uranium Survey Report: Kir	kcudbright Tra	aining Area 2010 Part 2 Ma	arine Environment		
4. Title Protective Markings incl. any	Caveats	UNCLASSIFIED			
5. Author Aaron Scarlett					
6. Originator's Name and Address		7. MOD Sponsor Name	and Address		
Aaron Scarlett		CESO(A) SO2 RAD			
DSTL Alverstoke		Headquarters Land Forces			
c/o Institute of Naval Medicine		IDL420, Ramillies Building			
Crescent Road		Marlborough Lines, Monks	ton Road		
Gosport		ANDOVER			
PO12 2DL		SP11 8HJ			
8. MOD Contract number and period	covered	701418			
9. Other Report Nos.					
10. Date of Issue	11. Paginatio	n	12. No. of References		
31 March 2011	40		22		
13. Abstract (A brief (approximately 1	50 words) fact	ual summary of the report)		
Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess any radiological impact of the firings on the terrestrial and marine environments and any associated risk to humans. This report presents the findings of the marine survey undertaken at KTA during 2010; the terrestrial survey is reported separately in Part 1. None of the samples analysed were radioactive within the meaning of the Radioactive Substances Act 1993, nor did they exceed a very small fraction of the Generalised Derived Limits advised by the Health Protection Agency (formerly the National Radiological Protection Board). In fact, the measured levels of uranium were found to be consistent with natural background levels. The results of the 2010 survey are consistent with historical survey findings. There is no evidence to indicate that members of the public are exposed to a radiological hazard from the marine environment as a result of test firing DU ammunition at Kirkcudbright.					
14. Abstract Protective Marking including any Caveats					
UNCLASSIFIED					
15. Keywords/Descriptors (Authors may provide terms or short phrases which identify concisely the technical concepts, platforms, systems etc. covered in the report.					
Kirkcudbright, Depleted Uranium, Radioactivity, Environmental, Survey, Marine.					

16. Report Announcement (refers to title/abstract being included in accessions lists e.g. Defence Reports Abstracts)					
Announcement of	this report is UNLIMITED				
If there are limitations can be announced (m	on the announcement of nore than one if required)	this report please indicate below the	groups to whom it		
Can be announce	d to MOD and its Agencie	S			
Can be announce	d to UK Defence Contract	tors			
Can be announce	d to Overseas Defence De	epartments			
Other (please spe	cify)				
17. Report Availability	/				
UNLIMITED distril	oution				
🛛 No Release witho	ut approval of the Release	e Authority (see box 7)			
If the above do not ap request <u>without</u> furthe	oply, please indicate below er Need-To-Know checks.	w the groups to whom the report may	be released upon		
Can be released t	o MOD and its Agencies				
Can be released t	o other UK Government D	Departments			
Can be released t	o UK Defence Contractors	S			
Can be released t	o Overseas Defence Depa	artments			
Other (please spe	cify)				
18. Downgrading Inst	ructions (check as approp	priate)			
This report may be	e automatically downgrad	ed to after years			
☐ This report may be	e reviewed years a	fter publication			
19. Authorisation					
	Name	Signature	Date		
Chief Scientist	Dr Adrian Baker	Addel	6/4/11		
Project Manager	Dee Emerson	Martin	61411		
Technical Reviewer	Ron Brown	RBrow	6/4/11		

This page is intentionally blank.