

UNCLASSIFIED



**Depleted Uranium Survey Report:
Kirkcudbright Training Area 2010
Part 1 Terrestrial Environment**

Aaron Scarlett

DSTL/TR51735 V1
31 March 2011

Environmental
Sciences Department
C/o INM
Crescent Road
Alverstoke
Hampshire
PO12 2DL

© Crown copyright 2011 Dstl

UNCLASSIFIED



Dstl is part of the
Ministry of Defence

UNCLASSIFIED




This document has been prepared for MOD and, unless indicated, may be used and circulated in accordance with the conditions of the Order under which it was supplied.

It may not be used or copied for any non-Governmental or commercial purpose without the written agreement of Dstl.

© Crown copyright 2011
Defence Science and Technology Laboratory UK

Approval for wider use or release must be sought from:

Intellectual Property Department
Defence Science and Technology Laboratory
Porton Down, Salisbury, Wiltshire, SP4 0JQ.

Authorisation			
	Name	Signature	Date
Chief Scientist	Dr Adrian Baker		6/4/11
Project Manager	Dee Emerson		6/4/11
Technical Reviewer	Ron Brown		6/4/11

UNCLASSIFIED

This page is intentionally blank

UNCLASSIFIED

Table of Contents

Executive Summary	3
List of Figures	6
List of Tables	6
1 Introduction	7
2 Background	7
3 Depleted Uranium (DU)	10
4 Differentiating DU From Natural Uranium	11
5 Reference Levels	12
6 Methodology	13
7 Results and Interpretation	18
8 Interpretation of Soil and Grass Isotopic Ratios	21
9 Evaluation of Potential Exposure Pathways	23
10 Conclusions	25
11 List of References	26
12 KTA Terrestrial Survey Results	28
Historical Data	33
ANNEX A Issues to be considered when interpreting or comparing uranium data	38
ANNEX B Change in the $^{238}\text{U}/^{234}\text{U}$ activity ratio of a medium containing natural uranium with the addition of depleted uranium	41
Initial Distribution	42

List of Figures

Figure 1. Number of DU projectiles fired at KTA between 1982 and 2010.	9
Figure 2. Approximate cumulative mass of DU projectiles fired at KTA between 1982 and 2010.	9
Figure 3. Terrestrial sampling locations at KTA 2010. Note: red markers indicate grass sampling; green markers indicate grass plus soil sampling.	14
Figure 4. Stream sampling locations - KTA 2010. Note: circular markings denote sediment sampling; square markings denote both sediment <i>and</i> water sampling.	16
Figure 5. Grass samples: isotopic ratios from alpha spectrometry analysis - KTA 2010.	22
Figure 6. Soil samples: isotopic ratios from alpha spectrometry analysis - KTA 2010.	22

List of Tables

Table 1. Approximate mass compositions of uranium isotopes in natural and depleted uranium.	10
Table 2. DUFERC investigation/action levels, RSA93 Schedule 1 activity concentrations and Generalised Derived Limits for uranium in soil and sediment.	12
Table 3. Summary of sample analyses - KTA 2010.	18
Table 4. Summary of soil, grass and animal indicator samples collected - KTA 2010.	28
Table 5. Summary of stream sediment and water samples collected - KTA 2010.	28
Table 6. Grass samples: alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.	29
Table 7. Soil samples: alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.	29
Table 8. Animal indicators (faeces): alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.	30
Table 9. Water samples: alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.	30
Table 10. Water samples (filter paper analysis): alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.	30
Table 11. Stream sediments: alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.	31
Table 12. Environmental gamma dose rates - KTA 2010.	31

1 Introduction

- 1.1 Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the extent of any radiological impact of the firings on the terrestrial and marine environments and any associated risk to humans [1 to13].
- 1.2 This report presents the findings of the terrestrial survey undertaken at KTA during 2010; the marine survey is reported separately in Part 2 [14]. The survey was undertaken to monitor the levels of any uranium in the terrestrial environment resulting from operations on the site and to identify the extent of any environmental transfer processes.
- 1.3 A review of historic environmental monitoring procedures was undertaken as part of an independent assessment of the DU firing at KTA (and Eskmeals) ranges in July 1995 [15]. During 1996, the environmental monitoring programme for KTA was revised in line with the recommendations of this assessment and the 1996 DU Baseline Survey report was published [16]. Environmental monitoring was undertaken in a consistent manner between 1996 and 2007.
- 1.4 Due to altered work activities at KTA (with only one battery-target combination now available for use and greatly reduced proof firing of DU munitions) the MOD and the Scottish Environment Protection Agency (SEPA) jointly agreed in 2007 that there should be greater emphasis on stream sediment sampling rather than soil sampling. The current terrestrial monitoring programme reflects this agreement and consists principally of the collection and analysis of grass and stream sediment/water samples, along with soil sampling around the active battery-target combination. Animal indicator samples are also collected where available. Further details of the revised sampling protocol are provided in the 2007 terrestrial survey report [11].

2 Background

- 2.1 The KTA range is located on the coast of Dumfries and Galloway, near Castle Douglas. In April 2006, the range became part of the Defence Training Estate (DTE).
- 2.2 DU has been released into the environment at KTA as a consequence of the test firing of DU ammunition during design and accuracy assessment trials. DU projectiles are fired through soft vertical targets and continue their trajectory, coming to rest in the Solway Firth.
- 2.3 Testing of projectiles historically has taken place at five locations on the KTA range. Strength of design trials were initially conducted at the Raeberry range using prototype ammunition. Functionality and accuracy trials were conducted until 2001 at the Balig and Doon Hill ranges, whilst confirmatory proving trials were carried out at

UNCLASSIFIED

the Silver Hill Low range. In 2003, rounds were fired from Challenger tanks placed at the Chapman 1000 metre firing point towards the India Target.

- 2.4 The current firing policy is to use the Balig Gun and India Target combination. There was no DU firing at KTA between 2003 and 2007, although a total of 20 rounds were fired in 2008 as part of a routine surveillance programme to ensure the safety and serviceability of the ammunition. No firing has taken place since 2008.
- 2.5 The number of DU rounds fired each year at KTA from the five firing locations and the cumulative mass of DU fired to date, are presented in Figure 1 and Figure 2 respectively.
- 2.6 DU penetrators do not fragment in air under normal circumstances. However, it was inevitable that some malfunctions would occur during the early test firings. Although the majority of malfunctioning penetrators still entered the Solway Firth, a small proportion impacted on land and some fragmented on exiting the gun barrel or in the early stage of flight. Whilst some of these fragments entered the sea, it is known that small quantities of particulate DU were deposited at a few discrete locations in the local terrestrial environment. Most of these locations are now fenced off, although the levels of DU present are below regulatory concern. The recovery of misfired DU penetrators has been attempted, although in most cases, penetrators are suspected to be buried at depth in the soil and therefore it has not been possible to locate or recover them.
- 2.7 An extensive radiological survey was carried out in 2002 using sensitive large area radiation monitoring equipment. Whilst identifying the expected isolated areas of low level DU contamination, the survey showed levels of radioactivity in most areas to be consistent with natural background levels. Most elevated radiation levels were due to naturally occurring radionuclides found in construction materials such as the granite chippings used on the range roads [17].

UNCLASSIFIED




Figure 1. Number of DU projectiles fired at KTA between 1982 and 2010.




Figure 2. Approximate cumulative mass of DU projectiles fired at KTA between 1982 and 2010.

3 Depleted Uranium (DU)

- 3.1 Uranium is a naturally occurring radioactive material which exists mainly as three isotopes: uranium-238 (^{238}U), uranium-235 (^{235}U) and uranium-234 (^{234}U). The typical mass composition of these isotopes is shown in Table 1. In natural uranium, ^{238}U and ^{235}U are in approximate radioactive equilibrium with their daughter products¹. Together these isotopes emit a range of alpha and beta particles along with gamma radiation. Being a heavy metal, the chemical toxicity of uranium is approximately equal to that of lead.
- 3.2 Uranium in an 'enriched' form is used as fuel in nuclear reactors. The enrichment process increases the concentration of ^{235}U (above 0.72%) in comparison to the natural form. The by-product of this process is 'depleted' uranium (DU), which has a reduced concentration of ^{235}U . Uranium-234 is also removed in the depletion process; meaning that DU is consequently less radioactive than natural uranium (the specific alpha activity of the DU fired at Kirkcudbright being approximately 1.4×10^7 milli becquerels per gram (mBq/g) compared to 2.5×10^7 mBq/g for natural uranium [18]. The mass compositions of DU and natural uranium are presented in Table 1 below.

Form of Uranium	^{238}U	^{235}U	^{234}U
Natural uranium	99.274%	0.72%	0.00554%
The DU used at Kirkcudbright	99.8%	0.20%	0.0008%

Table 1. Typical mass composition of uranium isotopes in natural and depleted uranium.

- 3.3 As discussed in paragraph 3.1, ^{234}U normally exists in approximate equilibrium with ^{238}U in the natural environment. In comparison, DU exhibits a $^{238}\text{U}/^{234}\text{U}$ activity ratio of between 7:1 and 8:1, dependent on the degree of depletion achievable by different methods of processing. This distinction is important in differentiating DU contamination from naturally occurring uranium in the environment (see Section 4). For the remainder of this report, isotopic ratios will be stated in terms of activity rather than mass and as a single value representing the ratio of becquerels of ^{238}U to 1 becquerel of ^{234}U (i.e. a $^{238}\text{U}/^{234}\text{U}$ ratio of 7 rather than 7:1).

¹A radioactive decay series occurs when a heavy radionuclide decays into successively lighter radionuclides known as daughter products. For example, ^{238}U decays to ^{234}Th , then $^{234\text{m}}\text{Pa}$, then ^{234}U and so on until a stable element is reached (^{206}Pb).

4 Differentiating DU From Natural Uranium

- 4.1 The fundamental requirement of the DU environmental monitoring programme is to quantify the radiological impact of DU firing. This is achieved partly by measuring the amount of total uranium in environmental materials and using this figure as an upper bound of DU contamination levels. However, as uranium is present at detectable levels in most environmental materials, this overestimates the risk. More sophisticated analyses involve the specific measurement of ^{238}U and ^{234}U isotopes (by activity and/or mass). Although isotope measurements are used in this survey, references to total uranium measurements are included for consistency with historic reports. The limitations of using total uranium concentrations are discussed further in Annex A.
- 4.2 A convenient fingerprint marker for DU contamination is the $^{238}\text{U}/^{234}\text{U}$ activity ratio. The DU fired at KTA has a $^{238}\text{U}/^{234}\text{U}$ activity ratio of approximately 7, whereas natural uranium in the environment typically has an activity ratio close to unity. Environmental samples are therefore analysed for isotopes of ^{238}U and ^{234}U to determine activity ratios and hence identify the origin of the uranium.
- 4.3 Substantial deposition of DU in the terrestrial environment (in addition to an existing natural uranium background) is required before the $^{238}\text{U}/^{234}\text{U}$ activity ratio diverges significantly from its natural ratio. An illustration of the impact of DU contamination on the isotopic ratio is given in Annex B. For the ratio to approach 7 in an analytical sample, the mass of DU would have to be approximately one hundred times the mass of the uranium that is naturally present. Hence, the lower the natural uranium background, the lower the levels of DU contamination that may be detected by isotopic analysis.
- 4.4 Isotopic quantification is achieved by techniques such as alpha spectrometry and mass spectrometry. Alpha spectrometry can detect uranium to parts per billion, which is equivalent to mBq per kg, or to lower levels if count times are increased. Mass spectrometry is more sensitive, but the lower levels detectable are of no recognised health significance. Isotopic information can also be yielded from gamma spectrometry analyses, although limits of detection are not generally sufficient for measurement of environmental levels.

5 Reference Levels

- 5.1 The Depleted Uranium Firing Environmental Review Committee (DUFERC), on which the MOD is represented, has agreed investigation/action levels for DU in soil². These are currently based on, and represent a small fraction of, the Generalised Derived Limits (GDLs) advised by the Health Protection Agency (formerly the National Radiological Protection Board) [19] and the Schedule 1 activity concentration for uranium laid down in the Radioactive Substances Act 1993 (RSA93) [20]. GDLs for uranium were last updated in 2000 and were referred to by the Royal Society in their studies of the potential health effects of using DU munitions [21]. The investigation/action levels are set intentionally low to ensure that any DU released into the environment is identified before it can accumulate to significant levels. In particular, action levels are set to less than 10% of the level at which control would be required under RSA 93.

Source	Reference Level	Activity concentration (mBq/g dry weight)
DUFERC	Investigation Level	300
DUFERC	Action Level	1,110
RSA 93	Level at which regulatory control is required.	11,100
NRPB (2000)	Generalised Derived Limit: ²³⁸ U in well-mixed soil	20,000
NRPB (2000)	Generalised Derived Limit: ²³⁸ U in freshwater sediment	400,000

Table 2. DUFERC investigation/action levels, RSA93 Schedule 1 activity concentrations and Generalised Derived Limits for uranium in soil and sediment.

- 5.2 There are no DUFERC agreed investigation/action levels for uranium in water. The alpha spectrometry system deployed by Dstl can detect uranium in water at levels down to about 10% of the World Health Organisation's (WHO) most restrictive recommendation for uranium levels in drinking water (2 µg per litre, which relates to approximately 50 mBq/l for natural uranium). This 10% level is commonly used as a 'trigger' in occupational health monitoring.

² Soil in areas of contamination above DUFERC investigation/action levels shall be managed in accordance with the KTA Depleted Uranium Management and Remediation Plan [22].

6 Methodology

- 6.1 As discussed in Section One, the current survey methodology consists primarily of the collection and analysis of grass and stream sediment/water samples, together with soil sampling around the active battery-target combination. Animal indicator samples are also collected where available and environmental gamma dose rates are recorded at each sampling location. Stream sediment/water sample points are located at areas where any potential contamination is likely to leach from the surrounding soil.
- 6.2 Full details of the methodology are provided in the following paragraphs. A list of the soil, grass and faecal samples collected is given in Table 4. A list of stream sediment and water samples collected is given in Table 5.

Terrestrial sampling sites

- 6.3 For each gun-target combination, sampling sites were chosen in the vicinity of the gun position, at the soft target stand, and at the mid-point of the range. In addition, samples collected from the Gypsy Point location were used for the purpose of background comparison. Grass samples were collected from all locations whereas soil samples were only collected from locations 4, 5 and 6 (Balig-India combination) and the background location (K13). The 13 sampling points are shown in Figure 3; the location names and Ordnance Survey of Great Britain grid references are provided below:

1.	Raeberry Target	NX 70449 43744
2.	Raeberry Bunker Midpoint	NX 70472 43836
3.	Raeberry Gun	NX 70521 43980
4.	India Target	NX 70631 43658
5.	Balig Gun/Target waypoint	NX 70964 44498
6.	Balig Gun	NX 71238 45452
7.	Zulu Gantry	NX 71848 43548
8.	Mullock Farm	NX 71188 44295
9.	Silver Hill (Low) Gun	NX 70355 44851
10.	Echo Target (Doon Hill)	NX 72291 43807
11.	Doonhill/Target waypoint	NX 72096 45429
12.	Doon Hill Gun	NX 71904 46947
13.	Gypsy Point	NX 68676 43789




Figure 3. Terrestrial sampling locations at KTA 2010. Note: red markers indicate grass sampling; green markers indicate grass plus soil sampling.

- 6.4 At each sampling site, grass samples were collected from three areas of approximately 1 m^2 in size located within 5 metres of each other. Where the grass was scarce or short, the sampling area was increased until the samples obtained were of the requisite mass for laboratory analysis (greater than 200 g). The grass was cut at a height of at least 2 cm above the ground to avoid including soil in the sample. The type of grass collected and the content of other plant species varied from site to site. The 3 unwashed grass samples from each site were combined into one composite sample and analysed by alpha spectrometry.
- 6.5 Soil samples were collected as undisturbed cores of 4.8 cm diameter and up to 30 cm in depth from the centre of each of the three 1 m^2 grass sampling areas. Each soil core was divided into sub-samples of two depth intervals (0-2 cm and 2-5 cm) in the laboratory and portions of the three sub-samples from each point combined to create

UNCLASSIFIED

one composite sample for each depth interval (0-2 cm and 2-5 cm). This system yielded a total of 8 composite samples from the 4 soil sample locations. These were analysed by alpha spectrometry to provide an indication of distribution with depth.

- 6.6 At each grass sampling area, gamma dose rates were derived from measurements taken using a Mini-Instruments 6-80/81 and a compensated MC71 Geiger-Muller tube set up at 1 m above the ground and allowed to record over a period of 300 seconds. A mean gamma dose rate was calculated from three replicate measurements using the methodology described in Reference [23].
- 6.7 Where available, samples of animal faeces (e.g. deer, rabbit, fox, badger, sheep and cattle) were collected close to the sampling sites. Only fresh samples were collected, taking care to minimise the amount of soil or grass that could inadvertently be sampled at the same time. Rabbit droppings were collected until enough material had been sampled, making one composite sample for that location. These samples were analysed by alpha spectrometry.
- 6.8 Deer culling takes place on the range to manage the deer population and biological samples (liver and/or kidneys) are routinely collected from the culled deer on an opportunistic basis. No biological samples were available for analysis in 2010 as no deer cull was conducted by site management. However, historical sample analysis has shown that levels of total uranium are very low and do not indicate any DU contamination.

Stream sediment and water sampling

- 6.9 Stream sediment was collected at fifteen locations from streams that collect run-off water from the soil surface of the range. Stream water samples were collected at five locations. The Ordnance Survey of Great Britain grid references for the fifteen stream sediment and water sampling sites used in 2010 are given below. The exact position of the sampling sites changes slightly from year to year due to changing stream conditions and access. The current locations are shown on a map of the site in Figure 4.

Stream sediments sampling grid references:

S1.	Burnfoot Bridge	NX 74199 44570
S2.	Netherlaw Wood	NX 74166 44631
S3.	Netherlaw Burn,	NX 73438 44779
S4.	Quatercake Burn	NX 72326 44363
S5.	Quatercake Burn	NX 71804 43917
S6.	Brandy Burn	NX 71264 44145
S7.	Dunrod & Overlaw Burns	NX 70820 43732
S8.	Dunrod Burn	NX 70958 44825
S9.	Dunrod Burn	NX 71082 45571
S10.	Ring Burn	NX 71152 45961
S11.	Overlaw Burn	NX 71166 44847
S12.	Overlaw Burn	NX 72120 46277
S13.	Overlaw Burn	NX 72507 46920
S14.	Balmae Burn	NX 69144 44643

S15. Balmae Burn NX 68556 43890

Stream water sampling grid references:

W1. Abbey Burn NX 74199 44570

W2. Netherlaw Burn NX 74202 44594

W3. Balmae Burn(Gypsy Point)NX 68556 43890

W4. Dunrod & Overlaw Burns NX 70820 43732

W5. Quartercake Burn NX 71804 43917




Figure 4. Stream sampling locations - KTA 2010. Note: circular markings denote sediment sampling; square markings denote both sediment *and* water sampling.

6.10 Stream sediment was collected from below the water where possible. Care was taken to sample undisturbed sediments which had not been disturbed by cattle, for example. The top layer of the sediment (up to 5 cm depth) was collected and any large stones were removed. Sample sizes ranged from 0.4 to 0.8 litres. The sediment samples were analysed by alpha spectrometry.

6.11 Five water samples (0.5 litre) were collected from watercourses that run through the range. Samples were collected at accessible locations along the stream (e.g. from bridges and easily accessible banks). To remove any suspended sediment, the water

UNCLASSIFIED

was filtered through a Whatman filter paper using a funnel previously rinsed in stream water into a rinsed sample bottle. Both the water and the filter paper were analysed by alpha spectrometry.

Sample Descriptors

- 6.12 Each sample was given a unique sample descriptor. Those samples collected from specific sites (e.g. grass samples) were given a prefix such as K5 ('K' for Kirkcudbright followed by the location number). This prefix was followed by 'S' for soil sample; a 'G' for grass samples or the type of faecal sample. Soil samples were also denoted with a 'U' or a 'L' for the upper and lower soil horizon composite samples respectively. Deer kidney and liver samples were simply named 'Deer kidney (a) or (b)' and water/sediment samples were given descriptors such as W3 and S13 respectively. Some examples of sample descriptors are provided below:

K5/S/U : Composite soil sample, upper horizon, collected from location 5

K8/G : Composite grass sample collected from location 8

K3 rabbit : Rabbit faecal samples collected from location 3

W2 : Water sample collected from water sample location 2

S12 : Stream sediment sample collected from stream location 12

Sample preparation and laboratory analysis

- 6.13 The samples were prepared and analysed by alpha spectrometry in the Dstl UKAS accredited radiochemistry laboratory following the procedure adopted for the analysis of the terrestrial samples during the 1996 Baseline Survey [16]. An outline of the approach is given below.
- 6.14 The solid samples (soil, grass and biological indicators) were dried to remove moisture and weighed (with results being reported as dry weight). The samples were ashed to remove organic material and homogenised. During these processes, the loss of uranium from the sample is deemed to be insignificant. The ashed samples were boiled in concentrated mineral acid (nitric acid and hydrochloric acid) to remove the 'loose' and leachable uranium from the sample. Recalcitrant matrices such as mineral grains were not broken down by the process and hence natural uranium bound up within them was not removed. The samples were filtered to remove solids. Water samples of 250 ml were boiled down to approximately 100 ml and acidified.
- 6.15 Uranium separation was carried out by extraction chromatography. Each eluted sample was electro-deposited onto a stainless steel planchette and the activity of each planchette was counted in a low background alpha spectrometer with a silicon surface barrier.
- 6.16 Uranium activity concentrations are reported in mBq/g (equivalent to Bq/kg) of dry weight for soils, grass and biological indicators and mBq/l for water samples.

7 Results and Interpretation

- 7.1 A summary of the results for all terrestrial samples collected in 2010 is given below in Table 3. The full terrestrial monitoring results are provided in Tables 6 to 12 in Section 12. Historical monitoring results for the KTA for the years 1996 to 2010 are presented in Historical Data section.
- 7.2 It should be noted that the summary below is provided as an overview of the data collected. Given that some samples were collected from specific areas of known contamination, the mean values should not be viewed as an indicator of average uranium concentrations across KTA as a whole.

Sample type	Number of samples	Number of samples containing detectable DU ³	Total uranium concentration (mBq/g or mBq/l)			
			Mean	Standard deviation of the mean	Minimum	Maximum
Soil upper	4	1	34.97	7.87	24.56	43.44
Soil lower	4	0	39.34	6.23	31.01	45.86
Grass	13	2	0.48	0.70	0.05	2.49
Water	5	0	4.06	2.35	1.83	7.85
Stream sediment	15	0	26.76	4.67	20.80	35.45
Faeces (Cow)	3	0	1.00	1.03	0.16	2.15
Faeces (Sheep)	3	0	0.39	0.18	0.18	0.50

Table 3. Summary of sample analyses - KTA 2010.

³ Samples are reported as containing detectable DU if the ratio of ²³⁸U to ²³⁴U (after subtraction of the associated uncertainty to give the 95% confidence level) is greater than 1.0 for soil, grass and biota samples.

UNCLASSIFIED

Grass sample analysis

- 7.3 Alpha spectrometry results for grass samples are shown in Table 6. The total uranium levels ranged from 0.05 ± 0.02 to 2.49 ± 0.31 mBq/g; the highest result originating from the enclosed Raeberry Target area (K01). The background level at Gypsy Point (K13) was found to be 0.35 ± 0.09 mBq/g.
- 7.4 The $^{238}\text{U}:$ ^{234}U ratio of two samples (K01 and K03) is indicative of depleted uranium (maximum 5.06 ± 1.66). However, the total uranium levels are consistent with the typical range for grasses in the UK (0.2 to 3.8 mBq/g [24]) and no sample was radioactive within the meaning of the RSA93.

Soil sample analysis

- 7.5 Alpha spectrometry results for soil samples are shown in Table 7. Levels of total uranium ranged from 34.55 ± 3.80 to 45.86 ± 4.88 mBq/g; this being consistent with the findings of previous surveys [1 to 13] and the levels found at the background site (Gypsy Point: 24.56 ± 3.10 to 31.01 ± 3.61 mBq/g). Results are also consistent with those reported in the literature for UK soil, where an upper bound of total uranium is estimated at 50 mBq/g [21]⁴.
- 7.6 Although there was some evidence of minor depletion at Balig Gun (K4) (isotopic ratio 1.51 ± 0.34), no sample was radioactive within the meaning of RSA93 and all samples were well below the GDL for well mixed soil (20,000 mBq/g). Furthermore, evidence of some minor depletion at this area is consistent with the findings of previous surveys [1 to 13].

Animal indicator sample analysis

- 7.7 Alpha spectrometry results for animal faeces samples are shown in Table 8. The levels of total uranium in faecal samples ranged from 0.16 ± 0.07 to 2.15 ± 0.37 mBq/g. The maximum background level recorded at Gypsy Point was 0.18 ± 0.06 mBq/g. Similar levels of uranium have been found in faecal samples in the past (see Historical Data Section) and it is emphasised that isotopic ratios indicate that the uranium is natural in origin.

Stream water sample analysis

- 7.8 Alpha spectrometry results for stream water and filter paper samples are presented in Table 9 and 10 respectively. Total uranium in water samples ranged from 1.83 ± 1.36 to 7.85 ± 2.96 mBq/l. All sample results were below the WHO drinking water limit of 2 µg/l, which is equivalent to approximately 50 mBq/l (natural uranium). Where it was possible to calculate an isotopic ratio from the low levels present, this was indicative of natural uranium.

⁴ Soil minerals containing uranium are widely distributed on the surface of the Earth's crust and the concentrations of natural uranium in the terrestrial environment can vary significantly between locations. Consequently, there is no single definitive reference level for natural uranium in soils. However, there is broad agreement in the range of values published in the literature: typical values in the UK range from 2 to 50 mBq/g wet weight [21], but concentrations of up to 100 times the typical range have been found in some locations.

UNCLASSIFIED

- 7.9 Analysis of filter paper samples, which represent the suspended particulates in the water, showed that total uranium levels ranged from 0.53 ± 0.38 to 1.24 ± 0.57 mBq/l. This is well below the WHO drinking water guidelines (50 mBq/l). The levels detected were so low that it was not possible to calculate isotopic ratios.

Stream sediment sample analysis

- 7.10 Alpha spectrometry results for stream sediment samples are presented in Table 11. The levels of total uranium present ranged from 20.80 ± 2.73 to 35.45 ± 4.13 mBq/g, with isotopic ratios indicating that the uranium present was natural in origin. No sample was radioactive within the meaning of RSA93 and all samples were well below the GDL for freshwater sediments (400,000 mBq/g).

Environmental dose rate measurements

- 7.11 Environmental gamma dose rate measurements recorded during 2010 are shown in Table 12 and range from 0.084 to 0.106 micro grays per hour ($\mu\text{Gy/h}$). The background measurement at Gypsy Point was recorded at $0.100 \mu\text{Gy/h}$. These measurements are consistent with those found during previous surveys [1 to 13].

8 Interpretation of Soil and Grass Isotopic Ratios

- 8.1 This section provides an assessment of the $^{238}\text{U}/^{234}\text{U}$ isotopic ratios within samples to ascertain the degree of depletion and hence the extent of DU contamination. Further discussion on the use of isotopic ratios to differentiate DU from natural uranium is given in Section 4. An illustration of the impact of increasing the DU mass in a sample on the isotopic ratio is provided in Annex B.
- 8.2 Provided below are graphical interpretations of grass and soil isotopic ratios together with an explanation of the findings. As no isotopic ratios significantly above unity were identified in other sampling types, graphs for animal indicators and stream sediment/water samples are omitted.

Isotopic ratios in grass samples

- 8.3 A graphical summary of the isotopic ratios for the grass samples collected in 2010 is provided in Figure 5. Where no isotopic value is shown in the graph, uranium isotopes have not been detected above the limits of detection.
- 8.4 Due to the low levels of uranium present, isotopic ratios could only be determined for 6 out of 13 grass samples. Most samples exhibited isotopic ratios indicative of natural uranium. However, samples from Raeberry Target (K01) and Raeberry Gun (K03) had isotopic ratios of 4.88 ± 1.32 and 5.06 ± 1.66 respectively. Although these ratios are indicative of DU contamination, it should be noted that the actual levels of uranium are consistent with typical background levels. Furthermore, these samples are taken from areas of known low level contamination resulting from past operations on site.

Isotopic ratios in soil samples

- 8.5 A graphical summary of the isotopic ratios for the soil samples analysed in 2010 is shown in Figure 6. Individual isotopic ratios ranged from 0.98 ± 0.23 to 1.51 ± 0.34 ; the highest ratio being from India Target (K04). The maximum isotopic ratio from the background location (Gypsy Point) was found to be 1.00 ± 0.23 . These results are consistent with previous findings and it is emphasised that the actual levels of uranium present are consistent with typical background levels.




Figure 5. Grass samples: isotopic ratios from alpha spectrometry analysis - KTA 2010.




Figure 6. Soil samples: isotopic ratios from alpha spectrometry analysis - KTA 2010.

9 Evaluation of Potential Exposure Pathways

9.1 Any contamination of the terrestrial environment with DU results in five potential exposure pathways for humans, as described below:

- External exposure from contaminated vegetation/soil or from DU fragments;
- Inhalation of re-suspended DU contamination;
- Ingestion of crops or animal products from DU contaminated pasture or soil;
- Ingestion of DU contaminated water from streams; and
- (Inadvertent) ingestion of DU contaminated soil or sediment.

9.2 As discussed in Section 7, the total uranium activity results for all samples represented a small fraction of the relevant GDLs and WHO limit for drinking water. Therefore, any associated radiation doses are deemed to be insignificant. For completeness, however, the potential doses from the highest level of uranium found in grass and soil samples are discussed below, together with a more general discussion of potential doses on the range as a whole.

External radiation exposure

9.3 Environmental gamma dose rates recorded across KTA during 2010 were consistent with natural background radiation levels. This is also the case where low levels of DU contamination have been found historically (e.g. Raeberry Target) and indicates that there is no increased risk from external exposure related to DU at KTA.

Inhalation of re-suspended DU

9.4 Any DU which has been deposited on soil, river sediment and vegetation may be re-suspended into the air and subsequently inhaled by range staff or members of the public. The risk associated with re-suspended DU would be greatest during maintenance work on site such as fence maintenance and strimming of vegetation. However, the levels of DU found in soil during 2010 are well below the GDL for well-mixed soil (20,000 mBq/g) which itself relates to a Committed Effective Dose of 1 mSv per annum (the current UK dose limit for members of the public). The maximum level of uranium found in soil during 2010 was at sample location K5, where total uranium levels were 45.86 ± 4.88 mBq/g (at 2 to 5 cm depth). Given that the level of uranium is consistent with natural levels reported in the literature (up to approximately 50 mBq/g), any radiological dose due to inhalation would be indistinguishable from natural background exposure.

Ingestion of DU contaminated foodstuffs

9.5 No agricultural crops intended for human consumption are grown on the range so the potential exposure route involving the ingestion of contaminated crops need not be considered further. However, as a small number of cattle, sheep and deer graze on the

UNCLASSIFIED

range, consideration must be given to the exposure pathway of ingestion of animal products derived from contaminated soil or pasture. It should be noted, however, that the known areas of contamination on site are enclosed within fenced areas and represent a small fraction of the total range area; any potential dose estimates therefore represent a worst case scenario.

- 9.6 No deer kidney or liver samples could be collected in 2010 as no deer cull took place. However, uranium levels in kidney and liver samples analysed during previous surveys have been very low. Any potential doses through this exposure pathway are therefore deemed to be indistinguishable from natural background exposure.

(Inadvertent) ingestion of DU contaminated soil or stream sediment

- 9.7 While there is a possibility that trace amounts of soil or stream sediment could be inadvertently ingested by those who come into contact with the material, the levels detected during 2010 are consistent with previous year's results and do not indicate any risk through this exposure pathway (they are well below the relevant GDLs). Furthermore, it should be noted that known areas of contamination on site are situated within fenced areas and access is strictly controlled.

Radiation exposure of representative person

- 9.8 The 'representative person' is a hypothetical individual who is deemed to receive the greatest radiation exposure from DU due to the critical pathway (e.g. external exposure and inhalation of re-suspended material). Two representative individuals have been identified for the terrestrial environment at KTA, as discussed below.
- 9.9 Workers employed to inspect and maintain the boundary fence around the known contaminated areas are likely to be at most risk due to their occasional presence and proximity to the contaminated areas. A representative person from this group of people may receive a radiation dose via external exposure and the inhalation of re-suspended DU contamination from soil or vegetation. They may also receive further exposure through the inadvertent ingestion of trace amounts of contaminated soil. However, since the findings of the 2010 survey indicate that total uranium levels are consistent with typical background levels, the dose implications are considered to be indistinguishable from natural background exposure.
- 9.10 For information, using the methodology provided in the HPA document W36 (Reference 25), the potential dose to a full time park worker exposed to disturbed patchy contamination at 1,100 mBq/g (the DUFERC Action Level) is estimated to be 1.4 µSv per annum. This represents just 0.1 % of the UK dose limit for members of the public.
- 9.11 The second representative person is a local individual who consumes meat from the wild animals that roam on the range. Although no deer samples could be obtained in 2010, the results of previous surveys indicate that the dose from this pathway would be indistinguishable from natural background exposure.

10 Conclusions

- 10.1 The 2010 Kirkcudbright terrestrial monitoring programme was undertaken to assess the levels of DU in the environment resulting from the firing of DU munitions. The monitoring programme consisted principally of the collection and analysis of grass and stream sediment/water samples, along with soil and grass sampling around the active battery-target combination. Animal indicator samples were also collected.
- 10.2 No sample was radioactive within the meaning of RSA93, nor did it exceed a small fraction of the relevant GDL. All samples contained levels of uranium which are consistent with those expected due to naturally occurring uranium. The survey results indicate that the existing areas of low-level DU contamination on site, which are maintained within fenced compounds, are not leading to the transfer of uranium to other parts of the environment.
- 10.3 Based on the findings of this survey – which are consistent with those of previous surveys – potential doses to representative persons, site personnel and members of the public are deemed to be consistent with natural background exposure.

11 List of References

1. Armstrong, V. (1999) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area - 1997. DRPS Report No 10/99.
2. Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area 1998. DRPS Report No 52/99.
3. Gooding, M. (2000) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area - 1999. DRPS Report No 18/00.
4. Liddle, S. and Mayes, D E. (2004) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area - 2000. DRPS Report No CR 11882.
5. Liddle, S. and Mayes, D E. (2004) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area - 2001. Dstl report CR 11888.
6. Liddle, S. and Mayes, D E. (2004) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area - 2002. Dstl report CR 12350.
7. Toqué, C. (2005) Terrestrial Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2003. Dstl Report CR 13038.
8. Toqué, C. (2006) Terrestrial Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2004. Dstl Report CR 17843.
9. Toqué, C. (2006) Terrestrial Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2005. Dstl Report CR19578.
10. Toqué, C. (2007) Terrestrial Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2006. Dstl Report CR24924.
11. Toqué, C. (2008) Terrestrial Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2007. Dstl Report DSTL/CR30510.
12. Scarlett, A. (2010) Depleted Uranium Survey Report: Kirkcudbright Training Area 2008 Part 1 Terrestrial Environment DSTL/TR37166 V1.
13. Scarlett, A. (2010) Depleted Uranium Survey Report: Kirkcudbright Training Area 2009 Part 1 Terrestrial Environment DSTL/TR40942 V1.
14. Scarlett, A. (2010) Depleted Uranium Survey Report: Kirkcudbright Training Area 2010 Part 2 Marine Environment DSTL/TR52797 V1.

UNCLASSIFIED

15. Environmental Assessment of the Firing of Depleted Uranium Projectiles at Eskmeals and Kirkcudbright Ranges, July 1995. WS Atkins Consultants Ltd.
16. Gibbs, K J. (1996) Environmental Depleted Uranium Baseline Survey Report DTEO Kirkcudbright. DERA Radiation Protection Services, DRPS 6/97 (1997).
17. Gooding, M. (2003) Radiological Survey of the Kirkcudbright Training Area. Dstl Radiological Protection Services report 224/2002.
18. Carter, T. (2002) Comparison of Kirkcudbright and Eskmeals Environmental Monitoring Data with Generalised Derived Limits for Uranium. Dstl 167/2002.
19. Generalised Derived Limits for Radioisotopes of Polonium, Lead, Radium and Uranium. Documents of the NRPB, Vol. 11, No 2 (2000).
20. Radioactive Substances Act 1993. HMSO (ISBN 0-10-54 1293-7).
21. The Health Hazards of Depleted Uranium Munitions (Part 2) The Royal Society (2002) ISBN 0854035745.
22. Kirkcudbright Training Area Depleted Uranium Management and Remediation Plan (2010) [IN DRAFT].
23. Her Majesty's Inspectorate of Pollution (1995) Technical Guidance Note (Monitoring) Routine Measurement of Gamma Ray Air Kerma Rate in the Environment. HMSO ISBN 0 11 753132 4.
24. Jenkins, D W K. Sandalls, F J. Hill, R J. (1989) Uranium and Thorium in British Crops and Arable Grasses. AERE-R13442. HMSO, London.
25. National Radiological Protection Board (2003) W-36 Methodology for Estimating the Doses to Members of the Public from the Future use of Land Previously Contaminated with Radioactivity.

12 KTA Terrestrial Survey Results

Sample station number	Sample type and number of samples			Environmental dose rates
	Soil	Grass	Animal indicators (faeces)	
K1	×	1	×	✓
K2	×	1	×	✓
K3	×	1	×	✓
K4	1	1	×	✓
K5	1	1	×	✓
K6	1	1	1	✓
K7	×	1	×	✓
K8	×	1	1	✓
K9	×	1	×	✓
K10	×	1	1	✓
K11	×	1	1	✓
K12	×	1	×	✓
K13	1	1	2	✓

Table 4. Summary of soil, grass and animal indicator samples collected - KTA 2010.

Note: × denotes sample not collected.

Sediment sample number	Water sample number	Location name	Burn
S1	W1 & W2	Burnfoot Bridge	Abbey and Netherlaw
S2	×	Netherlaw Wood	Netherlaw
S3	×	Cross roads	Netherlaw
S4	×	Craigraploch	Quatercake
S5	W5	Downstream	Quatercake
S6	×	Mullock Farm	Brandy
S7	W4	Mullock Bay	Dunrod and Overlaw
S8	×	Upstream	Dunrod
S9	×	Balig Gun turn off	Dunrod
S10	×	Dunrod Mill gate	Ring
S11	×	Overlaw Burn	Overlaw
S12	×	Bailey Bridge	Overlaw
S13	×	EM Gun	Overlaw
S14	×	Gypsy Barrier	Balmae
S15	W3	Gypsy Point	Balmae

Table 5. Summary of stream sediment and water samples collected - KTA 2010.

UNCLASSIFIED

Sample descriptor	Fresh weight (g)	Dry weight (g)	Ashed weight (g)	Measured activity of dry sample (mBq/g)				²³⁸ U/ ²³⁴ U ratio
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
K01	322.8	135.7	8.7	2.04 ± 0.30	< 0.04	0.42 ± 0.10	2.49 ± 0.31	4.88 ± 1.32
K02	313.8	85.2	6.7	< 0.11	< 0.11	< 0.11	0.08 ± 0.06	N/A
K03	363.7	136.3	9.7	0.99 ± 0.16	< 0.03	0.20 ± 0.06	1.20 ± 0.17	5.06 ± 1.66
K04	308.3	136.0	8.6	0.07 ± 0.04	< 0.04	< 0.04	0.10 ± 0.05	N/A
K05	353.3	127.8	6.9	< 0.02	< 0.02	0.02 ± 0.02	0.05 ± 0.02	N/A
K06	399.7	117.3	8.6	< 0.03	< 0.03	< 0.03	0.05 ± 0.03	N/A
K07	256.5	130.8	7.7	0.09 ± 0.03	< 0.02	0.10 ± 0.03	0.20 ± 0.05	0.84 ± 0.39
K08	398.6	181.6	9.3	< 0.02	< 0.02	0.03 ± 0.02	0.05 ± 0.02	N/A
K09	359.0	148.8	9.0	0.33 ± 0.10	< 0.05	< 0.05	0.37 ± 0.10	N/A
K10	343.6	159.7	15.6	0.41 ± 0.11	< 0.05	0.42 ± 0.11	0.86 ± 0.16	0.97 ± 0.37
K11	438.9	165.7	7.6	< 0.03	< 0.03	0.06 ± 0.03	0.09 ± 0.04	N/A
K12	477.2	154.4	8.3	0.17 ± 0.04	< 0.02	0.18 ± 0.05	0.35 ± 0.06	0.92 ± 0.34
K13	596.9	229.0	16.4	0.15 ± 0.06	< 0.04	0.19 ± 0.06	0.35 ± 0.09	0.77 ± 0.38

Table 6. Grass samples: alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.

Sample descriptor	Fresh weight (g)	Dry weight (g)	Ashed weight (g)	Measured activity of dry sample (mBq/g)				²³⁸ U/ ²³⁴ U ratio
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
K4/S/U	34.7	25.9	15.6	20.35 ± 3.06	0.76 ± 0.38	13.44 ± 2.22	34.55 ± 3.80	1.51 ± 0.34
K4/S/L	105.0	84.6	67.5	22.05 ± 3.36	1.16 ± 0.51	18.22 ± 2.89	41.42 ± 4.47	1.21 ± 0.27
K5/S/U	63.8	52.9	41.6	18.16 ± 2.97	0.75 ± 0.43	18.45 ± 3.01	37.35 ± 4.25	0.98 ± 0.23
K5/S/L	120.9	89.3	79.4	22.40 ± 3.42	0.95 ± 0.47	22.51 ± 3.44	45.86 ± 4.88	0.99 ± 0.21
K6/S/U	78.2	52.9	41.6	22.38 ± 3.52	0.76 ± 0.44	20.30 ± 3.26	43.44 ± 4.82	1.10 ± 0.25
K6/S/L	179.8	128.6	109.3	20.99 ± 3.13	< 0.40	17.72 ± 2.74	39.07 ± 4.16	1.18 ± 0.25
K13/S/U	84.0	57.9	42.0	11.44 ± 2.10	0.53 ± 0.35	12.58 ± 2.25	24.56 ± 3.10	0.91 ± 0.23
K13/S/L	163.5	132.2	115.4	15.39 ± 2.54	< 0.47	15.46 ± 2.55	31.01 ± 3.61	1.00 ± 0.23

Table 7. Soil samples: alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.

UNCLASSIFIED

Sample descriptor	Fresh weight (g)	Dry weight (g)	Ashed weight (g)	Measured activity of dry sample (mBq/g)				²³⁸ U/ ²³⁴ U ratio
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
K10 (cow)	59.0	25.3	2.4	0.25 ± 0.10	< 0.07	0.41 ± 0.13	0.68 ± 0.16	0.60 ± 0.30
K11 (cow)	45.2	21.1	3.0	0.88 ± 0.23	< 0.11	1.21 ± 0.28	2.15 ± 0.37	0.73 ± 0.26
K13 (cow)	70.9	17.2	1.5	0.08 ± 0.05	< 0.05	0.07 ± 0.04	0.16 ± 0.07	1.09 ± 0.92
K06 (sheep)	62.3	26.8	3.1	0.24 ± 0.12	< 0.13	0.22 ± 0.12	0.50 ± 0.18	1.09 ± 0.80
K08 (sheep)	49.6	19.5	2.2	0.18 ± 0.08	< 0.07	0.27 ± 0.10	0.48 ± 0.13	0.65 ± 0.38
K13 (sheep)	77.0	22.6	2.1	0.08 ± 0.04	< 0.05	0.09 ± 0.05	0.18 ± 0.06	0.89 ± 0.66

Table 8. Animal indicators (faeces): alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.

Sample descriptor	Measured activity of sample (mBq/l)				²³⁸ U/ ²³⁴ U ratio
	²³⁸ U	²³⁵ U	²³⁴ U	Total U	
W1	< 2.50	< 2.50	< 2.50	4.14 ± 2.20	N/A
W2	< 2.21	< 2.21	< 2.21	2.39 ± 1.57	N/A
W3	2.94 ± 1.81	< 2.35	4.35 ± 2.22	7.85 ± 2.96	0.68 ± 0.54
W4	< 2.17	< 2.17	2.39 ± 1.56	4.08 ± 2.03	N/A
W5	< 2.19	< 2.19	< 2.19	1.83 ± 1.36	N/A

Table 9. Water samples: alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.

Sample descriptor	Measured activity of sample (mBq/l)				²³⁸ U/ ²³⁴ U ratio
	²³⁸ U	²³⁵ U	²³⁴ U	Total U	
W1 (filter)	< 0.60	< 0.60	< 0.60	0.86 ± 0.49	N/A
W2 (filter)	< 0.60	< 0.60	0.77 ± 0.47	1.13 ± 0.57	N/A
W3 (filter)	< 0.57	< 0.57	< 0.57	0.69 ± 0.43	N/A
W4 (filter)	< 0.59	< 0.59	< 0.59	0.53 ± 0.38	N/A
W5 (filter)	< 0.55	< 0.55	0.94 ± 0.50	1.24 ± 0.57	N/A

Table 10. Water samples (filter paper analysis): alpha spectrometry results showing total uranium and isotopic ratios - KTA 2010.

UNCLASSIFIED

Sample descriptor	Fresh weight (g)	Dry weight (g)	Ashed weight (g)	Measured activity of dry sample (mBq/g)				²³⁸ U/ ²³⁴ U ratio
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
S1	205.4	142.0	136.2	12.95 ± 2.35	0.58 ± 0.38	12.93 ± 2.35	26.45 ± 3.35	1.00 ± 0.26
S2	228.3	186.5	181.0	15.99 ± 2.71	0.20 ± 0.22	19.26 ± 3.11	35.45 ± 4.13	0.83 ± 0.19
S3	171.6	138.6	134.5	13.24 ± 2.63	0.43 ± 0.39	11.81 ± 2.44	25.48 ± 3.61	1.12 ± 0.32
S4	159.9	122.8	117.5	10.08 ± 1.83	0.53 ± 0.33	11.43 ± 2.00	22.04 ± 2.73	0.88 ± 0.22
S5	150.1	72.0	64.7	13.05 ± 2.30	0.59 ± 0.38	15.10 ± 2.56	28.74 ± 3.47	0.86 ± 0.21
S6	140.4	51.4	44.0	14.42 ± 2.47	0.74 ± 0.43	15.44 ± 2.59	30.61 ± 3.61	0.93 ± 0.22
S7	110.8	78.1	75.1	10.42 ± 1.94	0.11 ± 0.15	10.27 ± 1.92	20.80 ± 2.73	1.01 ± 0.27
S8	87.6	51.2	48.2	10.81 ± 1.96	0.30 ± 0.25	10.55 ± 1.93	21.66 ± 2.76	1.02 ± 0.26
S9	168.5	85.4	79.2	10.20 ± 1.87	0.80 ± 0.42	11.09 ± 1.98	22.09 ± 2.76	0.92 ± 0.24
S10	146.4	98.5	94.6	12.14 ± 2.56	0.76 ± 0.53	11.95 ± 2.53	24.85 ± 3.64	1.02 ± 0.30
S11	128.8	72.6	67.7	10.58 ± 1.84	0.38 ± 0.27	11.95 ± 2.00	22.91 ± 2.73	0.89 ± 0.21
S12	151.1	95.8	90.2	11.71 ± 2.07	0.18 ± 0.19	13.34 ± 2.27	25.23 ± 3.07	0.88 ± 0.22
S13	93.8	40.2	35.7	15.45 ± 2.49	0.76 ± 0.40	15.77 ± 2.52	31.98 ± 3.57	0.98 ± 0.22
S14	84.7	23.1	18.3	16.86 ± 2.69	0.54 ± 0.35	16.37 ± 2.63	33.77 ± 3.77	1.03 ± 0.23
S15	98.2	56.4	53.2	13.13 ± 2.52	0.38 ± 0.34	15.77 ± 2.86	29.28 ± 3.83	0.83 ± 0.22

Table 11. Stream sediments: alpha spectrometry results showing total uranium and isotopic ratios – KTA 2010.

Note (for Tables 6 to 11): Activity results have been rounded to 2 decimal place. All uncertainties are stated at a 95% confidence level. Limits of Detection (LOD) are calculated by a 'modified Currie' formula⁵ at 95%. The total activity is calculated from the sum of the actual activities for each isotope, regardless of the LOD quoted for that isotope. Therefore, where activities are reported as less than LOD for any of the uranium isotopes, the total uranium value may not be equal to the sum of the individual isotopic values. Where the isotopic activities are below the LOD for more than one isotope, the total activity and the isotopic ratios could not be calculated and are reported as n/a.

Station number	Location	Average dose rate (µGy/h) (n = 3)
1	Raeberry Target	0.093
2	Raeberry Bunker	0.098
3	Raeberry Gun	0.106
4	India Target	0.091
5	Balig Gun/Target waypoint	0.095
6	Balig Gun	0.084
7	Zulu Gantry	0.089
8	Mullock Farm	0.085
9	Silver Hill (Low) Gun	0.103
10	Echo Target (Doon Hill)	0.089
11	Doonhill/Target waypoint	0.077
12	Doon Hill Gun	0.087
13	Gypsy Point	0.100

Table 12. Environmental gamma dose rates - KTA 2010.

⁵ Hurtgen C, Jerome S, Woods M. (2000) 'Revisiting Currie - how low can you go?' *Applied Radiation and Isotopes* 53 pp 45-50


UNCLASSIFIED

This page is intentionally blank.


UNCLASSIFIED

Historical Data

To allow year-to-year comparison, data from the 1996 to 2010 KTA environmental surveys are presented on the following pages. Data for soil, grass and animal indicator analyses are presented separately.



UNCLASSIFIED



UNCLASSIFIED

UNCLASSIFIED

Sample Type	1998	1999	2000	2002	2003	2004	2005	2006	2007	2008	2009	2010
Cow Faeces	1.4 (0.9)	0.8 (0.7)	31.5 (1.1)	0.2 (0.8)	1.6 (0.5)	0.5-1.5 (0.6-0.8)	0.4-7.7 (0.7-1.5)	1.1-1.5 (0.9-1.0)	1.0-2.6 (0.7-1.2)	0.9-1.5 (0.6-0.9)	0.7-14.3 (0.8-1.2)	0.2-2.2 (0.6-1.1)
Sheep Faeces	3.1 (0.6)	0.7-7.8 (0.9-1.7)	3.9 (1.3)	5.2 (0.7)	0.9 (1.5)	0.5 (1.5)	x	0.3 (1.0)	1.2-14.5 (0.8-0.9)	7.4 (1.0)	x	0.2-0.5 (0.7-1.1)
Fox Faeces	11.4 (0.7)	6.0 (0.8)	46.7 (6.7)	8.7 (0.6)	5.2 (2.2)	x	x	12.4 (1.8)	x	x	x	x
Deer Faeces	7.8 (2.7)	0.4-0.6 (1.7 ± 1.5)	x	1.0 (0.7)	0.5 (0.7)	x	1.4 (1.0)	0.5 (2.7)	x	0.5 (n/a)	4.9 (0.7)	x
Rabbit Faeces	0.3 (0.9)	x	3.9 (2.0)	42.6 (7.2)	0.5 (1.0)	0.5 (1.5)	1.2 (0.9)	1.3 - 5.8 (1.2 - 1.4)	1.2-13.2 (1.1-2.3)	0.7-1.4 (0.9-1.2)	1.2 (0.8)	x
Badger Faeces	x	x	x	x	x	x	x	x	0.1-1.2 (1.3 ± 0.5)	x	x	x
Deer Kidneys	x	x	x	x	x	x	0.1 ± 0.0 (n/a)	x	0.0-0.1 (n/a)	0.1 (n/a)	x	x
Mushroom	x	x	x	x	0.5 (n/a)	x	x	x	x	x	x	x
Black-berries	x	x	x	0.2 (n/a)	x	x	x	x	x	x	x	x
Water samples	3.8-16.6 (1.1-1.4)	9.0-14.6 (n/a)	12.3-13.0 (n/a)	2.7-7.1 (n/a)	1.7-4.7 (n/a)	<11 (n/a)	1.8-5.1 (n/a)	1.8-9.1 (n/a)	2.0-7.5 (n/a)	2.7-6.4 (n/a)	3.7-9.6 (0.4)	1.8-7.9 (0.7)

Total uranium results for all terrestrial biological indicator samples (mBq/g of dry weight) and water samples (in mBq/l) 1998-2010.

UNCLASSIFIED

Notes: * denotes no sample of this type was collected.

For simplicity, results for both the total uranium activity and the isotopic $^{238}\text{U}/^{234}\text{U}$ ratio are reported as follows:

- The activity is reported first in mBq/g, followed by the isotopic ratio (italicised and in parenthesis).
- Where the isotopic activities are below the Limit of Detection (LOD) for more than one isotope, the isotopic ratios could not be calculated and are reported as 'n/a'.
- When only one result is reported for a sample type; the result is reported together with its uncertainty. Where there is more than one result for a sample type, the range (min – max) is reported. The analytical error for these results, which is not reported here, is typically below 20%.

ANNEX A Issues to be considered when interpreting or comparing uranium data

A.1 There are a number of issues that may give rise to uncertainties when interpreting or comparing uranium data. These include:

- Analytical technique;
- Statistical variation;
- Spatial variability;
- Temporal variability; and
- Species variation (for plant and animal samples).

Analytical approaches

A.2 Soil sample results may be reported as either dry weight or wet weight depending on whether the masses of the samples were obtained prior to or after drying. This will have implications for comparison of results between the surveys at Kirkcudbright, which are reported as dry weight and other UK uranium in soil data, which may be reported as wet weight. Samples reported as dry weight will appear to have concentrations of uranium approximately 20% higher than those reported as wet weight (although this will depend on the moisture content).

A.3 For analysis techniques such as Inductively Coupled Plasma Mass Spectrometry (ICP-MS) or alpha spectrometry, the uranium present in a sample may be extracted into solution by either leaching the soil samples or totally dissolving them. Total dissolution will give rise to higher uranium results because the analysis will include all uranium including that which is contained within the mineral grains. Leached samples, in comparison, will only contain uranium that is either easily dissolved or is adhered to the surfaces of mineral grains. This limitation is acceptable as any DU contamination which may be present at Kirkcudbright is likely to be leachable. Total sample analysis techniques such as gamma spectrometry will give results similar to those for total dissolution. Given the differences between the results for total analysis and leached analysis, care should be taken when comparing sets of data to ensure that either the same approach has been used or that the differences are appropriately discussed.

A.4 Uranium concentrations in plants may be affected by contamination of foliage or roots with dust or soil particles. Preparation of plants for analysis may or may not involve a washing stage. It is therefore important to be aware of the preparation approaches that have been applied when comparing the results of different plant analyses.

Statistical variations

A.5 There is a statistical uncertainty associated with the result of a radiochemical analysis (identified in the counting statistics for the technique). However, this uncertainty is likely

to be small in comparison with the overall variation in activity between environmental samples.

Spatial variations

- A.6 DU contamination will not be uniformly distributed within a sampling area. Any DU firing malfunctions will probably result in small fragments distributed over an area. Within that area, sampling at some locations would indicate contamination, whereas at others the soil would appear to be clean. Hence, the repeat sampling and analysis of soils from within an area may give rise to a significant degree of variation.
- A.7 In addition to DU contamination due to firing at the range, there may be variations in uranium concentrations due to local anthropogenic or natural discharges. For example, natural uranium concentrations may be enhanced by the local application of phosphate based fertiliser to agricultural land. Most of the phosphate fertiliser applied to clay loam soils in England in the last 100 years can be still be found retained in the top 23 cm layer of soils, with no increase at greater depths [A1]. Veins rich in uranium minerals occur naturally along the coast of the Solway Firth, such as uraninite found at Needle's Eye, Caulkerbush, approximately 24 km away from KTA on the north coast of the estuary. These features are thought to be present across the region [A2, A3], although this has not been studied specifically.

Temporal variations

- A.8 There will be natural temporal variations in the uranium concentration and in the abundance of the various isotopes in the samples due to seasonal variations in rainfall. Rainfall can impact on dust re-suspension and deposition on grass as well as on the dissolution and migration of surface uranium down the soil profile. Some concentrations may be slightly elevated at the time of sampling, whilst others may be slightly below expected background levels.
- A.9 The activities of samples from any particular sampling site may vary from year to year. This may relate to temporal changes in uranium concentration, but will also be affected by spatial variation (see above).
- A.10 Plant uptake of radionuclides is affected by the period in the plant growing cycle. This is also mirrored in the animal uptake of radionuclides within their life cycle.

Species variations

- A.11 Plant uptake of radionuclides is affected by the soil characteristics (uranium concentration and speciation as well as other soil physico-chemical characteristics) and varies with plant species. In general, leafy vegetables take up higher concentrations than fruit and grain crops. Uranium tends to be preferentially distributed in the leaves and stems rather than in the roots, fruits or seeds [A1].
- A.12 Animal uptake of uranium is affected by their life habits, feeding patterns, physiology and the uranium concentration in their foodstuffs and the environment. Because it is clearly impractical to consider all species in a particular environment, the current approach in radiological dose assessments to non-human species is to advocate the use of reference organisms (see Reference A4). Although this environmental survey does not aim at

UNCLASSIFIED

compiling a dose assessment for non-human species, wherever possible, animal faeces samples have been collected and analysed.

Annex A References

- [A1] Mortvedt, JJ. (1992) Plant and soil relationships of uranium and thorium decay series radionuclides – A review, *Journal of Environmental Quality*, 23, 643.
- [A2] Basham, I R; Milodowski, A E; Hyslop, E K; Pearce, J M. (1989) The location of uranium in source rocks and sites of secondary deposition at the Needle's Eye natural analogue site, D&G, British Geological Survey Technical Report WE/89/56.
- [A3] Milodowski, A E. *et al.* (1990) Uranium-mineralised micro-organisms associated with uraniferous hydrocarbons in southwest Scotland. *Nature*, 347, 465.
- [A4] Environmental risk from ionising contaminants: Assessment and management (ERICA), EC 6th Framework Programme (Contract FI6R-CT-2003-508847)
[web reference: <http://www.ERICA-project.org>].

UNCLASSIFIED

ANNEX B Change in the $^{238}\text{U}/^{234}\text{U}$ activity ratio of a medium containing natural uranium with the addition of depleted uranium

Mass proportion of DU added*	Activity Concentration (mBq/kg)**			Ratio of total activity natural uranium to total activity	$^{238}\text{U}/^{234}\text{U}$ activity ratio
	U-238	U-235	U-234		
0	3.7×10^4	1.7×10^3	3.8×10^4	1.0	1.0
1	7.4×10^4	2.2×10^3	4.3×10^4	1.6	1.7
2	1.1×10^5	2.7×10^3	4.8×10^4	2.1	2.3
3	1.5×10^5	3.2×10^3	5.3×10^4	2.6	2.8
4	1.9×10^5	3.7×10^3	5.8×10^4	3.2	3.2
5	2.2×10^5	4.1×10^3	6.3×10^4	3.8	3.6
6	2.6×10^5	4.6×10^3	6.8×10^4	4.3	3.9
7	3.0×10^5	5.1×10^3	7.2×10^4	4.9	4.1
8	3.4×10^5	5.6×10^3	7.7×10^4	5.4	4.3
9	3.7×10^5	6.0×10^3	8.2×10^4	6.0	4.5
10	4.1×10^5	6.5×10^3	8.7×10^4	6.5	4.7
20	7.8×10^5	1.1×10^4	1.4×10^5	12.0	5.8
60	2.3×10^6	3.1×10^4	3.3×10^5	34.1	6.9
80	3.0×10^6	4.0×10^4	4.3×10^5	45.1	7.1
100	3.8×10^6	5.0×10^4	5.2×10^5	56.1	7.2
200	7.5×10^6	9.8×10^4	1.0×10^6	111.0	7.4
600	2.2×10^7	2.9×10^5	3.0×10^6	332.0	7.6
800	3.0×10^7	3.9×10^5	3.9×10^6	442.0	7.6
1000	3.73×10^7	4.8×10^5	4.9×10^6	552.0	7.6

Table reproduced from Volume 2 - Appendices, WS Atkins Environmental Assessment on DU Firings.

* The value represents the additional mass of depleted uranium added (all radionuclides) relative to the original mass of natural uranium present (3 mg U/kg soil).

** Table assumes 3 mg U/kg of natural uranium present in soil in following proportion: ^{238}U (2.978 mg /kg); ^{235}U (0.022 mg /kg); ^{234}U (2e-04 mg /kg), prior to addition of DU.

Initial Distribution

CESO(Army) SO2 RAD (2 hard copies plus electronic copy)

House of Commons Library (Electronic copy)

Head of Radioactive Waste - Environment Group, Scottish Government (8 copies, of which 6 copies are to be placed in the Scottish Government Library.)

Aaron Scarlett

Dee Emerson

Knowledge Services (Electronic copy)

Report Documentation Form

A copy of this form is to be completed by the principal author for all Dstl reports. When complete, it is to be bound as the last numbered pages of the finished report.

1. Originators Report Number incl. Version No		DSTL/TR51735 V1	
2. Report Protective Markings and any other markings e.g. Caveats, Descriptors, Privacy markings UNCLASSIFIED			
3. Title of Report Depleted Uranium Survey Report: Kirkcudbright Training Area 2010 Part 1 Terrestrial Environment			
4. Title Protective Markings incl. any Caveats		UNCLASSIFIED	
5. Author Aaron Scarlett			
6. Originator's Name and Address Aaron Scarlett DSTL Alverstoke c/o Institute of Naval Medicine Crescent Road Gosport PO12 2DL		7. MOD Sponsor Name and Address CESO(A) SO2 RAD Headquarters Land Forces IDL420, Ramillies Building Marlborough Lines, Monkston Road ANDOVER SP11 8HJ	
8. MOD Contract number and period covered		701418	
9. Other Report Nos.			
10. Date of Issue 31 March 2011	11. Pagination 44	12. No. of References 25	
13. Abstract (A brief (approximately 150 words) factual summary of the report) Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the radiological impact of the firings on the terrestrial and marine environments and any associated risk to humans. This report presents the findings of the terrestrial survey undertaken at KTA during 2010; the marine survey is reported separately in Part 2. No sample analysed was radioactive within the meaning of the Radioactive Substances Act 1993, nor did it exceed the Generalised Derived Limits advised by the Health Protection Agency. The survey results indicate that the existing areas of low-level DU contamination on site, which are maintained within fenced compounds, are not leading to the transfer of uranium to other parts of the environment. Based on the findings of this survey – which are consistent with those of previous surveys – potential doses to representative persons, site personnel and members of the public are deemed to be consistent with natural background exposure.			
14. Abstract Protective Marking including any Caveats UNCLASSIFIED			
15. Keywords/Descriptors (Authors may provide terms or short phrases which identify concisely the technical concepts, platforms, systems etc. covered in the report. Kirkcudbright, Depleted Uranium, Radioactivity, Environmental, Survey, Terrestrial.			

UNCLASSIFIED

16. Report Announcement (refers to title/abstract being included in accessions lists e.g. Defence Reports Abstracts)

Announcement of this report is UNLIMITED

If there are limitations on the announcement of this report please indicate below the groups to whom it can be announced (more than one if required)

- Can be announced to MOD and its Agencies
- Can be announced to UK Defence Contractors
- Can be announced to Overseas Defence Departments
- Other (please specify)

17. Report Availability

- UNLIMITED distribution
- No Release without approval of the Release Authority (see box 7)




If the above do not apply, please indicate below the groups to whom the report may be released upon request without further Need-To-Know checks.

- Can be released to MOD and its Agencies
- Can be released to other UK Government Departments
- Can be released to UK Defence Contractors
- Can be released to Overseas Defence Departments
- Other (please specify)

18. Downgrading Instructions (check as appropriate)

- This report may be automatically downgraded to _____ after _____ years
- This report may be reviewed _____ years after publication

19. Authorisation

	Name	Signature	Date
Chief Scientist	Dr Adrian Baker		6/9/11
Project Manager	Dee Emerson		6/4/11
Technical Reviewer	Ron Brown		6/11/11

UNCLASSIFIED

This page is intentionally blank

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED