

UNCLASSIFIED



**Depleted Uranium Survey Report:
Kirkcudbright Training Area 2008
Part 2 Marine Environment**

Aaron Scarlett

DSTL/TR37610 V1
04 February 2010

Environmental
Sciences Department
C/o INM
Crescent Road
Alverstoke
Hampshire
PO12 2DL

© Crown copyright 2010 Dstl

UNCLASSIFIED



Dstl is part of the
Ministry of Defence

UNCLASSIFIED




This document has been prepared for MOD and, unless indicated, may be used and circulated in accordance with the conditions of the Order under which it was supplied.

It may not be used or copied for any non-Governmental or commercial purpose without the written agreement of Dstl.

© Crown copyright 2010
Defence Science and Technology Laboratory UK

Approval for wider use or release must be sought from:

Intellectual Property Department
Defence Science and Technology Laboratory
Porton Down, Salisbury, Wiltshire, SP4 0JQ.

Authorisation			
	Name	Signature	Date
Lead Technical Reviewer	Dr David Smith		5/2/2010
Project Manager	Dee Emerson		5/2/10
Technical Reviewer	Ron Brown		5/2/10

Executive Summary

Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the extent of any environmental impact of the firings on the terrestrial and marine environments and any associated radiological risk.

This report presents the findings of the marine survey undertaken in the areas surrounding KTA during 2008; the terrestrial survey is reported separately in Part 1. The survey was undertaken to monitor the levels of any uranium in the marine environment resulting from operations on the site and to identify the extent of any environmental transfer processes.

None of the samples analysed were radioactive within the meaning of the Radioactive Substances Act 1993 (RSA93) nor did they exceed a very small fraction of the Generalised Derived Limits (GDLs) advised by the Health Protection Agency (formerly the National Radiological Protection Board).

The results of the 2008 survey agree with those from previous years and do not indicate any health or environmental impact from the firing of DU. There is no evidence to indicate that members of the public are exposed to a radiological hazard from the marine environment as a result of test firing DU ammunition at Kirkcudbright.

UNCLASSIFIED

This page is intentionally blank

UNCLASSIFIED

Table of Contents

Executive Summary	3
List of Figures	6
List of Tables	6
1 Introduction	7
2 Background	7
3 Depleted Uranium (DU)	9
4 Differentiating DU from Natural Uranium	10
5 Reference Levels	11
6 Methodology	12
7 Results and Interpretation	17
8 Evaluation of Potential Exposure Pathways	20
9 Conclusions	21
10 List of References	22
11 KTA Marine Survey Results	23
Historical Data	27
ANNEX A Issues to be considered when interpreting or comparing uranium data	29
ANNEX B Change in the $^{238}\text{U}/^{234}\text{U}$ activity ratio of a medium containing natural uranium with the addition of depleted uranium	32
ANNEX C Variability of uranium concentration and uranium isotopic ratios in marine environmental samples	33
ANNEX D Reference values for uranium in the Solway Firth and the UK	35
Initial distribution	37

List of Figures

Figure 1. Number of DU projectiles fired from KTA, from 1982 to 2008.	8
Figure 2. Approximate cumulative mass of DU projectiles fired from KTA, from 1982 to 2008.	8
Figure 3. Kirkcudbright inter-tidal sampling locations 2008.	12
Figure 4. Approximate underwater sampling locations 2008 (showing line of fire for each battery).	15

List of Tables

Table 1. Mass compositions of uranium isotopes in natural and depleted uranium.	9
Table 2. Reference levels relevant to the Kirkcudbright marine survey.	11
Table 3. Summary of inter-tidal sample collection and measurements 2008.	14
Table 4. Summary of sample analyses - KTA 2008.	17
Table 5. Inter-tidal environmental gamma dose rates - KTA 2008.	24
Table 6. Intertidal sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2008.	24
Table 7. Seaweed samples: alpha spectrometry analysis results showing total uranium and isotopic ratios – KTA 2008.	25
Table 8. Biota samples: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2008.	25
Table 9. Underwater sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios – KTA 2008.	26
Table 10. Historical summary of total uranium and isotopic ratios for sediment and biota samples 1996 to 2008.	28

1 Introduction

- 1.1 Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess the extent of any environmental impact of the firings on the terrestrial and marine environments and any associated radiological risk [1 to 11].
- 1.2 This report presents the findings of the marine survey undertaken in the areas surrounding KTA during 2008; the terrestrial survey is reported separately in Part 1 [12]. The survey was undertaken to monitor the levels of any uranium in the marine environment resulting from operations on the range and to identify the extent of any environmental transfer processes.
- 1.3 Since its inception in 1980, the monitoring programme has evolved to incorporate changes in best practice and increased knowledge of the local environment. The changes to sampling protocols over the years were fully explained in the marine report of 2005 [3]. Since 1996, the survey methodology has remained broadly consistent and involves the annual sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood is also sampled.

2 Background

- 2.1 The KTA range is located on the coast of Dumfries and Galloway, near Castle Douglas. In April 2006, the range became part of the Defence Training Estate.
- 2.2 DU has been released into the environment at Kirkcudbright as a consequence of the test firing of DU ammunition during design and accuracy assessment trials. DU projectiles are fired through soft vertical targets and continue their trajectory coming to rest in the Solway Firth.
- 2.3 Testing of projectiles historically has taken place at five locations on the KTA range. Each battery location had a designated target and line of fire and hence a predictable area of impact ranging from several hundred metres to approximately 7 kilometres offshore. Although a small number of the DU rounds malfunctioned and impacted on land, the vast majority entered the Solway Firth.
- 2.4 The number of DU rounds fired each year at KTA from the five firing locations and the cumulative mass fired to date, are presented in Figures 1 and 2 respectively.




Figure 1. Number of DU projectiles fired from KTA, from 1982 to 2008.




Figure 2. Approximate cumulative mass of DU projectiles fired from KTA, from 1982 to 2008.

3 Depleted Uranium (DU)

- 3.1 Uranium is a naturally occurring radioactive material which exists as three isotopes: uranium-238 (^{238}U), uranium-235 (^{235}U) and uranium-234 (^{234}U). The mass composition of these isotopes is shown in Table 1. In the environment, natural uranium exists in approximate equilibrium with the daughter products of the ^{238}U and ^{235}U decay series¹ in terms of radioactivity. Together these isotopes emit a range of alpha and beta particles along with gamma radiation. The chemical toxicity of uranium is approximately equal to that of lead.
- 3.2 Uranium in an 'enriched' form is used as fuel in nuclear reactors. The enrichment process increases the concentration of ^{235}U (above 0.72%) in comparison to the natural form. The by-product of this process is 'depleted' uranium (DU), which has a reduced concentration of ^{235}U . Uranium-234 is also removed in the depletion process; DU is consequently less radioactive than natural uranium (the specific alpha activity of the DU fired at Kirkcudbright being approximately 1.4×10^7 milli becquerels per gram (mBq/g) compared to 2.5×10^7 mBq/g for natural uranium). The mass compositions of DU and natural uranium are presented in Table 1 below.

Form of Uranium	^{238}U	^{235}U	^{234}U
Natural uranium	99.274%	0.72%	0.00554%
The DU used at Kirkcudbright	99.8%	0.20%	0.0008%

Table 1. Mass compositions of uranium isotopes in natural and depleted uranium.

- 3.3 As discussed in paragraph 3.1, ^{234}U normally exists in approximate equilibrium with ^{238}U in the natural environment. In comparison, DU exhibits a $^{238}\text{U}/^{234}\text{U}$ activity ratio of between 7:1 and 8:1, dependant on the degree of depletion achievable by different methods of processing. This distinction is important in differentiating DU contamination from naturally occurring uranium in the environment (see section 4). For the remainder of this report, isotopic ratios will be stated in terms of activity rather than mass and as a single value representing the ratio of becquerels of ^{238}U to 1 becquerel of ^{234}U (i.e. a $^{238}\text{U}/^{234}\text{U}$ ratio of 7 rather than 7:1).

¹ A radioactive decay series occurs when a heavy radionuclide decays into successively lighter radionuclides. For example, ^{238}U decays to ^{234}Th , then $^{234\text{m}}\text{Pa}$, then ^{234}U and so on until a stable element is reached (^{206}Pb).

4 Differentiating DU from Natural Uranium

- 4.1 The fundamental requirement of the DU environmental monitoring programme is to quantify the impact of DU firing. This is achieved partly by measuring the amount of total uranium in environmental materials and using this figure as an upper bound of DU contamination levels. However, as uranium is present at detectable levels in most environmental media, this overestimates the risk. More sophisticated analyses involve the specific measurement of ^{238}U and ^{234}U isotopes (by activity and/or mass). Although isotope measurements are used in this survey, references to total uranium measurements are included for consistency with historic reports. The limitations of using total uranium concentrations are discussed further in Annex A.
- 4.2 A convenient fingerprint marker for DU contamination is the $^{238}\text{U}/^{234}\text{U}$ activity ratio. The DU fired at KTA has a $^{238}\text{U}/^{234}\text{U}$ activity ratio of approximately 7, whereas natural uranium in the environment has an activity ratio close to unity. Environmental samples are therefore analysed for isotopes of ^{238}U and ^{234}U to determine activity ratios and hence identify the origin of the uranium.
- 4.3 Substantial deposition of DU in the environment (in addition to an existing natural uranium background) is required before the $^{238}\text{U}/^{234}\text{U}$ activity ratio diverges significantly from its natural ratio. An illustration of the impact of DU contamination on the isotopic ratio is given in Annex B. For the ratio to approach 7 in an analytical sample, the mass of DU would have to be approximately one hundred times the mass of the uranium that is naturally present. Hence, the lower the natural uranium background, the lower the levels of DU contamination that may be detected by isotopic analysis.
- 4.4 Isotopic quantification is achieved by techniques such as alpha spectrometry and mass spectrometry. Alpha spectrometry can detect uranium to parts per billion, which is equivalent to mBq per kg, or to lower levels if count times are increased. Mass spectrometry is more sensitive, but the lower levels detectable are of no recognised health significance. Isotopic information can also be yielded from gamma spectrometry analyses, although limits of detection are not generally sufficient for measurement of environmental levels.

5 Reference Levels

- 5.1 The Depleted Uranium Firing Environmental Review Committee (DUFERC), on which the MOD is represented, has agreed investigation/action levels for levels of DU in the terrestrial environment. Reference values for the marine environment are taken directly from legislation and guidance; as discussed below.
- 5.2 For uranium in sediment, the activity concentration can be compared to the Schedule 1 limit defined in the Radioactive Substances Act 1993 and Generalised Derived Limits (GDLs) advised by the Health Protection Agency (formerly the National Radiological Protection Board) [13 to 15]. GDLs for uranium were last updated in 2000 and were referred to by the Royal Society in their studies of the potential health effects of using DU munitions [16]. They are based on a 1 milli-sievert (mSv) potential exposure to a member of the public: the dose limit set by the International Commission on Radiological Protection (ICRP) and incorporated into UK statute [17].
- 5.3 Uranium levels in foodstuffs such as fish, crustaceans and molluscs can be compared against the relevant GDL (expressed as fresh mass of the edible fraction). For biota such as seaweed, the European Council Food Intervention Level (CFIL) for ‘other foodstuff’ is referred to [18].
- 5.4 Reference levels relevant to the Kirkcudbright marine survey are provided in Table 2.

Source	Reference Level	Activity concentration (mBq/g)
RSA 93	Schedule 1 limit - level at which regulatory control is required.	11,100
NRPB (2000)	Generalised Derived Limit: Marine sediment (dry weight)	100,000
	²³⁸ U in marine fish (fresh weight)	200
	²³⁸ U in crustaceans (fresh weight)	1,000
	²³⁸ U in molluscs (fresh weight)	1,000
CEC Regulation (Euratom) 2218/89	European Council Food Intervention Level (CFIL) for ‘other foodstuff’	1,250

Table 2. Reference levels relevant to the Kirkcudbright marine survey.

- 5.5 In addition to these reference values, it is also possible to compare the results with those of similar surveys carried in the UK. The Radioactivity in Food and the Environment (RIFE)[19] reports present the findings of an annual independent survey carried out by the Centre for Environment, Fisheries and Aquaculture Science (CEFAS) on behalf of the environment agencies and the Food Standards Agency. Where possible, monitoring results are compared against those of the RIFE reports and other relevant surveys such as

Reference 20. The latter includes monitoring data for Sandyhills Bay (also monitored by Dstl); which is sufficiently close to KTA to be similar in geology, but sufficiently far away to be considered unaffected by activities at the range.

6 Methodology

6.1 The survey methodology includes the sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood is also sampled. The methodology is outlined in the following paragraphs.

Inter-tidal zone sampling

6.2 Sampling and dose rate measurements were carried out at 11 pre-determined locations as shown at Figure 3. These locations have been chosen to provide a suitable distribution of sampling points spanning the areas likely to be affected by the DU firing at Kirkcudbright. It should be noted that sampling location No. 5 had to re-positioned due to problems with gaining access to the area (this is likely to be the case for subsequent surveys).



Figure 3. Kirkcudbright inter-tidal sampling locations 2008.

UNCLASSIFIED

6.3 The Ordnance Survey of Great Britain grid references for the sampling locations in 2008 are given below:

1.	South Carse	NX 99456 59431
2.	Sandyhills Bay	NX 89213 55172
3.	Port O'Warren Bay	NX 87901 53438
4.	Port Donnel	NX 84776 53714
5.	Balcarry Bay	NX 82182 49544
6.	Abbey Burn Foot	NX 74200 44413
7.	Mullock Bay	NX 70996 43765
8.	Lower Nunton Bay	NX 66081 48424
9.	Brighthouse Bay	NX 63508 45401
10.	Carrick Point	NX 57599 50722
11.	Mossyard Bay	NX 55207 52007

6.4 Inter-tidal sediment was collected from each sampling location for analysis by alpha spectrometry. The sediment was collected as close to the low water mark as possible, giving due consideration for the safety of the survey team. Due to the abundance of available inter-tidal sediment, composite samples were collected by taking the top few millimetres of sediment from a number of locations until enough material was collected to fill a 0.8 litre container. Shells, stones and other unwanted debris were removed from samples and any excess water was drained off.

6.5 Seaweed samples were collected from sampling locations where available for analysis by alpha spectrometry. The most recent growth of seaweed was collected by cutting the top 5cm of fronds from a number of locations and combining the material to form a composite sample (approximately 1.8 litre sample volume). The collected samples, which consisted of a single species (*Fucus vesiculosus*), were washed in sea-water to remove any sediment.

6.6 Mollusc samples were collected from each sampling location where found in abundance. A composite sample was collected in a 1.8 litre sample container and washed in sea-water to remove any sediment. In 2008, the only mollusc species available for sampling was the common mussel (*Mytilus edulis*). The samples were washed a second time and boiled on the day of collection to avoid them spoiling before analysis. The samples were placed in a re-sealable bag and frozen prior to transport to the laboratory for analysis by alpha spectrometry.

6.7 Environmental gamma dose rates were recorded at each sampling location using the Mini Instruments 6-80/81 fitted with an MC71 compensated Geiger-Muller tube. The instrument was allowed to record over a period of 300 seconds at three points along the low water mark and at one point at the high water mark (at one metre above the ground). The environmental gamma dose rate was derived by taking an average of the four measurements.

6.8 A summary of samples collected and dose rate measurements taken from the inter-tidal zone is provided in Table 3 overleaf.

UNCLASSIFIED

Sample area number	Sample type and number of samples			Environmental dose rates	
	Sediment	Seaweed	Mussel	LWM @ 1m height	HWM @ 1m height
1	✓	*	*	✓	✓
2	✓	✓	*	✓	✓
3	✓	✓	✓	✓	✓
4	✓	✓	✓	✓	✓
5	✓	✓	✓	✓	✓
6	✓	✓	*	✓	✓
7	✓	✓	*	✓	✓
8	✓	✓	✓	✓	✓
9	✓	✓	*	✓	✓
10	✓	✓	✓	✓	✓
11	✓	✓	*	✓	✓

Table 3. Summary of inter-tidal sample collection and measurements 2008.

Notes: LWM: low water mark, HWM: high water mark, *: sample not collected

Underwater sediment sampling

6.9 Underwater sediment samples were collected from 25 locations by use of a scooping device with a rubberised bag which was lowered into the water at each sampling point. The sample locations are shown on Figure 4 and the co-ordinates (latitude and longitude) are provided below:

- | | |
|-----------------------------|-----------------------------|
| 1. N 54 45 078 W 03 59 782 | 14. N 54 45 375 W 03 57 744 |
| 2. N 54 45 600 W 03 59 808 | 15. N 54 45 673 W 03 57 125 |
| 3. N 54 44 527 W 03 59 746 | 16. N 54 44 136 W 03 57 108 |
| 4. N 54 45 569 W 04 02 577 | 17. N 54 44 575 W 03 57 081 |
| 5. N 54 45 032 W 04 02 541 | 18. N 54 42 874 W 04 01 311 |
| 6. N 54 44 479 W 04 02 522 | 19. N 54 42 750 W 04 01 290 |
| 7. N 54 45 211 W 04 01 174 | 20. N 54 43 235 W 04 02 493 |
| 8. N 54 45 083 W 04 01 243 | 21. N 54 42 732 W 04 02 476 |
| 9. N 54 44 827 W 04 01 391 | 22. N 54 42 225 W 04 02 451 |
| 10. N 54 44 684 W 04 01 477 | 23. N 54 43 292 W 04 00 243 |
| 11. N 54 44 991 W 03 58 807 | 24. N 54 42 785 W 04 00 226 |
| 12. N 54 44 856 W 03 58 758 | 25. N 54 42 258 W 04 00 192 |
| 13. N 54 45 484 W 03 57 969 | |

6.10 The exact sampling locations may vary due to the effect of underwater currents on the sampling rig which was dragged along the sea bed for distances ranging from 50 to 100

metres. The co-ordinates given in paragraph 6.9 should therefore be considered as the approximate centre point of sampling areas of approximately 100 metres in radius.

6.11 Sediment collected from the sea-bed was screened for the presence of DU fragments by monitoring using a Mini Monitor and 44B probe. A portion of the sediment (approximately 0.8 litres) was sampled for analysis by alpha spectrometry.




Figure 4. Approximate underwater sampling locations 2008 (showing line of fire for each battery).

Locally sourced seafood

- 6.12 Locally sourced seafood is sampled as part of the Kirkcudbright survey to determine the extent of any potential transfer of uranium through the food chain. In 2008, a total of 3 lobsters (*Homarus gammarus*), 5 crabs (*Cancer pagurus*) and one bag of common whelks (*Buccinum undatum*) were purchased for analysis by alpha spectrometry. The samples were boiled on the day of collection before being frozen and transported to the laboratory.

Sample descriptors

- 6.13 Each sample was given a unique sample descriptor. For inter-tidal and underwater sediment, seaweed and mussel samples, the descriptor is followed by a number which relates to the specific sample point. For samples which do not come from a defined sample point (e.g. locally sourced lobster); no sample point is given. In this case, samples are given a letter suffix to distinguish between them. Some examples of typical sample descriptors are provided below:

I/T 4: Inter-tidal sediment sample from sample point 4

U/W 8: Underwater sediment sample from sample point 8

M. edulis 9: Mussel sample from sample point 9

C. pagurus A: Locally sources crab sample (sample A)

Sample preparation and laboratory analysis

- 6.14 Sample preparation took place either at the time of collection or at the Dstl UKAS accredited radiochemistry laboratory. Samples were analysed by alpha spectrometry with uranium concentration reported in mBq/g (equivalent to Bq/kg) of dry weight. An outline of the analysis procedure is provided below.
- 6.15 Mussel, whelk, crab and lobster samples were boiled prior to analysis and any shells were removed. The brown and white crab meat was used for analysis, but the digestive tracts were discarded. Mussels and whelks were analysed as composite samples (the whelks being divided into six equally sized portions). The crab and lobster samples were analysed individually.
- 6.16 All samples were weighed before drying to remove moisture and then weighed again (with results being reported as dry weight). The samples were ashed to remove organic material and homogenised by hand. During this process the quantity of uranium in each of the samples remains constant. The ashed samples were boiled in concentrated mineral acid (nitric acid and hydrochloric acid) to remove the 'loose' and leachable uranium from the sample. Recalcitrant matrices such as mineral grains are not broken down by the process and hence the natural uranium within them was not extracted. The samples were filtered to remove solids.
- 6.17 Uranium separation was carried out by extraction chromatography. Each eluted sample was electro-deposited onto a stainless steel planchette and the activity of each planchette was counted in a low background, silicon surface barrier, alpha spectrometer.

7 Results and Interpretation

7.1 A summary of the results for all marine samples collected in 2008 is given below in Table 4. The full marine monitoring results are provided in Tables 5 to 9 in Section 11. Historical monitoring results for the KTA for the years 1996 to 2008 are presented in Appendix A.

Sample Type	Number of Samples	No. of samples possibly containing detectable DU ²	Total uranium concentration (mBq/g)			
			Mean	Standard deviation of the mean	Minimum	Maximum
Inter-tidal Sediment	11	0	22.3	8.6	11.3	35.8
Seaweed	10	0	8.9	2.6	5.3	12.3
Mussels	5	0	7.7	1.5	5.7	9.6
Lobsters	3	0	0.2	0.1	0.1	0.3
Whelks	6	0	1.0	0.2	0.8	1.4
Crabs	5	0	0.3	0.1	0.2	0.5
Underwater Sediment	25	7	26.3	2.5	23.0	31.9

Table 4. Summary of sample analyses - KTA 2008.

Inter-tidal zone and biota sampling

7.2 Environmental gamma dose rate measurements for the inter-tidal monitoring locations are provided in Table 5. Measurements ranged from 115 to 163 nGy/h. Considering the natural variation in background dose rates, these results are generally consistent with those recorded in previous surveys of the inter-tidal zone [1-11]. Measurements recorded over salt marsh near Kirkcudbright and reported in the most recent RIFE report (average 75 nGy/h) [19] are slightly lower, although higher background dose rates are routinely observed in the inter-tidal areas due to the different underlying geology.

7.3 Alpha spectrometry analysis results for inter-tidal sediment samples are provided in Table 6. No sample was radioactive within the meanings of RSA93; nor exceeded 0.04% of the GDL for marine sediment [15]. In terms of total uranium content, the results lie within the typical range reported in the literature³ (considering measurement uncertainty)

² An underwater sediment sample may contain DU if it has a ²³⁸U/²³⁴U isotopic ratio above a value of 0.8 (after subtraction of the measurement uncertainty). However, ratios of around 1.0 were recorded even before DU firings began; further information is provided in Annex C and D. For inter-tidal sediment and biota, samples are assumed to contain a degree of depletion if the isotopic ratio exceeds 1.0 (following subtraction of measurement uncertainty).

UNCLASSIFIED

and are consistent with the results of previous surveys [1-11]. The results also compare favourably with the level of ^{238}U found at Sandyhills Bay during an environmental survey of UK coastal regions (14 ± 0.4 mBq/g dry weight) [20] and with the sediment sample results reported by CEFAS in 2008 [19] (see Annex D).

- 7.4 Within the range of measurement uncertainty, the isotopic ratios for all the inter-tidal sediment samples were below 1.0; indicating that the uranium present is of natural origin.
- 7.5 Analysis results for the 10 inter-tidal seaweed samples (*F. vesiculosus*) are provided in Table 7. The levels of total uranium detected were low (5.3 ± 2.2 to 12.3 ± 3.9 mBq/g) and represent less than 1% of the CFIL for 'other foodstuffs'. The measured levels of ^{238}U were similar to that found previously at Sandyhills Bay (3.8 ± 0.1 mBq/g)[20] and lie within the typical UK range reported in the literature (3.8 to 18.6 mBq/g) (see Annex D). Isotopic ratios are indicative of naturally occurring uranium.
- 7.6 Alpha spectrometry analysis results for biota samples are presented in Table 8. It should be noted that GDLs and biota results reported by CEFAS [19] are expressed as activity per fresh mass. In contrast, Dstl results are expressed as activity per dry weight; resulting in higher activity concentrations. In order to directly compare the results, it is necessary to convert the reported result to fresh mass activity concentrations using the recorded wet:dry ratio (the preparatory drying stage leads to an average reduction in mass of approximately 70%). The biota results are discussed further in the following paragraphs.
- 7.7 A total of five mussel (*M.edulis*) samples were analysed in 2008. The total uranium content ranged from 5.7 ± 0.6 to 9.6 ± 0.9 mBq/g (dry weight) and the level of ^{238}U ranged from 2.7 ± 0.4 to 4.3 ± 0.6 mBq/g (dry weight). Converting these activity concentrations to wet weight using the wet:dry ratio shows that no sample exceeded 0.1% of the GDL for molluscs (1000 mBq/g fresh mass). The mussel sample results are slightly above that reported previously from Sandyhills Bay (1.1 ± 0.1 mBq/g dry weight)[20] but are consistent with the natural background levels of ^{238}U in mussels reported in the most recent RIFE report, considering conversion to activity per fresh mass (0.89 mBq/g fresh weight)[19]. Isotopic ratios are indicative of natural uranium.
- 7.8 Similarly, the levels of ^{238}U in the 6 whelk (*B.undatum*) samples represent less than 0.01% of the GDL for molluscs (converting to fresh mass). They are also consistent with the natural background levels reported in RIFE [19]. The total uranium and ^{238}U values ranged from 0.8 ± 0.2 to 1.4 ± 0.5 mBq/g and 0.4 ± 0.1 to 0.7 ± 0.3 mBq/g respectively. Isotopic ratios are indicative of natural uranium.
- 7.9 Of the five crab (*C. pagurus*) samples analysed, only two had detectable levels of ^{238}U . These relate to less than 0.01% of the GDL for crustaceans, considering a conversion to fresh weight, and are broadly consistent with the background levels for crabs reported in RIFE [19]. Where a calculation was possible, isotopic ratios indicate that the uranium was of natural origin.

³ Minerals containing uranium are widely distributed throughout the Earth's crust and the concentrations of natural uranium found locally can vary significantly due to changes in the underlying geology. Consequently, there is no single, definitive, reference level for uranium in soils and sediments; although there is broad agreement in the typical range of values published in the literature. Typical values in the UK range from 3.6 to 32.3 mBq/g (dry weight) [see Annex D], but values exceeding 100 times this typical range can be found locally.

- 7.10 No ^{238}U was detected above the lower limit of detection in any of the three lobster (*H. gammarus*) samples analysed.

Underwater sediment sampling

- 7.11 A total of 25 underwater sediment samples were collected in 2008. Screening of these samples with a Mini Monitor and 44B probe did not indicate the presence of any fragments of DU. Alpha spectrometry analysis results are shown in Table 9. No samples were radioactive within the meaning of RSA93 or exceeded 0.02% of the GDL for marine sediment. The measured levels of ^{238}U (11.1 ± 1.8 to 18.1 ± 4.5 mBq/g) lie within the typical UK coastal sediment range of 3.6 to 32.3 mBq/g (see Annex D). The total uranium levels are generally consistent with those found in previous surveys [1-11].
- 7.12 The $^{238}\text{U}/^{234}\text{U}$ ratio for 18 of the 25 samples were, within the range of measurement uncertainty, in agreement with values generally reported in marine sediment (see Annex C). Seven of the samples had isotopic ratios which were greater than the typical value of 0.8 (ranging from 0.9 to 1.0 after subtraction of the uncertainty). However, this is not unexpected as isotopic ratios of around 1.0 have been recorded in sediments from around Kirkcudbright even before DU munitions testing began. In any case, the radiological implications of these results are insignificant as the levels of ^{238}U in these samples do not exceed 0.02% of the relevant GDL.

8 Evaluation of Potential Exposure Pathways

8.1 Any contamination of the marine environment with DU could result in three potential exposure pathways for humans, as described below:

- External radiation exposure from contaminated sediment or seaweed;
- Inhalation of re-suspended DU contamination and
- Ingestion of seaweed or food products contaminated with DU

External radiation exposure

8.2 The measured radiation levels on the inter-tidal zone are consistent with natural background levels. Furthermore, background levels of uranium isotopes have been found by alpha spectrometry of inter-tidal sediment samples. The measurements carried out are sensitive enough to detect radiation at levels far below anything which could be considered as a health risk; it is therefore concluded that there is no risk from external radiation resulting from the firing of DU munitions at KTA.

Inhalation of re-suspended DU

8.3 DU which has been deposited on sediment may become re-suspended in the air especially if it is attached to items which are subject to disturbance (e.g. fisherman's netting). Once the DU has been re-suspended in the air, it is then possible to be inhaled by persons in close proximity. However, the levels of uranium identified in this report are consistent with those expected as a result of natural background radioactivity; the amount of ^{238}U representing a very small fraction of relevant GDLs. It is concluded that the risk from inhalation of re-suspended DU is insignificant

Ingestion of DU contaminated foodstuffs

8.4 The levels of uranium isotopes found in biota samples were consistent with those expected due to natural background radioactivity and represented a very small fraction of the relevant GDLs. This indicates that, in terms of potential DU contamination, there is no risk associated the consumption of locally sourced foods in the Kirkcudbright area.

8.5 Although the inadvertent consumption of seawater by members of the public is possible, it was recommended by SEPA in 2001 to discontinue the sampling of seawater. The basis of this recommendation was that the immense dilution of the Solway would never give rise to detectable levels of DU or to any significant radiation exposures.

9 Conclusions

- 9.1 The 2008 annual Kirkcudbright marine monitoring programme was undertaken to assess the levels of any DU in the environment resulting from firing of DU munitions. The monitoring programme involved the sampling of inter-tidal sediment and biota, together with the measurement of environmental gamma dose rates along the Dumfries coastline. Underwater sediment and locally caught seafood were also sampled.
- 9.2 The findings of the survey indicate that the levels of uranium present in the marine environment are largely indistinguishable from natural background radioactivity. None of the samples were radioactive within the meaning of the Radioactive Substances Act 1993; nor did they exceed a very small fraction of the Generalised Derived Limits advised by the Health Protection Agency.
- 9.3 The results of the 2008 survey agree with those from previous years and do not indicate any health or environmental impact from the firing of DU. There is no evidence to indicate that members of the public are exposed to any radiological hazards from the marine environment as a result of test firing DU ammunition at Kirkcudbright.

10 List of References

1. Toqué, C. (2008) Terrestrial Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area, 2007. Dstl/CR30510.
2. Toqué, C. (2007) Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2006, Dstl/CR26985 V1.
3. Toqué, C. (2007) Marine Environmental Depleted Uranium Survey Report, Kirkcudbright Training Area, 2005. Dstl/CR19579.
4. Toqué, C. (2006) Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area – 2004, Dstl/CR18154 V1.
5. 2003 Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area, Dstl/CR 13981 (2005).
6. 2002 Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area, Dstl/CR113980 (2005).
7. 2001 Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area, Dstl/CR12350.
8. 2000 Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area, Dstl/CR12021.
9. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area. DRPS DERA/CBD/DRPS/17/00.
10. Marine Environmental Depleted Uranium Survey Report Kirkcudbright Training Area, DRPS 9/99.
11. 1996 Marine Environmental Survey Report DTEO Kirkcudbright, DRPS 10/97.
12. Scarlett, A. (2009) Depleted Uranium Survey Report: Kirkcudbright Training Area 2008. Part 1 Terrestrial Environment. DSTL/TR37166 V1.
13. Carter, T. (2002) Comparison of Kirkcudbright and Eskmeals Environmental Monitoring Data with Generalised Derived Limits for Uranium. Dstl 167/2002.
14. Radioactive Substances Act 1993. HMSO (ISBN 0-10-54 1293-7).
15. Generalised Derived Limits for Radioisotopes of Polonium, Lead, Radium and Uranium. Documents of the NRPB, Vol. 11, No 2 (2000).
16. The Health Hazards of Depleted Uranium Munitions (Part II), (ISBN 0854035745), The Royal Society (2002).
17. Ionising Radiations regulations 1999 Statutory Instrument 1999 No. 3232.
18. CEC Council Regulation (Euratom) No 2218/89 amending Regulation (Euratom) No 3954/87 laying down maximum permitted levels of radioactive contamination of foodstuffs and feedingstuff following a nuclear accident or any other case of radiological emergency. Official Journal of the European Community. L211/1 (1989).
19. Radioactivity in Food and the Environment 13 (2007) Centre for Environment, Fisheries and Aquaculture Science ISSN 1365-6414.
20. McDonald P., Cook G.T. & Baxter M.S (1991) Natural and Artificial Radioactivity in Coastal Regions of UK. in Radionuclides in the Study of Marine Processes (Ed. Kershaw P.J. & Woodhead D.S.), Elsevier Applied Science, London.

11 KTA Marine Survey Results

UNCLASSIFIED

Station number	Location	Average dose rate (nGy/h) (n = 3)
1	South Carse	115
2	Sandyhills Bay	134
3	Port o'Warren Bay	154
4	Port Donnel	143
5	Balcarry Bay	163
6	Abbey Burn Foot	141
7	Mullock Bay	163
8	Lower Nunton Bay	126
9	Brighthouse Bay	130
10	Carrick Point	128
11	Mossyard Bay	142

Table 5. Inter-tidal environmental gamma dose rates - KTA 2008.

Sample Descriptor	Wet weight (g)	Dry weight (g)	Ashed weight (g)	Measured Activity of Dry Sample (mBq/g)				Ratio of $^{238}\text{U} / ^{234}\text{U}$
				^{238}U	^{235}U	^{234}U	Total U	
IT1	499.0	125.1	120.0	7.9 ± 1.5	< 0.5	7.1 ± 1.4	15.4 ± 2.0	1.1 ± 0.3
IT2	419.5	50.9	49.6	15.4 ± 2.4	0.7 ± 0.4	17.8 ± 2.7	33.9 ± 3.7	0.9 ± 0.2
IT3	429.1	23.0	22.6	12.6 ± 1.9	0.4 ± 0.2	12.2 ± 1.9	25.2 ± 2.7	1.0 ± 0.2
IT4	467.8	16.1	15.7	6.1 ± 1.3	< 0.6	4.9 ± 1.1	11.3 ± 1.7	1.3 ± 0.4
IT5	451.7	10.9	10.1	13.7 ± 2.2	< 0.5	13.1 ± 2.1	27.2 ± 3.0	1.1 ± 0.2
IT6	649.8	10.6	10.3	6.7 ± 1.3	< 0.5	8.7 ± 1.6	15.7 ± 2.0	0.8 ± 0.2
IT7	509.9	10.3	9.4	17.9 ± 2.8	0.7 ± 0.4	17.2 ± 2.7	35.8 ± 3.9	1.0 ± 0.2
IT8	490.5	12.9	12.4	9.0 ± 1.7	< 0.7	9.3 ± 1.8	18.7 ± 2.5	1.0 ± 0.3
IT9	469.2	11.7	11.3	6.7 ± 1.4	< 0.6	7.9 ± 1.5	14.7 ± 2.1	0.9 ± 0.2
IT10	483.8	10.9	10.6	8.4 ± 1.6	< 0.6	8.0 ± 1.6	16.6 ± 2.2	1.1 ± 0.3
IT11	512.5	14.4	13.3	14.8 ± 2.5	0.7 ± 0.4	15.2 ± 2.5	30.6 ± 3.6	1.0 ± 0.2

Table 6. Intertidal sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2008.

UNCLASSIFIED

Sample Descriptor	Wet weight (g)	Dry weight (g)	Ashed weight (g)	Measured Activity of Dry Sample (mBq/g)				Ratio of ²³⁸ U/ ²³⁴ U
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
<i>F. vesic</i> ' 2	163.1	35.3	9.8	4.9 ± 0.9	0.3 ± 0.2	5.8 ± 1.0	11.0 ± 1.4	0.8 ± 0.2
<i>F. vesic</i> ' 3	191.2	45.7	19.0	4.3 ± 0.9	< 0.4	4.7 ± 0.9	9.1 ± 1.3	0.9 ± 0.3
<i>F. vesic</i> ' 4	188.8	46.6	11.4	< 2.3	< 2.3	3.1 ± 1.7	5.3 ± 2.2	N/A N/A
<i>F. vesic</i> ' 5	234.3	54.0	12.7	5.0 ± 1.3	< 0.7	5.7 ± 1.4	10.9 ± 1.9	0.9 ± 0.3
<i>F. vesic</i> ' 6	213.4	43.2	9.6	5.2 ± 0.9	0.4 ± 0.2	5.7 ± 0.9	11.3 ± 1.3	0.9 ± 0.2
<i>F. vesic</i> ' 7	163.8	45.6	10.4	3.5 ± 0.6	< 0.2	3.8 ± 0.7	7.4 ± 0.9	0.9 ± 0.2
<i>F. vesic</i> ' 8	162.7	34.9	7.6	2.4 ± 0.5	< 0.2	2.8 ± 0.5	5.3 ± 0.7	0.9 ± 0.2
<i>F. vesic</i> ' 9	197.6	46.8	11.1	3.0 ± 0.6	< 0.2	3.4 ± 0.6	6.6 ± 0.8	0.9 ± 0.2
<i>F. vesic</i> ' 10	179.5	50.8	15.7	4.7 ± 0.8	< 0.3	4.8 ± 0.8	9.5 ± 1.2	1.0 ± 0.2
<i>F. vesic</i> ' 11	155.9	35.2	8.8	5.1 ± 2.5	< 2.5	6.1 ± 2.8	12.3 ± 3.9	0.8 ± 0.6

Table 7. Seaweed samples: alpha spectrometry analysis results showing total uranium and isotopic ratios – KTA 2008.

Sample Descriptor	Wet weight (g)	Dry weight (g)	Ashed weight (g)	Measured Activity of Dry Sample (mBq/g)				Ratio of ²³⁸ U/ ²³⁴ U
				²³⁸ U	²³⁵ U	²³⁴ U	Total U	
<i>M edulis</i> 3	113.1	28.8	2.5	2.7 ± 0.4	0.1 ± 0.1	2.9 ± 0.5	5.7 ± 0.6	0.9 ± 0.2
<i>M edulis</i> 4	96.3	23.7	1.8	3.9 ± 0.6	0.2 ± 0.1	4.8 ± 0.8	8.8 ± 1.0	0.8 ± 0.2
<i>M edulis</i> 5	81.5	19.5	1.7	4.3 ± 0.6	0.2 ± 0.1	5.1 ± 0.7	9.6 ± 0.9	0.8 ± 0.2
<i>M edulis</i> 8	98.5	25.3	2.0	3.3 ± 0.7	< 0.2	4.1 ± 0.9	7.5 ± 1.2	0.8 ± 0.2
<i>M edulis</i> 10	116.5	29.5	2.4	3.3 ± 0.5	0.2 ± 0.1	3.5 ± 0.5	7.0 ± 0.7	1.0 ± 0.2
<i>B. undatum</i> A	157.1	38.6	3.0	0.6 ± 0.2	< 0.1	0.6 ± 0.2	1.2 ± 0.2	1.0 ± 0.4
<i>B. undatum</i> B	144.0	35.3	2.9	0.5 ± 0.2	< 0.3	0.4 ± 0.2	0.9 ± 0.3	1.4 ± 1.0
<i>B. undatum</i> C	150.7	38.9	3.1	0.4 ± 0.1	< 0.1	0.4 ± 0.1	0.8 ± 0.2	0.9 ± 0.5
<i>B. undatum</i> D	147.2	38.3	2.9	0.5 ± 0.1	< 0.1	0.5 ± 0.2	1.0 ± 0.2	0.8 ± 0.3
<i>B. undatum</i> E	142.6	34.0	3.2	0.7 ± 0.3	< 0.3	0.6 ± 0.3	1.4 ± 0.5	1.1 ± 0.8
<i>B. undatum</i> F	149.3	39.1	3.0	0.5 ± 0.2	< 0.1	0.5 ± 0.1	1.0 ± 0.2	1.1 ± 0.5
<i>C. pagarus</i> A	80.5	19.9	1.9	< 0.2	< 0.2	< 0.2	0.3 ± 0.1	N/A
<i>C. pagarus</i> B	128.9	37.5	3.4	0.3 ± 0.1	< 0.1	0.3 ± 0.1	0.5 ± 0.1	1.0 ± 0.4
<i>C. pagarus</i> C	131.3	33.4	2.4	< 0.2	< 0.2	< 0.2	0.2 ± 0.1	N/A
<i>C. pagarus</i> D	105.0	29.2	2.7	0.1 ± 0.1	< 0.1	0.2 ± 0.1	0.3 ± 0.1	0.6 ± 0.3
<i>C. pagarus</i> E	156.7	60.3	5.5	< 0.1	< 0.1	0.1 ± 0.1	0.2 ± 0.1	N/A
<i>H. gammarus</i> A	134.2	36.4	2.5	< 0.1	< 0.1	0.1 ± 0.1	0.2 ± 0.1	N/A
<i>H. gammarus</i> B	106.2	25.9	2.2	< 0.2	< 0.1	0.3 ± 0.1	0.3 ± 0.1	N/A
<i>H. gammarus</i> C	151.4	40.6	2.7	< 0.2	< 0.1	< 0.1	0.1 ± 0.0	N/A

Table 8. Biota samples: alpha spectrometry analysis results showing total uranium and isotopic ratios - KTA 2008.

UNCLASSIFIED

Sample descriptor	Wet weight (g)	Dry weight (g)	Ashed weight (g)	Measured Activity of Dry Sample (mBq/g)				Ratio of $^{238}\text{U}/^{234}\text{U}$
				^{238}U	^{235}U	^{234}U	Total U	
UW1	530.1	19.4	18.6	13.2 ± 2.1	0.6 ± 0.3	10.7 ± 1.8	24.5 ± 2.7	1.2 ± 0.3
UW2	462.7	10.5	10.0	14.5 ± 2.3	0.9 ± 0.4	14.7 ± 2.4	30.1 ± 3.4	1.0 ± 0.2
UW3	516.7	12.8	12.2	11.6 ± 2.0	< 0.6	11.9 ± 2.0	24.1 ± 2.8	1.0 ± 0.2
UW4	525.2	13.9	13.3	11.6 ± 1.9	< 0.5	11.0 ± 1.8	23.0 ± 2.7	1.1 ± 0.3
UW5	535.0	11.0	10.6	12.9 ± 2.1	< 0.6	15.1 ± 2.4	28.4 ± 3.2	0.9 ± 0.2
UW6	525.0	20.2	19.3	13.0 ± 2.1	< 0.5	11.9 ± 1.9	25.3 ± 2.8	1.1 ± 0.3
UW7	535.4	18.4	17.6	12.2 ± 2.0	0.5 ± 0.3	11.9 ± 2.0	24.6 ± 2.8	1.0 ± 0.2
UW8	520.5	26.6	25.4	14.2 ± 2.3	0.7 ± 0.4	12.6 ± 2.1	27.5 ± 3.2	1.1 ± 0.3
UW9	565.4	18.7	17.9	13.6 ± 2.3	0.8 ± 0.4	13.6 ± 2.3	27.9 ± 3.3	1.0 ± 0.2
UW10	521.6	30.1	29.0	13.0 ± 2.1	0.7 ± 0.4	10.5 ± 1.8	24.1 ± 2.8	1.2 ± 0.3
UW11	503.8	16.1	15.4	15.3 ± 2.5	0.6 ± 0.4	12.9 ± 2.2	28.8 ± 3.4	1.2 ± 0.3
UW12	508.2	15.2	14.6	14.0 ± 2.4	< 0.6	11.6 ± 2.1	26.1 ± 3.1	1.2 ± 0.3
UW13	514.0	15.1	14.0	14.5 ± 2.5	0.8 ± 0.5	14.1 ± 2.5	29.4 ± 3.5	1.0 ± 0.3
UW14	535.2	21.6	20.8	12.6 ± 2.2	< 0.6	11.6 ± 2.0	24.7 ± 3.0	1.1 ± 0.3
UW15	503.8	12.1	11.6	13.1 ± 2.2	0.6 ± 0.4	12.3 ± 2.1	26.0 ± 3.1	1.1 ± 0.3
UW16	539.5	32.9	31.7	12.1 ± 2.1	< 0.6	11.3 ± 2.0	23.6 ± 2.9	1.1 ± 0.3
UW17	540.6	16.2	15.5	12.6 ± 2.2	< 0.7	10.3 ± 1.9	23.2 ± 3.0	1.2 ± 0.3
UW18	513.3	14.0	13.3	11.4 ± 1.9	0.8 ± 0.4	11.8 ± 2.0	24.0 ± 2.8	1.0 ± 0.2
UW19	504.2	15.0	14.3	18.1 ± 4.5	< 2.0	13.3 ± 3.6	31.9 ± 5.8	1.4 ± 0.5
UW20	503.6	12.7	12.1	14.4 ± 2.2	0.9 ± 0.4	14.7 ± 2.3	30.0 ± 3.2	1.0 ± 0.2
UW21	515.2	20.0	19.0	13.9 ± 2.2	0.8 ± 0.4	13.7 ± 2.2	28.4 ± 3.2	1.0 ± 0.2
UW22	505.2	17.8	16.9	13.8 ± 2.1	0.5 ± 0.3	12.6 ± 2.0	26.9 ± 2.9	1.1 ± 0.2
UW23	522.1	18.9	18.1	11.1 ± 1.8	0.9 ± 0.4	11.3 ± 1.8	23.3 ± 2.6	1.0 ± 0.2
UW24	534.1	11.9	11.4	12.8 ± 2.0	1.3 ± 0.5	11.6 ± 1.9	25.8 ± 2.8	1.1 ± 0.3
UW25	508.7	19.2	18.3	13.6 ± 2.1	0.7 ± 0.4	12.4 ± 2.0	26.7 ± 2.9	1.1 ± 0.3

Table 9. Underwater sediment: alpha spectrometry analysis results showing total uranium and isotopic ratios – KTA 2008.

Note: Activity results have been rounded to one decimal place. All uncertainties are stated at a 95% confidence level. Limits of Detection (LoD) are calculated by a ‘modified Currie formula’ at 95% (Hurtgen C., Jerome S. & Woods M. (2000) ‘Revisiting Currie – how low can you go?’ *Applied radiation and Isotopes* 53 pp 45-50). The total activity is calculated from the sum of the actual activities for each isotope, regardless of the quoted LoD. Therefore, in Tables 6 to 9, where activities are reported as less than LoD for any of the uranium isotopes, the total uranium value may not be equal to the sum of the individual isotopic values.

Historical Data

To allow historical comparison, data from KTA marine surveys from 1996 to 2008 is presented in the following Table 10. This includes inter-tidal sediment, underwater and biota samples.

UNCLASSIFIED

Sample Type	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Inter-tidal sediment	11.9-38.1 (10) (0.9-1.1)	14.3-19.0 (10) (0.8-1.2)	12.2-31.8 (10) (0.6-1.2)	11.0-32.6 (10) (0.9-1.1)	2.2- 44.7 (10) (0.8-1.2)	10.7-31.1 (9) (0.9-1.4)	12.0-31.9 (10) (0.9-1.2)	9.8-714.4 (10) (0.8-1.3)	4.2-35.0 (11) (0.9-1.3)	9.8-24.5 (11) (0.8-1.3)	7.5-29.9 (11) (0.8-1.3)	6.9-27.6 (11) (0.9-1.1)	11.3-35.8 (11) (0.8-1.3)
Seaweed	8.7-26.7 (9) (0.7-0.9)	9.8-22.7 (9) (0.8-1.1)	5.0-19.6 (9) (0.8-1.2)	7.8-14.1 (9) (0.8-1.3)	1.3 -5.1 (9) (0.9 -1.1)	6.8-12.1 (7) (0.8-1.3)	0.2-16.7 (9) (0.7-1.2)	7.2-34.2 (9) (0.7-0.9)	9.2-22.6 (9) (0.8-1.0)	4.9-18.2 (10) (0.8-1.2)	6.9-16.0 (9) (0.8-1.0)	10.7-25.6 (10) (0.5-1.1)	5.3-12.3 (10) (0.8-1.0)
Mussels	6.8-14.9 (8) (0.8-1.1)	4.3-11.4 (8) (0.8-1.1)	7.6-12.2 (5) (0.8-1.1)	6.6-11.1 (5) (0.8-1.1)	3.6 -7.2 (5) (0.8-0.9)	2.9-11.6 (7) (0.3-0.9)	6.4-10.5 (5) (0.9-1.0)	5.6-7.4 (5) (0.7-0.9)	4.0-7.8 (4) (0.8-0.9)	5.1-10.7 (4) (0.9-1.0)	2.6-12.6 (7) (0.9-1.1)	4.8-12.0 (5) (0.6-0.9)	5.7-9.6 (5) (0.8-1.0)
Whelks	x	x	2.5-3.0 (3) (0.8-1.0)	0.2-0.4 (4) (0.3-1.3)	0.6-1.3 (5) (0.5-1.6)	2.0-3.7 (8) (0.5-1.4)	1.1-39.4 (8) (0.9±0.5)	x	x	0.7-2.9 (6) (0.8-1.6)	0.1-0.1 (4) (n/a)	0.4-1.2 (8) (0.9-1.1)	0.8-1.4 (6) (0.8-1.4)
Scallops	x	2.8-3.5 (4) (0.8-1.0)	2.8-3.2 (3) (0.8-1.1)	0.7-1.1 (4) (1.3-1.5)		2.8-13.3 (8) (0.7-1.6)	x	1.5-33.0 (6) (n/a)	2.8-3.1 (2) (0.9-9.9)	2.8-13.3 (3) (n/a)	0.1-0.2 (4) (n/a)	x	x
Crabs	x	0.6 ± 0.3 (1) (0.9 ± 0.5)	x	0.3-0.4 (2) (1.5-1.7)	0.2-1.4 (8) (0.1-3.7)	1.2-2.3 (2) (0.9-1.0)	0.8-23.8 (4) (0.9 0.5 ^a)	0.1-2.3 (3) (n/a)	x	0.4-1.6 (6) (n/a)	0.1-0.6 (5) (n/a)	0.4-0.8 (3) (1.6±0.8)	0.2-0.5 (5) (0.3-0.4)
Lobsters	x	x	0.2 ± 0.1 (1) (n/a)	0.3-0.3 (2) (0.1-0.5)		2.4-3.2 (4) (0.7-1.6)	1.2-1.8 (2) (n/a)	0.3-3.4 (3) (n/a)	0.4 ± 0.1 (3) (1.0 ± 0.4)	0.4-0.9 (4) (n/a)	0.1-0.1 (2) (n/a)	0.1-0.2 (4) (n/a)	0.1-0.3 (3) (n/a)
Underwater sediment	22.3-30.0 (6) (0.9-1.2)	22.3-27.2 (6) (0.9-1.1)	21.1-28.0 (6) (0.9-1.1)	15.4-32.1 (6) (0.5-1.1)	0.2-12.8 (6) (0.8-1.3)	19.3-31.0 (34) (0.8-1.3)	15.2-30.4 (33) (0.9-1.2)	20.2-30.9 (36) (0.9-1.2)	17.0-33.2 (64) (0.8-1.2)	16.3-24.3 (18) (0.9-1.3)	16.6-26.6 (25) (0.8-1.3)	19.5-42.3 (25) (0.8-1.3)	23-31.9 (25) (0.9-1.4)

Table 10. Historical summary of total uranium and isotopic ratios for sediment and biota samples 1996 to 2008.

Notes: Values provided on the top row show the range of total uranium values in mBq/g (e.g. 1.5 – 3.2). The value in bold shows the total number of sample collected (e.g. **(12)**). The values in the bottom row show the range of ²³⁸U/²³⁴U isotopic ratios (e.g. (0.8 – 1.1)). Measurement uncertainties can be found in the annual reports.

ANNEX A Issues to be considered when interpreting or comparing uranium data

A.1 There are a number of issues that may give rise to uncertainties when interpreting or comparing uranium data as shown below. Further information is available from Reference [A1].

- analytical technique;
- statistical variation;
- spatial variability;
- temporal variability; and
- species variation (for plant and animal samples).

Analytical approaches

A.2 Sediment sample results may be reported as either dry weight or wet weight depending on whether the masses of the samples were obtained prior to or after drying. This will have implications for comparison of results between the surveys at Kirkcudbright, which are reported as dry weight and other UK uranium in sediment data, which may be reported as wet weight. Samples reported as dry weight will appear to have concentrations of uranium approximately 20% higher than those reported as wet weight (although this will depend on the moisture content).

A.3 For analysis techniques such as inductively coupled plasma mass spectrometry (ICP-MS) or alpha spectrometry the uranium present in a sample may be extracted into solution by either leaching the soil samples or totally dissolving them. Total dissolution will give rise to higher uranium results, because the analysis will include all uranium, including that which is contained within the mineral grains, whereas leached samples will only contain uranium that is either easily dissolved or is fixed to the surfaces of mineral grains. This limitation is acceptable as the primary purpose of the environmental survey is to assess any levels off DU in addition to naturally occurring uranium. However, this limitation causes a conservative bias on the $^{238}\text{U}/^{234}\text{U}$ isotopic ratios for mineral samples and other recalcitrant matrices. Total sample analysis techniques such as gamma spectrometry will give results similar to those for total dissolution. Given the differences between the results for total analysis and leached analysis care should be taken when comparing sets of data to ensure that either the same approach has been used or that difference are appropriately discussed.

A.4 Uranium concentrations in seaweed may be affected by contamination of surfaces with sediment particles. Preparation of seaweed for analysis may or may not involve a washing stage; hence it is important to be aware of the preparation approaches that have been applied when comparing the results of different seaweed analyses.

A.5 Seaweed and marine biota sample results may be reported as either dry weight or wet weight depending on whether the masses of the samples were obtained prior to or after

UNCLASSIFIED

drying. For marine biota, samples reported as dry weight will appear to have concentrations of uranium higher than those reported as wet weight. The relative increase in uranium concentration will depend on the moisture content, and can be as much as 300% for biota with high water content such as molluscs.

Statistical variations

- A.6 There will be minor variations between the true uranium content of a sample and results produced by analysis. This variation is highlighted in the counting statistics for the technique. The statistical uncertainties of laboratory results are likely to be small in comparison with the true variation in activity between samples.

Spatial variations

- A.7 DU contamination will not be uniformly distributed within a sampling area, but will depend on factors such as water flow, tidal movements and sediment drift. Repeat sampling and analysis of sediments from within an area may therefore give rise to a significant degree of variation.
- A.8 In addition to DU contamination due to firing at the range, there may be variations in uranium concentrations due to local anthropogenic or natural discharges. For example, natural uranium concentrations may be enhanced by the local application of phosphate based fertiliser to agricultural land. Veins rich in uranium minerals occur naturally along the coast of the Solway Firth, such as uraninite found at Needle's Eye, approximately 24 km away from KTA on the north coast of the estuary. These features are thought to be present across the region, although this has not been studied [A2,A3].

Temporal variations

- A.9 There will be natural temporal variations in the uranium concentration and the abundance of the various isotopes in the samples, due to seasonal variations in rainfall. Rainfall can impact on the dissolution of uranium and its migration in surface waters.
- A.10 The activities of samples from any particular sampling site may vary from year to year. This may relate to temporal changes in uranium concentration, but will also be affected by spatial variation (see above).
- A.11 Marine plant uptake of radionuclides is affected by the period in the plant growing cycle. This phenomenon also occurs in animal uptake of radionuclides within their life cycle.

Species variations

- A.12 Plant uptake of radionuclides is affected by the substrate characteristics (uranium concentration and speciation as well as other soil physico-chemical characteristics) and varies with plant species.
- A.13 Animal uptake of uranium is affected by their life habits, feeding patterns, physiology and the uranium concentration in their foodstuffs and the environment.

Annex A References

- [A1] International Commission on Radiation Units and Measurements Report 75 (2006) Sampling for Radionuclides in the Environment in *Journal of the ICRU Volume 6 No 1*
- [A2] Basham, I R; Milodowski, A E; Hyslop, E K; Pearce, J M. (1989) The location of uranium in source rocks and sites of secondary deposition at the Needle's Eye natural analogue site, D&G, British Geological Survey Technical Report WE/89/56.
- [A3] Milodowski, A E. *et al.* (1990) Uranium-mineralised micro-organisms associated with uraniferous hydrocarbons in southwest Scotland. *Nature*, 347, 465.

UNCLASSIFIED

ANNEX B Change in the ²³⁸U/²³⁴U activity ratio of a medium containing natural uranium with the addition of depleted uranium

Mass proportion of DU added *	Activity concentration (mBq/kg) **			Ratio of total activity natural uranium to total activity	²³⁸ U/ ²³⁴ U activity ratio
	U-238	U-235	U-234		
0	3.7 x10 ⁴	1.7 x10 ³	3.8 x10 ⁴	1.0	1.0
1	7.4 x10 ⁴	2.2 x10 ³	4.3 x10 ⁴	1.6	1.7
2	1.1 x10 ⁵	2.7 x10 ³	4.8 x10 ⁴	2.1	2.3
3	1.5 x10 ⁵	3.2 x10 ³	5.3 x10 ⁴	2.6	2.8
4	1.9 x10 ⁵	3.7 x10 ³	5.8 x10 ⁴	3.2	3.2
5	2.2 x10 ⁵	4.1 x10 ³	6.3 x10 ⁴	3.8	3.6
6	2.6 x10 ⁵	4.6 x10 ³	6.8 x10 ⁴	4.3	3.9
7	3.0 x10 ⁵	5.1 x10 ³	7.2 x10 ⁴	4.9	4.1
8	3.4 x10 ⁵	5.6 x10 ³	7.7 x10 ⁴	5.4	4.3
9	3.7 x10 ⁵	6.0 x10 ³	8.2 x10 ⁴	6.0	4.5
10	4.1 x10 ⁵	6.5 x10 ³	8.7 x10 ⁴	6.5	4.7
20	7.8 x10 ⁵	1.1 x10 ⁴	1.4 x10 ⁵	12.0	5.8
60	2.3 x10 ⁶	3.1 x10 ⁴	3.3 x10 ⁵	34.1	6.9
80	3.0 x10 ⁶	4.0 x10 ⁴	4.3 x10 ⁵	45.1	7.1
100	3.8 x10 ⁶	5.0 x10 ⁴	5.2 x10 ⁵	56.1	7.2
200	7.5 x10 ⁶	9.8 x10 ⁴	1.0 x10 ⁶	111.0	7.4
600	2.2 x10 ⁷	2.9 x10 ⁵	3.0 x10 ⁶	332.0	7.6
800	3.0 x10 ⁷	3.9 x10 ⁵	3.9 x10 ⁶	442.0	7.6
1000	3.73 x10 ⁷	4.8 x10 ⁵	4.9 x10 ⁶	552.0	7.6

Table reproduced from Volume 2 - Appendices, WS Atkins Environmental Assessment on DU Firings.

* The value represents the additional mass of depleted uranium added (all radionuclides) relative to the original mass of natural uranium present (3 mg U/kg soil).

** Table assumes 3 mg U/kg of natural uranium present in soil in following proportion: ²³⁸U (2.978 mg /kg); ²³⁵U (0.022 mg /kg); ²³⁴U (2e-04 mg /kg), prior to addition of DU.

ANNEX C Variability of uranium concentration and uranium isotopic ratios in marine environmental samples

- C.1 There are few specific examples that demonstrate the variability of uranium concentrations and isotopic ratios within marine environmental samples. In order to understand this variability, it is useful to consider the flux of uranium between the various components of the marine environment.
- C.2 Uranium occurs naturally in seawater and its concentration generally varies in proportion to salinity. It is present in open seawater at an average concentration of $82.5 \pm 5 \text{ Bq/m}^3$, with a $^{238}\text{U}/^{234}\text{U}$ activity ratio of 0.88 ± 0.03 at a salinity of 35 ‰. The isotopic ratio of seawater is below unity as ^{234}U is preferentially mobilised from the soil during chemical weathering, thus enhancing its presence in seawater [C1 & C2]. The physical mixing of low uranium river water¹ and high uranium sea water in estuary environments generally results in a dilution of uranium in the estuarine waters. As a result, uranium concentrations in estuarine water are mostly lower than open ocean values and increase linearly with salinity [C2].
- C.3 Concentrations of uranium in marine sediment are variable (from 32.5 to 1,625 mBq/g dry weight) and vary primarily as a function of the geology of the area.

Sample Type	Uranium Concentration	Typical $^{238}\text{U}/^{234}\text{U}$ ratio
Sea water	82.5 Bq/m^3	0.88
Estuarine water	$<82.5 \text{ Bq/m}^3$	<0.88
Marine sediment (^{238}U only)	32.5-1,625 mBq/g	0.81

- C.4 Levels of uranium also vary depending on the type of sediment present, as the physical and chemical characteristics of the sediment determine the amount of uranium which is concentrated from marine waters [C2 & C3]. Both low oxygenation² and low salinity³ provide favourable conditions for uranium scavenging from the water column by a variety of processes. These include the precipitation of uranium rich colloids into the low salinity zone and the reduction of uranium into insoluble forms [C3 & C4].

¹ Concentrations of uranium in rivers vary considerably, with carbonate and dissolved solids concentrations, with an average of $7.5\text{-}15 \text{ Bq/m}^3$, and an isotopic $^{238}\text{U}/^{234}\text{U}$ activity ratio of $0.77\text{-}0.83$ as ^{234}U is preferentially mobilised during chemical weathering.

² Low oxygenation is found with increasing depth and increasing organic content

³ Low salinity is dependent on river rate of flow, proximity to river outflow and depth of water

UNCLASSIFIED

Annex C References

- [C1] Murphy, W M; Shock, E L. (1999) Environmental Aqueous Geochemistry of Actinides, in Uranium: Mineralogy, Geochemistry and the Environment, edited by Burns, P and Finch, R. Reviews in Mineralogy, Vol 38, The Mineralogical Society of America.
- [C2] Cochran, J K. (1982) The Oceanic Chemistry of the U- and the Th-series Nuclides, in Uranium Series Disequilibrium: Applications to Environmental Problems, edited by Ivanovich, M and Harmon RS. Clarendon Press, Oxford.
- [C3] Van den Berg, C M G.; Huang, Z Q. (1984) Analytica Chemica Acta, 164, 209-222.
- [C4] Ragnarsdottir V; Charlet L. (2000) Uranium Behaviour in Natural Environments in: Environmental Mineralogy: Microbial Interactions, Anthropogenic Influences, Contaminated Land and Waste Management, Mineralogical Society Series Vol. 9, Mineralogical Society of Great Britain & Ireland.

ANNEX D Reference values for uranium in the Solway Firth and the UK

- D.1 For UK coastal sediment, ^{238}U concentrations generally range from 3.6 to 32.3 mBq/g of dry weight [D1]. Values reported for inter-tidal sediment uranium concentrations in the Solway Firth vary with location and the specific uranium rich geology of the area.
- D.2 Values for ^{238}U , ^{235}U and ^{234}U for sediment collected at Kirkcudbright in 2007 were reported by CEFAS as 10, 0.92 and 13 mBq/g of dry weight respectively[D2].
- D.3 Values reported for ^{238}U concentrations at Sandyhills Bay, in southwest Scotland, were lower than worldwide average values for seawater and sediment reported in ANNEX C. Seawater and sediment from Sandyhills Bay had ^{238}U concentrations of 40 ± 3 Bq/m and 14 ± 0.4 mBq/g dry weight, respectively [D3]. No other uranium isotopes were measured, so no indication of isotopic ratio could be provided.
- D.4 However, semi-quantitative analysis of shore sediment samples obtained from the mudflats offshore from the uraninite vein, located approximately 24 km east from KTA at Needle's Eye, yielded uranium values of up to several hundred parts per million. These activities were measured both around open, oxygenated root channels and in near surface peat material. This analysis has suggested that uranium disperses seawards and in ground waters, and is retarded in this by organic material in the sediment [D4].

Sample	Location	Activity per fresh weight (mBq/g)	Activity ^{238}U per dry weight (mBq/g)
Seaweed	[D3] Sandyhills Bay [D3] UK		3.8 ± 0.1 3.8 to 18.6
Mussel	[D3] Sandyhills Bay [D3] UK		1.1 ± 0.1 1.01 to 37.1
Mollusc	[D3] Sandyhills Bay(winkle) [D3] UK [D2] UK (mollusc & winkle) [D2] Parton (winkle)	0.89 1.3	2.72 ± 0.01 1.36 to 18.9
Crab	[D2] UK [D2] Parton	0.046 0.052	
Lobster	[D2] UK [D2] Parton	0.035 0.028	

- D.5 Literature values reported for ^{238}U in seaweed and marine biota samples for the Solway Firth area are shown in the table above; also reported are estimated values of ^{238}U from natural sources in aquatic foodstuff for the UK given by CEFAS [D2, D3]. Analysis results from Parton (near Whitehaven) are also included, although it should be noted that the uranium inventory at Parton is dominated by historical anthropogenic input of natural uranium from a local phosphate processing plant.

Annex D References

- [D1] Ragnarsdottir V; Charlet L. (2000) Uranium Behaviour in Natural Environments in: Environmental Mineralogy: Microbial Interactions, Anthropogenic Influences, Contaminated Land and Waste Management, Mineralogical Society Series Vol. 9, Mineralogical Society of Great Britain & Ireland.
- [D2] Radioactivity in Food and the Environment 13 (2007) Centre for Environment, Fisheries and Aquaculture Science ISSN 1365-6414.
- [D3] McDonald, P; Cook, G T; Baxter, M S. (1991) Natural and Artificial Radioactivity in Coastal Regions of UK. In: Radionuclides in the Study of Marine Processes, edited by Kershaw, P J and Woodhead, D S. Elsevier Applied Science, London.
- [D4] Milodowski, A E et al. Uranium-mineralised Micro-organisms Associated with Uraniferous Hydrocarbons in Southwest Scotland. Nature, 347, 465, (1990).

Initial distribution

CESO(Army) SO2 RAD (2 hard copies plus electronic copy)

House of Commons Library (Electronic copy)

Head of Radioactive Waste - Environment Group, Scottish Government (8 copies, of which 6 copies are to be placed in the Scottish Government Library.)

Aaron Scarlett

Dee Emerson

Knowledge Services (Electronic copy)

Report Documentation Form

A copy of this form is to be completed by the principal author for all Dstl reports. When complete, it is to be bound as the last numbered pages of the finished report.

1. Originators Report Number incl. Version No		DSTL/ V1	
2. Report Protective Markings and any other markings e.g. Caveats, Descriptors, Privacy markings UNCLASSIFIED			
3. Title of Report Depleted Uranium Survey Report: Kirkcudbright Training Area 2008 Part 2 Marine Environment			
4. Title Protective Markings incl. any Caveats		UNCLASSIFIED	
5. Author Aaron Scarlett			
6. Originator's Name and Address Aaron Scarlett DSTL Alverstoke c/o Institute of Naval Medicine Crescent Road Gosport PO12 2DL		7. MOD Sponsor Name and Address CESO(A) SO2 RAD Headquarters Land Forces Airfield Camp Netheravon SALISBURY SP4 9SF	
8. MOD Contract number and period covered		701418	
9. Other Report Nos.			
10. Date of Issue 04 February 2010	11. Pagination 39	12. No. of References 20	
13. Abstract Depleted uranium (DU) ammunition has been test fired at the Kirkcudbright Training Area (KTA) since 1982. Routine environmental monitoring has been carried out at KTA since 1980 to assess any environmental impact of the firings on the terrestrial and marine environments and any associated radiological hazards. This report presents the findings of the marine survey undertaken at KTA during 2008; the terrestrial survey is reported separately in Part I. No sample analysed was radioactive within the meaning of the Radioactive Substances Act 1993 nor exceeded a fraction the Generalised Derived Limits set by the National Radiological Protection Board. The findings are consistent with those of previous surveys and it is concluded that there is no radiological risk resulting from the firing of DU munitions at Kirkcudbright.			
14. Abstract Protective Marking including any Caveats UNCLASSIFIED			
15. Keywords/Descriptors (Authors may provide terms or short phrases which identify concisely the technical concepts, platforms, systems etc. covered in the report. Kirkcudbright, Depleted Uranium, Radioactivity, Environmental, Survey.			

UNCLASSIFIED

16. Report Announcement (refers to title/abstract being included in accessions lists e.g. Defence Reports Abstracts)

Announcement of this report is UNLIMITED

If there are limitations on the announcement of this report please indicate below the groups to whom it can be announced (more than one if required)

- Can be announced to MOD and its Agencies
- Can be announced to UK Defence Contractors
- Can be announced to Overseas Defence Departments
- Other (please specify)

17. Report Availability

- UNLIMITED distribution
- No Release without approval of the Release Authority (see box 7)




If the above do not apply, please indicate below the groups to whom the report may be released upon request without further Need-To-Know checks.

- Can be released to MOD and its Agencies
- Can be released to other UK Government Departments
- Can be released to UK Defence Contractors
- Can be released to Overseas Defence Departments
- Other (please specify)

18. Downgrading Instructions (check as appropriate)

- This report may be automatically downgraded to _____ after _____ years
- This report may be reviewed _____ years after publication

19. Authorisation

	Name	Signature	Date
Lead Technical Reviewer	Dr David Smith		5/2/2010
Project Manager	Dee Emerson		5/2/10
Technical Reviewer	Ron Brown		5/2/10

When complete the form is to be bound into the report to which it refers and is to form the last numbered pages of the report. Dstl Knowledge Services, Porton Down will enter an abstract and other details onto the relevant report management systems.

UNCLASSIFIED

This page is intentionally blank

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED