



5th July 2018

British Survey of Fertiliser Practice Fertiliser use on farm for the 2017 crop year

The British Survey of Fertiliser Practice is an annual survey that collects information on usage and application rates of nitrogen, phosphate, potash, sulphur, organic manures and lime on the major crops and grass grown in mainland Britain. It also includes the official statistics on annual fertiliser consumption in the UK (Table B2.6 of the full report, see link below).

This release gives key results from the 2017 survey. Full results and methodological details are published in a separate report [here](#).

Key results – Overall application rates by nutrient (compared to 2016)

The overall application rate (in kg per hectare) is based on the proportion of the crop area treated and the actual field rate of application used.

Nutrient	Tillage crops		Grassland		All crops and grass	
	Change	Rate (kg/ha)	Change	Rate (kg/ha)	Change	Rate (kg/ha)
Nitrogen	↓	137	↓	54	↓	91
Phosphate	↑	30	↓	8	↔	18
Potash	↓	37	↔	12	↓	23
Sulphur	↑	34	↔	3	↑	17

Enquiries on this publication to: David Fernall, Department for Environment, Food and Rural Affairs, Foss House, Kings Pool, 1-2, Peasholme Green, York YO1 7PX. Tel: 020 802 66202 Email: david.fernall@defra.gsi.gov.uk

These statistics are produced free from any political interference. You can find information about National Statistics on the internet at www.statistics.gov.uk.


© Crown copyright 2018. You may re-use this information free of charge under the terms of the Open Government Licence v.3. To view this licence visit <http://www.nationalarchives.gov.uk/doc/open-government-licence/version/3/>

Defra statistical releases, including details of future publication dates, and can be viewed on the Defra website at: https://www.gov.uk/government/statistics/announcements?utf8=%E2%9C%93&keywords=&topics%5B%5D=&organizations%5B%5D=department-for-environment-food-rural-affairs&from_date=&to_date=&commit=Refresh+results

1. Overall fertiliser use on crops and grass

Figure 1 shows the overall application rates of nitrogen (N), phosphate (P₂O₅) and potash (K₂O) on tillage crops and grass from 1983. Overall application rates are driven by a combination of the number of fields that receive a dressing and the rate of application for that field. Maximum usage was seen in the 1980s but there has been a general downward trend since then. The long-term decline in total nitrogen over this period is mainly due to decreased use on grassland.

Figure 1: Overall fertiliser use (kg/ha) on all crops and grass, Great Britain 1983 - 2017



2. Nitrogen

2017 saw a 3 kg/ha decrease in total nitrogen use on all crops and grassland (Table 1). This was driven by a 4 kg/ha fall in the overall rates on tillage crops and a 2 kg/ha fall in the overall rate to grassland. However, the total nitrogen use of 137 kg/ha on tillage crops remained close to the typical 140-150 kg/ha range seen over the majority of the 30 years of the survey.

Table 1: Overall nitrogen use (kg N/ha), Great Britain 2013 - 2017

	Tillage crops	Grass	All crops and grass
2013	136	59	94
2014	146	60	99
2015	146	56	98
2016	141	56	94
2017	137	54	91

3. Phosphate and potash

Table 2: Overall phosphate and potash use (kg/ha), Great Britain 2013 - 2017

Total phosphate (P ₂ O ₅)			Total potash (K ₂ O)				
	Tillage crops	Grass	All crops and grass		Tillage crops	Grass	All crops and grass
2013	28	9	18	2013	40	13	25
2014	29	10	18	2014	39	14	25
2015	29	9	18	2015	38	12	24
2016	29	9	18	2016	39	12	24
2017	30	8	18	2017	37	12	23

Table 2 shows overall phosphate and potash application rates for the past five years. Overall, fertiliser phosphate and potash use on all crops and grass has varied little over this period with rates on tillage crops about three times higher than those used on grassland.

4. Sulphur

Table 3: Overall sulphur use (kg SO₃/ha), Great Britain 2013 - 2017

	Tillage crops	Grass	All crops and grass
2013	27	2	13
2014	31	4	16
2015	31	3	16
2016	31	3	16
2017	34	3	17

Overall applications of sulphur on tillage crops increased by 3 kg/ha to 34 kg/ha in 2017. However applications on grass remained unchanged at 3 kg/ha (Table 3). This low overall rate on grass is a result of the low dressing cover, with only 10% of grass receiving a sulphur dressing.

5. Organic manures

Organic manures¹ applied to agricultural land may be produced on farm by livestock as slurries, farmyard manure (FYM) and poultry manures or imported from other sources such as treated sewage sludge (also called bio-solids) and some industrial ‘wastes’ such as compost, paper waste or brewery effluent. The nutrient levels in organic manures vary but can provide a valuable source of nitrogen, phosphorus and potassium.

In 2017, 63% of farms in the survey used organic manures on at least one field on the farm² (Table 4). Cattle manure from beef and dairy farms represents by far the largest volume of manure type generated in Great Britain. The proportion of farms using cattle FYM and cattle slurry has remained relatively stable over the last 5 years and was 47% and 16% of farms respectively in 2017.

Table 4: Numbers and percentage (%) of farms using each type of manure in Great Britain, 2017

	Farms in sample	Farms in population	Farms in population %	Volume (Mt: Mm ³)	Volume %
None	363	33,208	37%	n/a	n/a
Cattle FYM	595	42,980	47%	34.4	38%
Cattle slurry	204	14,115	16%	43.5	48%
Pig FYM	38	1,622	2%	1.5	2%
Pig Slurry	18	949	1%	1.5	2%
Layer manure	29	1,125	1%	0.6	1%
Broiler/ turkey litter	24	1,131	1%	0.4	0%
Other FYM	43	4,307	5%	1.6	2%
Other farm	13	781	1%	1.5	2%
Bio-solids	45	1,729	2%	2.9	3%
Other non-farm	30	1,395	2%	2.1	2%
Total with manure	796	57,281	63%	90.0	100%

Note: some farmers may use more than one type of manure. Mt: Mm³ are million tonnes and cubic metres.

In 2017, organic manure was applied to 25% of the area of tillage crops whereas this was 31% for grass of five years and over and 46% for grass under five years old. The majority of cattle manure and slurry was applied to grassland, reflecting the practice of utilising the manure on the farm on which it is produced.

Broadcast application is by far the predominant method of applying slurry being mostly spread on grassland. Manures applied to fields for winter sown crop are primarily treated in August and September (prior to drilling) whereas spring sown and grass fields are predominantly treated in the spring.

Where organic manures are used, applications of manufactured fertiliser can usually be reduced. Whilst the survey did not specifically ask farmers whether they adjusted manufactured fertiliser inputs because of manure use, an indication of this is possible by comparing fields that received manure with those that did not. This shows that for the major tillage crops the overall application rate of nitrogen was lower on fields which received manure. A similar trend is also seen for phosphate and potash use.

¹ The underlying sample design of the BSFP is constructed to measure manufactured fertiliser usage and therefore may not represent the population of farmers using organic manures as robustly.

² Not all manure generated by a farm is necessarily retained for use by that farm and excess manure/slurry may be exported for use elsewhere.

Further information

Survey Background and Methodology

The British Survey of Fertiliser Practice (BSFP) is the primary source of data on inorganic and organic fertiliser use in Great Britain. Its main purpose is to estimate average application rates of nitrogen, phosphate and potash used for agricultural crops and grassland. Information is also collected on applications of sulphur fertilisers, organic manures and lime. The survey data are used by Government, industry and the wider agricultural community to monitor best practice, to assess potential environmental impacts and mitigation strategies and provide important evidence to estimate greenhouse gas emissions from agriculture to inform policy.

The full Report with detailed methodological information plus separate key datasets are available on the GOV.UK [website](#).

The BSFP is a voluntary annual survey of a sample of farmers selected from a population of agricultural holdings compiled using the June Agricultural Survey. The target sample size is 1,300 farms; this sample size has been designed to achieve a statistically representative sample at the national level. Holdings of less than 20 hectares are not included in the sample. While these smaller holdings account for a significant proportion of all holdings in terms of numbers, they cover a much smaller proportion of the total area of crops and grass.

Data collection is undertaken mainly through face to face interviews with individual farmers and in 2017 the achieved sample size was 1,160. The underlying sample design of the BSFP is constructed to measure manufactured fertiliser usage and therefore may not as reliably represent the population of farmers using organic manures. The standard errors are relatively small for tillage crops, all crops and the main arable crops of wheat, oilseed rape and barley. Detailed methodology is provided in the full report.

All calculations of fertiliser rates are based on sown area of crops rather than field areas³ and results are expressed in terms of the equivalent nutrient. The overall application rate takes into account the proportion of the crop area treated and the actual field rate of application used.

Feedback

Feedback on the publication and the survey is welcome. Contact information for feedback or questions is provided on the front page of this statistical notice.

Other statistics of interest

Defra also run other surveys which may be of relevance and interest to fertiliser use and related practices through its Farm Practices Survey for England which is available of the Defra [website](#).

Data on fertiliser use are also a key element of soil nutrient balances. Soil nutrient balances provide a method for estimating the annual nutrient loadings of nitrogen and phosphorus to agricultural soils. They give an indication of the potential risk associated with losses of nutrients to the environment; losses which can impact on air and water quality and on climate change. Soil nutrient balances estimates are published under the heading “Soils” here:- <https://www.gov.uk/government/collections/agri-environment-analysis>

³ This reflects cross-compliance and environmental scheme measures where field margins remain uncropped.