










# A visual guide to vaccines

used in the routine immunisation schedule

Vaccine trade name/abbreviation	Disease(s) protected against	Age due
 <b>Infanrix hexa</b> DTaP/IPV/Hib/HepB	Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b (Hib) and hepatitis B	8,12,16 weeks
<b>Prevenar 13</b> PCV	Pneumococcal (13 serotypes)	8, 16 weeks and 1 year
 <b>Bexsero</b> MenB	Meningococcal group B	8, 16 weeks and 1 year
<b>Rotarix</b> Rotavirus (oral)	Rotavirus gastroenteritis	8 and 12 weeks
 <b>Menitorix</b> Hib/MenC	<i>Haemophilus influenzae</i> type b (Hib) and meningococcal group C	1 year
<b>M-M-RvaxPro</b> or <b>Priorix</b> MMR	Measles, mumps and rubella	1 year and 3 years 4 months
 <b>Fluenz Tetra</b> LAIV (nasal)	Influenza	2 to 9 years (routine) 10 to <18 years (children in clinical risk group)*
<b>Repevax</b> dTaP/IPV	Diphtheria, tetanus, pertussis and polio	3 years 4 months
 <b>Gardasil</b> HPV	Cervical cancer caused by human papillomavirus (types 6, 11, 16 and 18)	Girls aged 12 to 13 years
<b>Revaxis</b> Td/IPV	Tetanus, diphtheria and polio	14 years (school year 9)
 <b>Nimenrix</b> MenACWY	Meningococcal groups A, C, W and Y	14 years (school year 9)
<b>Boostrix-IPV</b> dTaP/IPV	Diphtheria, tetanus, pertussis and polio	Pregnant women from 16 weeks gestation
 <b>Pneumococcal Polysaccharide Vaccine PPV</b>	Pneumococcal (23 serotypes)	65 years
<b>Zostavax</b> Shingles	Shingles	70 years

\*Images of inactivated influenza vaccines not shown on this poster as several different products are available. Full details of the routine immunisation schedule are available at [www.gov.uk/government/collections/immunisation](http://www.gov.uk/government/collections/immunisation)