

# Having cervical screening

An easy guide about cervical screening for women aged 25 to 64



Contents: what is in this booklet		Page
	Cervical screening	3
	Cervical cancer	5
	Your choice	7
	Having the test	8
	Your test results	11
	Changes to look for	12
	More information	13

# Cervical screening



This booklet tells you about having cervical screening.



The test looks for changes in your cervix.

The cervix is inside your body at the top of your vagina. You cannot see it.

People use different words for their vagina such as private parts, front bottom or down below.



Cervical screening looks for a virus in the cervix.

For a very small number of women the virus can be serious and cause cervical cancer.

This is a cancer that starts in the cervix. It is a very serious illness that you can die from.



Having the test can stop you getting cervical cancer.

Having cervical screening saves thousands of lives every year.



If you are aged 25 to 49, the NHS will offer you the test every 3 years.

111111



If you are aged 50 to 64, the NHS will offer you the test every 5 years.

# Cervical cancer



The virus that can cause cervical cancer is called HPV.

Most women will get the virus at some time in their lives.

Most women who have the virus do not know they have it.





You can get the virus from any kind of sex or sexual touching with a man or a woman.

You can have cervical screening even if you have never had sex.





Children are offered the HPV vaccine when they are 12 or 13.

This helps to protect against the virus that causes cervical cancer.

You may have had the HPV vaccine when you were younger.



Having cervical screening is important even if you had the HPV vaccine.

It is the only way to find out if your cervix is healthy.

# Your choice





It is your choice to have cervical screening or not.



Good points

Screening can find the virus that causes cervical cancer.

Screening can find unhealthy cells.

We can remove unhealthy cells before they turn into cancer.



#### Bad points

Sometimes screening can miss unhealthy cells.

Some women are sent for more tests and treatment they do not need.



You can talk to your family or carer. You can talk to a nurse or your doctor.

# Having the test



You will get a letter asking if you want to book your cervical screening.



You should try to book a time and day when you are not having your period.

Do not use any creams on your vagina in the 2 days before your test.



Your cervical screening is usually done by a female nurse or doctor.

You can ask to make sure a female does your test when you book your appointment.





Wearing a skirt or dress can make the test easier.



You can take a female friend, carer or relative with you.

You can ask for a longer appointment.



You will go into a private room.

The nurse will check your name and age.

You can ask questions.



The nurse or doctor will ask you to take your knickers off.





The nurse or doctor will ask you to lie on a bed with your knees bent.

They will ask you to open your legs.

You will have a paper sheet over you.



The nurse or doctor will use a speculum to put in your vagina so they can see your cervix.



They will then use a small brush to take a sample of cells from your cervix.

It should not be painful but it may be uncomfortable.

You can stop the test at any time.



The test takes only a minute or two.

The whole appointment usually takes about 10 minutes.

## Your results





Your sample is tested to look for the virus and any unhealthy cells.



You will get your results in a letter or someone will telephone you.

9 out of every 10 women have a normal result.



This means your cervix is probably healthy.

1 out of 10 women might need to have more tests.

This means you may have unhealthy cells or the virus.



If you need to have more tests, your result letter will tell you what to do.

# Changes to look for





You should go to your doctor if you notice any of the following changes at any time.



Bleeding from your vagina when it is not your period.



Bleeding after your periods have stopped.



Pain or bleeding during or after having sex.



Any unusual fluids from your vagina.

# More information





You can telephone your doctor.



You can visit the NHS.UK website: www.nhs.uk/cervical



You can talk to your doctor or nurse.

You can talk to your family or carer.

First published: 2019 This version: June 2024 © Crown copyright 2024 Re-use of Crown copyright material (excluding logos) is allowed under the terms of the Open Government Licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/version/2/ for terms and conditions. Images courtesy of: Public Health Wales: illustrations p12 Photosymbols: first image p4, first p5, second and third p7, second and third p8 Inspired Services: second image p3, second, third and fourth p5