

Animal & Plant Health

Procedures for Official Examination of Valuet for Cultivation and Use (VCU) Harvest 2005 Sugar Beet for Cultivation and Use (VCU) Harvest 2029 Sugar Beet April 2019 Changes from Harvest 2018 VCU procedures

- 1. p1, A.3.2.2, Trials Organiser: added responsibility
- p4, A.4.1, Resistance to harmful organisms: percospora and Stemphyllium added.
 p6, C.2.8, paragraph added.
- 4. p7, C.3.2.1, amended for clarity.
- 5. p8, C.3.4.7, amended for clarity
- 6. p8 & p10, C.4.2 & C.4.3, Fertiliser amended to Pesticide.
- 7. p 11, C.4.10, Post triates bandry: added.
- 8. p11, C.5.1.2, paragraph deleted and subsequent re-numbering.
- 9. p17, C.6.3.3 to 15, "Assessment to be made before the application of fungicide" deleted.
- 6.3.3.16, Cercospora added.
- C.6.3.3.17, Stemphyllum added.
- 2. p23, Appendix 2, Seed Treatment: amended.
 - 13.p25, Appendix 4, Trial Locations: Fotheringhay replaces Holme, Theveton replaces Hibaldstow and Brockford replaces Sandringham.
 - 14.p26, Appendix 5, Controls: Cantona KWS, BTS 860, Daphna and BTS 3325 replace Cayman, Hornet, Salamanca KWS and Aurora.
 - 15.p32, Appendix 10 Control of Groundkeepers: added

Contents

| Section A - General Information | 1 |
|--|----|
| Section A - General Information A.1 Purpose A.2 Scope A.3 Responsibilities A.4 Summary of Growing Trials, Tests and Assessments Procedures Section B – Seed Handling Procedures B.1 Responsibilities B.2 Seed Handling Procedures B.3 Authentication of VCU Seed Section C – Growing Trial Procedures C.1 Responsibilities C.2 Site Suitability | 1 |
| A.2 Scope | |
| A.3 Responsibilities | |
| A.4 Summary of Growing Trials, Tests and Assessments Procedures | 5 |
| Section B – Seed Handling Procedures | 5 |
| B.1 Responsibilities | ξ |
| B.2 Seed Handling Procedures | Ę |
| B.3 Authentication of VCU Seed | ξ |
| Section C – Growing Trial Procedures | 6 |
| C.1 Responsibilities | 6 |
| C.2 Site Suitability | 6 |
| C.3 Sowing the Trial | 6 |
| C.3 Sowing the Trial. | |
| C.5 Harvesting | 1(|
| C.6 Records | |
| Section D - Disease Testing Procedures | 18 |
| D.1. Assessment of Natural Infection | 18 |
| Section E - Quality Testing Procedures | 19 |
| E.1 Responsibilities | 19 |
| Quality Assessment Methodology for Obligatory and Additional Tests | 19 |
| Quality Assessment Methodology for Obligatory and Additional Tests | 20 |
| F.1 Plan Validation and Storage | |
| F.2 Data Recording | 20 |
| F.3 Other Tests and Trials | 2(|

| Appendix 1 Approved Trial Organisers/ Operators for Sugar Beet | 21 |
|--|-------------|
| Appendix 2 Seed Treatment Products for Use on NL Trials – to be advised | 22 |
| Appendix 3 Seed Delivery Deadline Dates | 23 |
| Appendix 4 VCU Growing Trial Operators and Trial Locations for Sugar Beet | 24 |
| Appendix 4 VCU Growing Trial Operators and Trial Locations for Sugar Beet Growing Trial Operators/Seed Handling Operators Appendix 5 Control Varieties for VCU Assessments for Sugar Beet Appendix 6 Fungicide Programme for Sugar Beet Appendix 7 Dates by which Records Should be Sent to the Trial Design and Data Ha | 24 |
| Appendix 5 Control Varieties for VCU Assessments for Sugar Beet | 25 |
| Appendix 6 Fungicide Programme for Sugar Beet | . |
| Appendix 7 Dates by which Records Should be Sent to the Trial Design and Data Ha | nding 27 |
| | 28 |
| Leaf diseases | 28 |
| Infection Disease Severity Description | 28 |
| Appendix 9 Growth Stage Key of Beet | 29 |
| Longer in USE. See | |
| this document is no longer in | |

Section A - General Information

A.1 Purpose

For the latest procedure A.1.1 This document sets out the approved procedures to be used for growing trials, tests and assessments as required by the current Protocol for Official Examination of Value for Cultivation and Use for Sugar Beet.

A.2 Scope

A.2.1 These procedures apply to all varieties of Sugar Beet.

A.3 Responsibilities

A.3.1 Procedures Development Group

The Procedures Development Group is responsible for reviewing these procedures annually and making amendments for which it has responsibility, in accordance with the provisions of the VCU Protocol.

A.3.2 Organisers and Operators

A.3.2.1 Trials Organiser

,ee G British Society of Plant Breeders Ltd (BS BSPB House 114 Lancaster Way Business Tel No: Elv 01353 653846 Cambs Fax No: 01353 661156 Email: jeen y.widdowson@bspb.co.uk CB6 3NX

A.3.2.2 The Trials Organiser is responsible for ensuring all VCU Protocol and Procedures requirements are followed and liaison with all Operators carrying out trials for National List purposes, including any new characters, supply of seed and data handling.

A.3.2.3 Data Nandling Operator. The Data Handling Operator identified by the trials organiser is esponsible for trial design and data validation in accordance with the VCU protocolorid associated procedures.

4 Growing Trial Operators, Seed Handling Operators and Quality Testing Operators. Trials Organiser is responsible for identifying potential Growing Trial Operators and Quality Testing Operators to carry out trials and tests as determined by the Procedures Development annual review in accordance with the VCU Protocol, and these Procedures. The Trials Organiser is also responsible for finding Seed Handling Operators who are able to carry out seed handling. Seed Handling Operators prepare trial seed for sowing on behalf of any Growing Trial Operator in accordance with the VCU Protocol and these Procedures.

A.3.2.5 A list of all approved Organisers and Operators is shown in Appendix 1.

A.3.3 VCU Protocol and Procedures non-compliance

A.3.3.1 The Trials Organiser will forward any reports on VCU Protocol or Procedures noncompliance to APHA within 1 week of receipt. The Trials Organiser will obtain A.3.5.1 The Trial Design and Data Hand¹¹ Deprator and Seed Handling 5 arry out their of authorisation from APHA for any actions, including those necessary to remedy non-

A.3.5.2 The Seed Treatment Operator is responsible or organising the processing of seed of candidate varieties submitted by the applicant, and seed of control, or other reference varieties, in accordance with the requirements set out in these Procedures and the current VCU Protocol. The Trials Organiser will ensure that any seed treatments or additives are approved for the purpose. Approved products are listed in Appendix 2.

A3.5.3 The Seed Handling Operator with receive seed from the Seed Treatment Operator for distribution to the Growing Trials Operators.

A.3.6 Dispatch of Seed

A.3.6.1 The Seed Handling perator will arrange for seed to be dispatched by the agreed deadlines to the Growing Trial Operators, and, for authentication, to the DUS testing centres including, where appropriate, foreign testing authorities. Dates are given in Appendix 3.

A.3.7 Monitoring of Growing Trial Operators and Seed Handling Operators -Documentation

The Trials Organiser will take any necessary action to enforce deadline dates and quality standards for required documentation.

XA.3.7.2 The Trials Organiser will ensure Growing Trial Operators and Seed Handling Operators have access to all current protocols and procedures relevant to them and that they are notified of any amendments.

A.3.8 Seed Quantities

A.3.8.1 The Trials Organiser will determine the quantity of seed required for all VCU tests and trials in each annual series, including authentication, and will notify the applicant of quantities and delivery addresses.

A.3.9.1 The Seed Handling Operator is responsible for ensuring all seed is clearly labelled cedure with variety name/breeders reference and AFP number.
A.3.10 Seed quality
A.3.10.1 Seed submitted for VCU testing must meet the standards for the standards for the appropriate seed results. of seed given in the appropriate seed regulations, in respect of germination, analytical purity and content of other seeds and any other impurities. A.4 Summary of Growing Trials, Tests and

Assessments Procedures

A.4.1 The number of trials and site locations are as detailed in Appendix 4.

A.4.2 Control varieties are listed in Appendix 5.

A.4.3 The Trials Organiser is responsible for informing the Growing Trial Operators of the additional characters, which must be recorded as and when requested by applicants, and any samples that may be required for analysis.

rec for an instance in us

A.4.4 VCU trial assessments required

Bold = Obligatory *Italics = Additional only if requested by the applicant*

A.4.4.1

| Type of Character | Reference | Description of assessment Plot root yield Emergence % Plant uniformity Early vigour |
|--------------------------------------|---|---|
| Behaviour with respect to factors in | Section C Section C | Plot root yield Emergence % |
| the physical environment. | Section C | Plant uniformity |
| ine physical charteria. | | Early vigour |
| | | Population count (required for Bolters%) |
| | | Bolters % |
| | | Drought stress |
| | | Bolters % Drought stress Top size Crown height |
| | | |
| * Resistance to harmful organisms | Section D | Virus yellows % |
| | | Downy mildew % |
| | | Powdery mike % |
| | | Rust % Ramulaya % |
| | | Cercospora % |
| | | Stemphyllium % |
| | | |
| Quality characteristics | Section E | Sugar content % |
| | Q. | Total estimated impurities |
| | | |
| I here is a requirement to record c | - | dance with these VCU Procedures. |
| urther Measurements | SO. | |
| | 15 | |
| | v recorded in al | trials following procedures in Section |
| be following must be measured | | |
| | ••••••••••••••••••••••••••••••••••••••• | ý 31 |
| The following must be measured of | •••••••••• | |

Sowing Date Harvest date Plot size Harvest losses (where present at levels which will affect results)

Section B – Seed Handling Procedures

B.1 Responsibilities

procedure B.1.1 The Seed Treatment and Seed Handling Operators are responsible for carrying out the following procedures.

B.1.2 Raw seed is supplied direct to the Seed Treatment Operator for treating and pelleting.

B.1.3 Treated and pelleted seed is supplied by the Seed Treatment Operator to the Seed Handling Operator for distribution to the Growing Trials Operators.
B.2 Seed Handling Procedures

B.2.1 The Seed Treatment and Seed Handling Operators will receive a sowing list from the Trials Organiser, along with instructions as to which seed treatments or additives may be used. A list of chemicals approved by the Procedures Development Group is at Appendix 2.

B.2.2 The Seed Treatment and Seed Handling Operators must record receipt of seed from applicants by checking it off against the sowing list as it arrives. APHA should be notified of any damage to the packaging, loss of seed or problems that would affect the validation of the trials.

B.2.3 Once seed has been treated, it must be kept safely until required for drilling, authentication and quality control.

B.2.4 The Seed Treatment Operator must record use of treatment chemicals in accordance with best practice and in full observance of all manufacturers' recommendations and relevant statutory obligations.

B.2.5 Any seed treatment equipment used must be fit for the purpose, properly calibrated, set up and operated in accordance with the manufacturer's recommendation.

B.2.6 Cross, contamination must be avoided by ensuring equipment is clean between weighing and treatments.

Cord must be kept of chemicals used and date of treatment.

B23 Seed treatment should take place as near to the drilling date as possible.

B.3 Authentication of VCU Seed

B.3.1 Results from the second year's submission will be compared, by the DUS centre, with the first year for authentication purposes.

Section C – Growing Trial Procedures

C.1 Responsibilities

... one trials according to providing a suitable site, which proceeding to providing a suitable site, which meets the following criteria:
C.2.2 Previous cropping - no Sugar beet or other *Beta* species to be grown with the training of t

fertility and texture should be uniform across the site. The soil should be sufficiently uniform to avoid variation in the growth of the trial.

C.2.4 The trial should be sited away from trees, hedges headlands and other features, which are likely to cause uneven growth or encourage gazing damage from fauna.

C.2.5 The trial area should be cultivated in the dipection of ploughing and drilled across the direction of ploughing and cultivation such that ach plot receives similar treatments. Cultivations should follow best local practice

C.2.6 The frequency, direction and according to the state of all cultivations carried out since the last crop should be recorded in the site details record sheet.

C.2.7 Organic manure should not be applied to the trial area after the preceding crop unless they can be applied durately and evenly across the trials area. The use of Fresh Farm Yard manure is not permitted.

It remains the responsibility of the trials manager to maintain the integrity of the trial.

C.2.8 Be aware that some varieties may have stewardship requirements. These will be provided in Appendix 1.

ving the Trial

3.1 Plot Size

C.3.1.1 Three rows are to be drilled at 0.5m row width, with the same row width between plots. Variation in row spacing of more than 10% between adjoining plots should be notified to the Trials Organiser. All rows of the plot will be harvested for yield and the plot size should be sown to allow a minimum target harvest plot, after trimming, of 10m². A minimum of 3m pathway between plot ends is required to facilitate machine harvesting. There will be four replicates sown.

To allow access for harvesting equipment a headland of a minimum of 24m is required around the trial

C.3.2 Plant population

C.3.2.2 Trials should be gapped as near as possible at the 2-4 true leaf stage to give a uniform plant population of approximately 100,000 evenly-distributed plants per hectare.

establishment is uneven the Growing Trials Operators should contact the Trials Organiser for guidance.

C.3.3 Trial layout

*01 C.3.3.1 The Trials Organiser, following consultation with APHA produces provisional sowing lists. The Trials Organiser will make final sowing lists available to Growing Trial Operators, along with the trial plans produced by the Trial Design and Data Handling Operator.

C.3.3.2 The trial should be sown according to the plan produced by the Trial Design and Data Handling Operator and may be an incomplete block design. In an incomplete block design each replicate is split into a number sub-blocks. Any splitting of replicates must be between sub-blocks and not through sub-blocks. Varieties can be moved within a subblock but must not be moved from their sub-block. Varieties must not be moved around within the plan eg if drilling errors occur. If plots are moved out of their original sub-block they will have to be treated as missing plots. The Trials Organiser must be informed immediately if there are any departures from the original plan or if there any other anomalies.

C.3.3.3 If there is a new replace a planned variety eg if varieties are withdrawn, affected plots must be sown with any of the standard control varieties. Any such replacements must be agreed with the Trials Organiser. The control varieties are listed in Appendix 5

C.3.4 Drik

Trills must be set up and calibrated in the field before commencing drilling.

4.2 Care must be taken with drill settings and drilling speed to ensure satisfactory and Uniform establishment and plant population from plot to plot. It is also important to ensure that there is no-carry over of seed between plots.

C.3.4.3 Six rows of discard of the same variety, or 2 x 3 rows of 2 different varieties, should be drilled on either side of the trial with the same drill and at the same time that the trial is drilled

C.3.4.4 Precautions must be taken to avoid any missing rows. Any missing rows or parts of rows must be noted on the drilling plan and reported to the Trials Organiser within one month of emergence.

C.3.4.5 Seed is supplied for trial purposes only. Unused seed must not be supplied to

C.3.4.6 Seed disposal All surplus packets and discard trial seed must be returned to the Seed Handling Operator for disposal and the date and quantity returned noted in the Trial Diary. C.3.4.7 Pathways A gap (pathway) between plots of at least 3m is required to avoid carried to avoid carried harvesting equipment. The gap (pathway) should effects on the beet at the effects on the beet at the end of each plot and improves their harvestablity. Beet in the pathways must either be harvested or destroyed by any appropriate method that aims to create a level pathway which is free from beet and any weeds which will affect harvesting. Rotavating or cultivating pathways late in the season should to be carried out as it may create soft ground conditions that adversely affects harvesting. Only under exceptional circumstances may this be considered and only after seeking agreement from the Trials Organiser. Pathways must be gleaned pre-harvest for beet or beet fragments that are of a harvestable size.

At least one row of the cross drilling not adjacent to plot ends should contain a rhizomania susceptible variety. Seed to be supplied by the Trials Organiser.

C.3.5 Confirmation of trial layout

C.3.5.1 After the trial has been drifted, the Growing Trial Operator must:

- Confirm the drived plan by transmission of the sowing date and plan to the a) Trial Design and Data Handling Operator with any amendments to the plan clearly indicated.
- Despetch a map of the site location, showing major roads and entry point to b) e site, as well as a detailed ground plan of the trial to the Trial Design and this document Data Handling Operator.

C.4 Husbandry

C.4.1 Agronomy

All spraying activities must take account of The Plant Protection Products (Sustainable of Practice for Using Plant Protection Products (as amended). Applications of pesticides and sprays should be uniform. These applied across the direction of the plots.

C.4.3 Fertiliser application

"the Applications of fertilisers should be uniform and take into account in Kerent fertility, previous cropping, winter rainfall, the best local practice. All fertiles applications should take account of the AHDB Nutrient Management Guide (RB209), the corresponding advisory publications in England, Wales, Scotland and Northern Ireland and past trialling experience.

C.4.4 Herbicides

Chemicals must not be used if there are any known varietal sensitivities. If in doubt, the Trials Organiser should be consulted.

15⁰

C.4.5 Pest and Disease Control

C.4.5.1 Pest Control

Appropriate seed dressings of the applied as approved by the Trials Organiser. The chemical seed treatment a bied to the trial seed should control some soil-borne pests and may provide some early season control of insect-borne-virus vectors. However, appropriate pesticide meatments should be undertaken to control virus vectors through the season. Precautions should be taken against attacks by, for example, birds, deer, rabbits, hares, mice and insects.

C.4.5.2 Disease control

Seeding diseases should be controlled by the routine seed-dressings used and viruses ted with a fungicide according to the instructions in Appendix 6. should be controlled by targeting their insect-vectors (see C.4.5.1 above). Trials should be

C.4.6 Irrigation

Irrigation will not be permitted without the specific agreement of the Trials Organiser.

C.4.7 Plot assessment

Plots should be assessed between the 6-8 and 8-10 true leaf stage and scored on a scale

1= treat as missing plot 5= may be reliable 9= likely to be reliable

Please provide comments to explain the 1 and 5 scores.

C.4.7.1 Missing plots - Plots with gaps or poor uniformity may occur

If plots are weak due to mechanical or agronomic problems throughout their entire length it may be necessary to make the plots missing. The adjacent plots may have to be missing due to unfair advantage to their growth. These plots should be entered in subsequent data records as "*" (see C.6.2.5) and should be clearly indicated when the data is sent to the Trial Design and Data Handling Operator.

Where possible 'gapping-up' outside rows of missing plots should be considered to try to avoid an edge effect on the unaffected neighbouring plot. The plots should be clearly marked when the data is sent to the Trial Design and Data Nandling Operator.

C.4.8 Weed beet

Weed beet should be removed from plots before they become competitive.

C.4.9 Trials not taken to plot harvest

It is the responsibility of the Growing Totals Operator to discuss with the host grower the forward management of trials not taken to plot harvest. If herbicide tolerant varieties are included in the trial then a procedure must be put in place to ensure that bolters are removed and destroyed to prevent pollen release and regrowth. The method and visits should be noted in the Trial Wary.

C.4.10 Post trial husbandry (See Appendix 10)

C.5 Harvesting

C.5.1 Timing of harvesting

C.5.1 Date of harvesting will be according to a schedule which will be drawn up by the Trial Harvesting Contractor after consultation with the Growing Trial Operators.

C.5.1.2 Pathways (see C.3.4.7 above).

procedure

C.5.2 Harvesting method

The harvesting operation is based on a single pass system comprising a tractor with a front mounted 6 row flail defoliator and a trailed 6 row lifter specifically designed to harvest est procedure and bag two 3 row plots at a time. The plot bags are sealed, labelled and delivered to the Trials Tarehouse, which is located at the Commercial Tarehouse, for washing, yield and quality analysis.

C.5.3 Samples

- 1. Within **24 hours** of being harvested the plots are delivered to the Trials Tarehouse.
- 2. Within **48 hours** of being received at the Trials Tarehouse the plot washed, assessed for yield and a sub-sample taken. This sub-sample is used to produce the brei sample for quality analysis.
- 3. Within **24 hours** of being washed, weighed and sample the washed plot subsample is assessed for Quality through the Commercia Tarehouse Laboratory.

Details of the quality analysis process are described in sec

C.5.4 Submission of data

C.5.4.1 Appendix 7 lists the records, with deadlines, to be sent to the Trials Organiser. Diary sheets and other field records should be returned to the Trials Organiser within 5 working days of harvest.

C.5.4.2 All plot records should be transmitted to the Trial Design and Data Handling Operator following the deadlines set out in Appendix 7. The Growing Trial Operator should ensure that data are free for errors before transmission. After scrutiny, copies of results will be returned to the Growing Trial Operator for action as agreed by the Trial design and Data Handling

uing, uing, phis document is no lon

C.6 Records

Records should be clear and self-explanatory so that the trial can be carried on at a moment's notice by another person without difficulty.

C.6.1 There are four components:

- 1.
- Field notes of trial status, recording and inspections
 Site data 1 Site details including site sketch map and location, previous cropping, soil analysis, cultivations and drilling.
 Site data 2 Site details including fertiliser and sprays, herbicker, fungicides, insecticides and harvest.
 lot records Plot data.
 y in the Dimensional status, recording and inspections 2.
- 3.
- 4. Plot records Plot data.

C.6.1.1 An entry in the Diary sheet should be made on every tral visit and anv observations relevant to variety performance should be recorded. If the trial is in good condition, with no problems, this should be recorded.

C.6.2 Plot records

C.6.2.1 Plot data may be recorded direct onto a stata logger using a system approved by the Trials Organiser or recorded on paper the entered and validated onto a computer using an approved system. A system of ensuring that data are recoverable, in the event of loss of original data, must be implemented, eg copy and safe storage. Whichever method is used, individual plot data will only be accepted by the Trial Design and Data Handling Operator in an approved format using the measure names and units as listed in Section C.6.3.

C.6.2.2 All observations should be checked at the time of recording to ensure that they lie within acceptable limits for the character recorded. Observations that have been designated as exceptional by the recorder should be identified with a note on the approved data file or hard commedium describing the possible causes together with a recommendation for their exclusion or inclusion in the trial analysis.

C.6.2.3 Ple numbers on record sheets must correspond with the numbering on the field plan.


4 If a character is not recorded or is missing the Growing Trial Operator should in the diary or on the recording sheet the reason why it has been excluded.

C.6.2.5 Where a plot record is missing the Growing Trial Operator should enter "*" in the approved data file or hard copy medium and, unless the non-recording of the plot has already been agreed with the Trials Organiser, append a note to the file explaining why a missing value has been entered for that plot. The Growing Trial Operator must not enter "0" for missing plots.

C.6.2.6 Specific plot records should be made as counts or on the scales shown for each character.

C.6.2.7 All records should be returned to the Trial Design and Data Handling Operator immediately after recording. Indicative deadlines are given in Appendix 7.

C.6.3.1 The following procedures must be followed for measuring all characters to be used in NL decision-making. C.6.3.2 Characters to be Recorded for DUS Purposes. These records will be undertaken by the DUS Operator. The Triple Decision and the triple Decision of the Grand the triple Decision of the Grand the triple Decision of the triple D and that their sites should be made available to the DUS Operator. NB and of the sites will also be used to obtain leaf measurements for DUS purposes which will involve access to the site by a team from the DUS Operator over several days.



9 = dark green

C.6.3.2.5 **LEAF WAVING** from all plots (OBLIGATORY) (1-9)Relatest procedure Record the waviness of the leaf margin in July 1 = smooth9 = very wavy.C.6.3.2.6 LEAF BLISTERING (OBLIGATORY) from all plots Record the blistering of the leaf surface in July 1 = smooth9 = very blistered.C.6.3.3 VCU Characters to be Recorded for VCU Purposes. C.6.3.3.1 The following procedures must be followed for measuring characters to be used in NL decision-making. C.6.3.3.2 EMERGENCE from all plots (COUNT) When the most advanced variety has 2 true leaves, count the number of plants in the centre row of each plot and record the length of row dounted and seed spacing so that percentage emergence can be calculated. from all plots (OBLIGATORY) C.6.3.3.3 PRE-GAPPING POPL (COUNT) Record the number of plants from the cantre row in the harvested plot area at the 2 to 4 true leaf stage pre gapping. Doubles hould be counted as one. Record the plot length and indicate any rows that have a we population. ULATION COUNT from all plots (OBLIGATORY) C.6.3.3.4 POST-GAPPING (COUNT) Record the number of Tants of the 3 rows in the harvested plot area, post gapping, from the 6-8 true leaf stage onwards. Record the plot length and indicate any rows that have a low population *NT UNIFORMITY* from all plots (ADDITIONAL) (1-9)C.6.3.3.5 A plant uniformity within the plot in late May/June on the scale: large variation $\overline{9} = \text{small variation}$

C.6.3.3.6 EARLY VIGOUR

(1-9)

Record on the basis of relative plant size in late May/early June on the scale:

1 = very small 9 = very large.

Record the size of the plants at each end of the scale.

C.6.3.3.7 BOLTER NUMBER from all plots (OBLIGATORY)

(COUNDOCECITIC Bolters are a potential source of weed beet if flowering occurs early enough for viable s to be produced. The following procedures ensure that the number of bolters occorring are recorded and pollen release, seed set and shed is prevented.

If herbicide tolerant varieties are entered into the trial these plots should be inspected for bolters every 2 weeks until harvest to prevent pollen release. Bolters to be stripped and flowering parts destroyed to prevent any further regrowth or pollen release. Bolter inspection visits and any actions should be noted in the Trial Diar

The number of bolters in the harvested area should be counted at:

- The end of June count all bolters. Flowering branches to be stripped to a) leave the main stem. Ensure that flowering parts are destroyed to prevent any further regrowth or pollen release. The root should remain as part of the harvest plot.
- The end of July count all being including those previously counted. b) Flowering branches to be stripped to leave the main stem. Ensure that flowering parts are destroyed to prevent any further regrowth or pollen release. The root should remain as part of the harvest plot.
- About the time charvest count all bolters including those previously c) counted.

If extra bolting counts are been say follow procedures b) above and send the additional records, clearly indicating the date on which the counts were taken, to the Trial Design and Data Handling Operator.

IT STRESS C.6.3.3.8 DRC

from all plots

(ADDITIONAL)

This should be assessed on a plot basis when varietal differences become apparent and Ought stress is uniform across the trial. where

vere symptoms no symptoms

If areas of the trial become severely affected by drought, these should be recorded on a trial layout and returned to the Trial Design and Data Handling Operator as soon as possible. The Trials Organiser should also be informed.

C.6.3.3.9 TOP SIZE

from all plots (ADDITIONAL)

Estimate the relative top size on a plot basis prior to harvest.

the latest procedure 1 = very small 9 = very largeEnsure that records are made before severe frost. C.6.3.3.10 HEIGHT OF CROWN from all plots (ADDITIONAL) Record height above soil level on a plot basis prior to harvest 1 = Lowest leaf scar at soil level 9 = Lowest leaf scar very high above soil level Measure the actual height for extreme values used. C.6.3.3.11 VIRUS YELLOWS % from all plots (OBLIG Y if present) (%) Virus yellows % should be assessed on a plot basis by estimating the percentage number of plants showing infection in August. **WELIGATORY** if present) (COUNT) C.6.3.3.12 **DOWNY MILDEW** from all plots Downy mildew should be assessed on a plot basis by recording the number of plants showing any symptoms. Should the lever infection become severe inform the Trials Organiser. C.6.3.3.13 POWDERY MILDEW from all plots (OBLIGATORY if present) (%) Inspect in July/August/September and record on a plot basis if the disease is > 5% in the most susceptible variety or imprection loci present. Use the NIAB Sugar beet foliar disease key in Appendix (OBLIGATORY if present) C.6.3.3.14 RUST from all plots (%) Inspect in July/August/September and record on a plot basis if the disease is > 5% in the most susceptible variety or if infection loci present. Use the NIAB Sugar beet foliar disease ken h Appendix 8. from all plots (OBLIGATORY if present) 5 RAMULARIA (%) pect in July/August/September and record on a plot basis if the disease is > 5% in the most susceptible variety or if infection loci present. Use the NIAB Sugar beet foliar disease key in Appendix 8. C.6.3.3.16 CERCOSPORA from all plots (OBLIGATORY if present) (%)

Inspect in July/August/September and record on a plot basis if the disease is > 5% in the most susceptible variety or if infection loci present. Use the NIAB Sugar beet foliar disease key in Appendix 8.

C.6.3.3.17 STEMPHYLLIUM LEAF SPOT from all plots (OBLIGATORY if present) (%)

Inspect in July/August/September and record on a plot basis if the disease is > 5% in the most susceptible variety or if infection loci present. Use the NIAB Sugar beet foliar disease key in Appendix 8.

C.6.3.3.17 ROOT YIELD from all plots (OBLIGATORY)

redure Root yield of each plot should be recorded after washing at the trials tarehouse. Record the weight in kilograms of the clean beet.

Yield data should be sent to the Trial Design and Data Handling Operator within 2014 harvesting the trial.

The following information should accompany the yield data. This should eithe submitted with the yield data to the Data Handling Operator or where appropriate before harvest by the Growing Trials Operator:

- Lifting date of the trial and processing date in the tarehouse. 1.
- 2. Plot length: the plot length harvested in metres.
- Plot width: the width of the harvested plot in methes from outer row to outer row 3. plus half of the inter-plot gap on either side
- Missing plot information. 4.
- All records and material changes made during the lifting and tarehouse 5. process.
- Any other information that may 6. affected the yield.

C.6.4 Site Factors

Any factors which may have affected the yield of the trial or individual plots must be noted and taken into account when validating the trial.

Attacks are observed, then plot records should be made in If any other pest or disease accordance with the processive in Section D for disease.

Plot records for other scores should be taken as plants affected on a 1 to 9 scale. Include definitions for each rating on the 1 to 9 scales.

C.6.5 Trial hospection

All triats will be inspected by the Trial Inspection Operator and Technical Validation Operator and, in some cases, it may be necessary to visit on more than one occasion.

Trial Operators in respect of inspections are to:

- Give inspectors reasonable access to trials
- Provide the inspector with information (for example pesticide sprays applied etc) at the time of inspection if requested.
- Co-operate with the inspector in making any non-routine assessments required to establish the validity of the trial (for example population counts)
- Carry out any action agreed in consultation with the inspector.

(kg)

<section-header><section-header><section-header>

Section E - Quality Testing Procedures

E.1 Responsibilities

Procedure E.1.1 The Quality Testing Operator appointed by the Trials Organiser are responsible for conducting approved quality tests according to these procedures.

E.2 Quality Assessment Methodology for Obligatory and Additional Tests

E.2.1 The Tarehouse Laboratory operated by the Quality Testing Operator shareholde the automated analysis of the brei samples for sugar and impurity measurement Sodium, Potassium and Amino-nitrogen).

E.2.2 Operation of the Tarehouse Laboratory shall be in accordance with the British Sugar Procedure for Tarehouse Laboratory Operation (PR-BP-060). 51

E.2.3 In this process the brei shall be analysed by the 'cold water digestion method'. This is achieved by the proportional addition of dilute basic lead acetate solution to a given weight of brei, followed by digestion and filtration. Chemical analysis shall then take place on the clarified solution.

E.2.4 The above shall be accomplished using a dutomated Laboratory System, comprising of

- Manipulator / Brei Transporter (STD-BP-061 App A)
 - Brei Sampler unit (STD-BP-061 App B)
 - Dispenser / Clarifie
- (STD-BP-061 App C)
- Sample track / Filtration unit (STD-BP-061 App D)
- An Impurity Meter
- (STD-BP-061 App E)
- A Polarimeter
- (STD-BP-061 App F)

E.2.5 Once data for a trachas been analysed it should be sent by the approved methods to the Trial Design and Data Handling Operator.

E.2.6 These characters are also required for DUS: sugar and impurity measurement. this document

Section F - Trial Design and Data Handling **Procedures**

F.1.3 The Trial Design and Data Handling Operator will check these for statistical conditions and once this has been done, will load the plan on the database.
F.2.1 Data are recorded using the methods and characters given in sections of the sections of

F.3 Other Tests and Trials

F.3.1 Any additional or alternative designs required for the assessment of additional VCU characters not detailed in Appendix 3 of the TRIAL PROTOCOL for sugar beet will be added to these **Procedures** as and when approved by the NLSC.

this document is no longer in use

Appendix 1 Approved Trial Organisers/ Operators for Sugar Beet

| rganisers/Operators Responsible |
|--|
| AB ermains AB Project Board Inspectors IAB BRO ritish Sugar AB Control of the second |
| AB ermains AB Project Board Inspectors IAB BRO ritish Sugar AB CONTRACTOR CONTRACTOR AB |
| AB Project Board Inspectors AB BRO ritish Sugar AB Cover the second secon |
| AB Project Board Inspectors IAB BRO ritish Sugar AB Conversion of the second sec |
| Project Board Inspectors AB BRO ritish Sugar AB Cover the second se |
| AB BRO ritish Sugar AB Covint for the covint for the covint for the covint for th |
| BRO ritish Sugar AB COVINTO |
| AB AB COVINTO |
| AB |
| 2 GOVINK FOR THE |
| |
| |

Appendix 2 Seed Treatment Products for Use on NL Trials – to be advised

The following procedure is carried out to pellet and treat sugar beet seed for trials

Seed shall be treated with the following PPPs at the registered rates applicable and in accordance with the manufacturer's label recommendation.

- a) Tachigaren 70 WP To be applied as a surface coating to the pellet at 20 g product per unit / 100,000 seeds

 hymexazol total 20g commercial product = 14g ai/unit · EPAC
 The hymexazol used shall be MAPP not 1775

 b) Force ST To be applied as a surface coating to the pellet at 20 g product / (100,000) section
- - i. 200g tefluthrin per litre = 10g ai/unit; Group 3A IRAC
 - ii. The Force SB used shall be MAPP no: 11752
- c) Vibrance SB To be applied as a surface coating to the vellet at 33.3ml product per unit / 100,000 seeds.
 - i. Sedaxane @ 15g per litre = 0.5 g ai/unit; Group C2 FRAC code 7
 - ii. Fludioxonil @ 22.5g per litre = 0.75g a/upit; Group E2 FRAC code 12
 - iii. Metalaxyl-M @14.4g per litre = 0.48g ji/unit; Group A1 FRAC code 4

in: 185 in: 185 in user The Vibrance SB used shall be MAPP no: 1850

this document is no longer muse. See GOV. W. for the latest procedure **Appendix 3 Seed Delivery Deadline Dates**

Appendix 4 VCU Growing Trial Operators and Trial Locations for Sugar Beet

Growing Trial Operators/Seed Handling Operators

| Growing Trial Operator | Seed Handling Operator (If not Trial Operator) | Location of Trial |
|------------------------|---|--|
| NIAB | | Morley |
| BBRO | NIAB | Bracebridge |
| BBRO | NIAB | Bracebridge Fotheringhay Thelveton |
| BBRO | NIAB | Thelveton |
| BBRO | NIAB | Rougnam |
| SESVANDERHAVE | NIAB | Caythorpe |
| SESVANDERHAVE | NIAB | Holbeach |
| SESVANDERHAVE | NIAB | Walsham le Willows |
| SESVANDERHAVE | NIAB | |
| SESVANDERHAVE | NIAB | Brockford |
| (WS | NIAB | Swaffham Prior |
| KWS KWS | NIAB | Hitcham Fulbourn |
| | USE. SEE | |
| ocumentisnolomoe | NIAB NIAB See | |

this document is no longer in use. see GOV. UN for the latest procedure **Appendix 5 Control Varieties for VCU**

<section-header><section-header><text>

Appendix 7 Dates by which Records Should be Sent to the Trial Design and Data Handling Organiser

| | Latest date of receipt |
|--|--|
| Site data part 1 (including site sketch) | Within 5 days of drilling trial |
| Site data part 2 plus diary | Within 5 days of harvesting the trial |
| Harvest date | Within 2 days of harvest |
| Confirmation of Trial Layout | Within 5 days of drilling the trial |
| Plot records | Within 2 days of harvest Within 5 days of drilling the trial Within 5 days of record being taken Within 5 days of harvesting the trial |
| Harvest records | Within 5 days of harvesting the trial |
| oumentienolonoerimue | Etest date of receipt |

Appendix 8 Assessment Keys for Sugar Beet Diseases

Leaf diseases

- 1.
- 2.
- 3.
- 4.

Infection Disease Severity Description

- 0
- 0.1
- 1
- 5
- Include all necrosis and chlorosis attributable to disease to be assessed Estimate % infection using the description below, interpolating values if necessary of the average % infection from the 3 areas
 Ction Disease Severity Description
 No infection observed
 Older leaves with a trace of infection, other leaves uninfected.
 Older leaves with up to 10% infection, other leaves largely prinfected.
 Older leaves with up to 25% infection, middle aged to 0 other leaves with up to 25% infection, middle aged to 0 other leaves with up to 25% infection. 10 uninfected.
- Leaves of all ages appear 50% infected 50% green on average 25
- this document is no longer in use. See Leaves of all ages appear more infected than green on average

Appendix 9 Growth Stage Key of Beet

Beet Meier et al., 1993

For the latest procedure Phenological growth stages and BBCH-identification keys of beet (Beta vulgaris L. ssp. vulgaris)

Code Description

Principal growth stage 0: Germination

00 Dry seed 01 Beginning of imbibition: seeds begins to take up water 03 Seed imbibition complete (pellet cracked)

- 05 Radicle emerged from seed (pellet)
- 07 Shoot emerged from seed (pellet)
- 09 Emergence: shoot emerges through soil surface

Principal growth stage 1: Leaf development (youth stage

- 10 First leaf visible (pinhead-size): cotyledons horizontally boblided
- 11 First pair of leaves visible, not yet unfolded (pea-size)
- 12 2 leaves (first pair of leaves) unfolded
- 14 4 leaves (2nd pair of leaves) unfolded
- 15 5 leaves unfolded
- Stages continuous till . . .
- 19 9 and more leaves unfolded

See Principal growth stage 3: Rosette growth (crop cover)

31 Beginning of crop cover: leaves over 10% of ground

32 Leaves cover 20% of ground

33 Leaves cover 30% of ground

34 Leaves cover 40% of ground

35 Leaves cover 50% of a synd

36 Leaves cover 60% opround

37 Leaves cover 70% of ground 38 Leaves cover 80% of ground

39 Crop cover complete: leaves cover 90% of ground

49 Beet root has reached harvestable size Principal growth stage 4: Development of harvestable vegetative plant parts Beet

Appendix 10 - Control of Groundkeepers from ALS tolerant plots in subsequent years.

vvnilst good gleaning at harvest will minimize the risk of groundkeepers in the following year the following control measures should be adopted. Measure 3 should be regarded to obligatory.
1. Ploughing post-harvest. This will bury and this been present. Your head

- Using a non-ALS herbicide. In the following crop a non-ALS herbicide that will kill 2. any beet present including ALS-tolerant beets. Your host growershould be made aware of this.
- Inspecting the trial field. This must be done in the following season for the presence 3. of groundkeepers. If any ALS-tolerant groundkeepers are present they must be removed, normally by hand. The location of the ALS plots should be recorded and mean set in mean set in mean set in mean set in mean set is no longer in use. See these specific areas can be targeted for inspection. A diary entry should be made of inspection date, findings and control measures taken.



© Crown copyright 2018

misdocumentis no longer in use. See