

Routine childhood immunisations from September 2012

When to immunise	Diseases protected against		Vaccine given	Immunisation site**
Two months old	Diphtheria, tetanus, pertussis, polio and <i>Haemophilus</i> <i>influenzae</i> type b (Hib)		DTaP/IPV/Hib (Pediacel)	Thigh
	Pneumococcal disease		PCV (Prevenar 13)	Thigh
Three months old	Diphtheria, tetanus, pertussis, polio and Hib		DTaP/IPV/Hib (Pediacel)	Thigh
	Meningococcal group C disease (MenC)		MenC (Menjugate or Neisvac C)	Thigh
Four months old	Diphtheria, tetanus, pertussis, polio and Hib		DTaP/IPV/Hib (Pediacel)	Thigh
	MenC		MenC (Menjugate or Neisvac C)	Thigh
	Pneumococcal disease		PCV (Prevenar 13)	Thigh
Between 12 and 13 months old – within a month of the first birthday	Hib/MenC		Hib/MenC (Menitorix)	Upper arm/thigh
	Pneumococcal disease		PCV (Prevenar 13	Upper arm/thigh
	Measles, mumps and rubella (German measles)		MMR (Priorix or view Valence)	Upper arm/thigh
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio		dTaP/I/V v epevax or DTaP/IPV (Infan ix-IPV)	Upper arm
	Measles, mumps and rubella		wine 8 (Penrix or MMR VaxPRO) (check first dose has been given)	Upper arm
Girls aged 12 to 13 years old	Cervical cancer caused by human papillomavirus types 16 and 18 (and genital warts caused by types 6 and 11)	~	Garaasil	Upper arm
Thirteen to 18 years old	Tetanus, diphtheria and polio		Td/IPV (Revaxis), and check MMR status	Upper arm

** Where two or more injections are required at once, these she did ideally be given in different limbs. Where this is not possible, injections in the same limb should be given 2.5cm apart. For more details see Chapters 4 and 11 in the Green Book.

Non-routine immunisations for at the babies

At birth, 1 month old, 2 months old and 12 months old	Нерать В	Нер В		Thigh
At birth	Interculosis	BCG		Upper arm (intradermal)
	Prevenar 13	Menjugate*	Neisvac C*	Menitorix
Priorix* MMR	VaxPRO* Repe	vax* Infanrix-IF	V* Gardasil	Revaxis

*NB Where a vaccine is manufactured by more than one supplier, it may, on occasion be necessary to substitute an alternative brand. All vaccines for use in the routine childhood programme can be ordered free of charge at www.immform.dh.gov.uk



