



This report summarises the information from the surveillance systems which are used to monitor the Coronavirus Disease 2019 (COVID-19) pandemic in England. More information on the surveillance systems are available [here](#).

The report is based on data from week 37 (between 7 August and 13 September 2020) and for some indicators daily data up to 15 September 2020. References to COVID-19 represent the disease name and SARS-CoV-2 represent the virus name.

Data is reported from week 27 (week beginning 29 June 2020) onwards. For reports with data prior to week 27, consult previous reports [here](#).

Summary

Several surveillance indicators suggest that COVID-19 activity at a national level has continued to increase during week 37.

Case detections in England increased from 15,231 in week 36 to 15,693 in week 37. Case rates remain highest in North West and Yorkshire and Humber. Further increases in detections for week 37 are expected as more results for the most recent samples become available. Positivity rates have increased across most age groups particularly in the 85+ year olds tested through Pillar 2 and those in the 15-44 years in Pillar 1. Positivity by regions remains highest in the North. At a local authority level, incidence continued to be highest in Bolton. Case rates continued to be highest in the 20-29 year age group.

The following local authorities have been included in the watchlist following the weekly Local Action Committee meeting as areas of intervention: Bolton, Blackburn with Darwen, Oadby and Wigston, Hyndburn, Preston, Warrington, Tameside, Sunderland, Oldham, Birmingham, Bradford, Liverpool, Wirral, Burnley, Knowsley, St. Helens, Bury, Salford, Leicester, South Tyneside, Rochdale, Manchester, Gateshead, Solihull, Sandwell, Newcastle upon Tyne, Pendle, Halton, Kirklees, Wolverhampton, Calderdale, Rossendale, South Ribble, Sefton, North Tyneside, West Lancashire, County Durham, Trafford, Chorley, Wyre, Fylde, Northumberland, Lancaster and Ribble Valley.

The overall number of acute respiratory infection incidents reported to PHE Health Protection Teams increased notably from 246 in the previous week to 729 in week 37. The highest increases were noted in the number of incidents in care homes, educational and workplace settings in comparison to the previous week. Around half of the incidents in educational settings were confirmed as COVID-19 outbreaks. There has also been an increase in rhinovirus activity in school aged children which may account for some of the acute respiratory infection incidents reported.

Community and syndromic indicators continued to increase during week 37. This included large increases in NHS 111 cold/flu calls and GP out of hours consultations for influenza like illness.

Through the GP swabbing scheme, an increase in positivity was noted in week 37 at 7.6% compared to 1.4% in the previous week.

There were small increases in emergency department attendances with a COVID-19-like diagnosis. Increases continued to be seen in hospital admission rates for confirmed COVID-19 at national level, particularly in the 85+ year olds by age group and in the North West and Midlands by region. ICU/HDU admission rates increased slightly with highest rates seen in the 45-64 year olds.

COVID-19 deaths were stable in week 37 and no excess mortality was observed overall in week 36.

New adjusted seroprevalence estimates based on samples from adult blood donors in the North East and Yorkshire was relatively stable at 3.9%. In the Midlands seroprevalence was estimated at 6.8%, an increase on the previous sampling period which may reflect a higher proportion of samples from Birmingham. Among adult blood donors, seroprevalence is highest in the youngest age groups.

Following this week's meeting of the Local Action Committee, the Secretary of State for Health and Social Care, drawing on epidemiological advice from the CMO, NHS Test and Trace, JBC and PHE, has determined the following Watchlist (Table 1 and 2), highlighting the local authorities of greatest concern.

The Watchlist is produced by first considering the lower tier local authorities with the highest weekly incidence rate and its trend, combined with a range of other indicators including the test positivity rate, an assessment of the local response and plans, and the trend of other metrics such as healthcare activity and mortality. The classification decision is therefore a blended assessment drawing on professional judgement.

Whilst this list is determined at the granularity of lower tier local authority, the Contain Framework places responsibility for local action at the level of the upper tier local authority. Later in this report, we list the UTLA with the highest incidence rate in the country from a purely statistical viewpoint (Figure 11).

The Watchlist classification uses definitions as set out in the Contain Framework:

- area(s) of concern—for areas with the highest incidence, where the local area is taking targeted actions to reduce prevalence e.g. additional testing in care homes and increased community engagement with high risk groups
- area(s) for enhanced support—for areas at medium/high risk of intervention where there is a more detailed plan, agreed with the national team and with additional resources being provided to support the local team (e.g. epidemiological expertise, additional mobile testing capacity)
- area(s) of intervention—where there is divergence from the measures in place in the rest of England because of the significance of the spread, with a detailed action plan in place, and local resources augmented with a national support

Maps representing the areas from this week's Watchlist (Table 1 and 2) by Lower Layer Super Output Area (LSOA) are available [here](#).



Table 1: Local Authority watchlist areas - Areas of intervention

Lower Tier Local Authority	Individuals tested per day per 100,000 population (7 day moving average)	Trend	Incidence per 100,000 population (weekly)	Trend	Contain Framework Watchlist Status - week beginning 4 September	Change in Watchlist Status from previous week	Area with household mixing prohibited?
Bolton	257.4	↑	212.7	↑	Intervention	→	YES
Blackburn with Darwen	266.6	↑	122.9	↑	Intervention	→	YES
Oldby and Wigston	243.4	↑	119.2	↑	Intervention	↑	YES
Hyndburn	299.8	↑	117.6	↑	Intervention	↑	YES
Preston	285.4	↑	105.1	↑	Intervention	→	YES
Warrington	217.8	↑	105.0	↑	Intervention	↑	YES
Tameside	203.7	↑	103.5	↑	Intervention	→	YES
Sunderland	324.7	↑	103.1	↑	Intervention	↑	YES
Oldham	236.2	↑	98.9	↑	Intervention	→	YES
Birmingham	164.8	↑	98.0	↑	Intervention	→	YES
Bradford	171.5	↑	97.5	↑	Intervention	→	YES
Liverpool	182.7	↑	95.8	↑	Intervention	↑	YES
Wirral	270.0	↑	95.6	↑	Intervention	↑	YES
Burnley	315.8	↑	93.8	↑	Intervention	→	YES
Knowsley	210.4	↑	92.9	↑	Intervention	↑	YES
St. Helens	176.5	↑	91.6	↑	Intervention	↑	YES
Bury	199.3	↑	90.5	↑	Intervention	→	YES
Salford	170.4	↑	88.8	↑	Intervention	→	YES
Leicester	216.1	↑	86.7	↑	Intervention	→	YES
South Tyneside	169.6	↓	86.5	↑	Intervention	↑	YES
Rochdale	223.4	↑	84.1	↑	Intervention	→	YES
Manchester	167.6	↑	83.6	↑	Intervention	→	YES
Gateshead	174.8	↑	77.5	↑	Intervention	↑	YES
Solihull	181.3	↑	77.2	↑	Intervention	→	YES
Sandwell	164.2	↑	72.1	↑	Intervention	→	YES
Newcastle upon Tyne	169.7	↑	69.6	↑	Intervention	↑	YES
Pendle	245.5	↑	61.3	↑	Intervention	→	YES
Halton	183.5	↑	60.7	↑	Intervention	↑	YES
Kirklees	135.0	↑	60.4	↑	Intervention	→	YES
Wolverhampton	204.6	↑	60.3	↑	Intervention	↑	YES
Calderdale	151.3	↑	59.5	↑	Intervention	→	YES
Rossendale	290.2	↑	57.8	↓	Intervention	↑	YES
South Ribble	197.4	↑	52.5	↑	Intervention	↑	YES
Sefton	169.0	↑	49.0	↑	Intervention	↑	YES
North Tyneside	176.9	↑	48.5	↑	Intervention	↑	YES
West Lancashire	149.3	↑	47.4	↑	Intervention	↑	YES
County Durham	196.0	↑	46.7	↑	Intervention	↑	YES
Trafford	159.3	↑	45.7	↑	Intervention	↑	YES
Chorley	167.5	↑	35.1	↑	Intervention	↑	YES
Wyre	128.1	↑	34.2	↑	Intervention	↑	YES
Fylde	134.3	↑	28.8	↑	Intervention	↑	YES
Northumberland	164.2	↑	24.7	↑	Intervention	↑	YES
Lancaster	118.8	↑	22.9	↑	Intervention	↑	YES
Ribble Valley	165.3	↑	18.3	↓	Intervention	↑	YES
ENGLAND	149.8	↑	33.8	↑			



Table 2: Local Authority watchlist areas - Areas of enhanced support and concern

Lower Tier Local Authority	Individuals tested per day per 100,000 population (7 day moving average)	Trend	Incidence per 100,000 population (weekly)	Trend	Contain Framework Watchlist Status – week beginning 4 September	Change in Watchlist Status from previous week	Area with household mixing prohibited?
Leeds	166.4	↑	75.5	↑	Enhanced Support	→	NO
Blaby	251.2	↑	65.7	↑	Enhanced Support	↑	NO
Stockport	155.4	↑	48.7	↑	Enhanced Support	→	NO
Selby	186.8	↑	65.1	↑	Concern	↑	NO
Hartlepool	199.8	↑	55.8	↑	Concern	→	NO
Sheffield	161.7	↑	53.7	↑	Concern	→	NO
Spelthorne	157.8	↑	53.4	↑	Concern	↑	NO
Corby	273.5	↑	50.8	↑	Concern	→	NO
Middlesbrough	250.6	↑	47.0	↑	Concern	→	NO
Northampton †	211.7	↑	42.6	↑	Concern	→	NO
Scarborough	141.0	↑	42.3	↑	Concern	↑	NO
Hertsmere	174.4	↑	37.4	↓	Concern	→	NO
Peterborough	119.8	↑	30.3	↑	Concern	→	NO
Stoke-on-Trent	153.9	↑	27.4	↑	Concern	→	NO
ENGLAND	149.8	↑	33.8	↑			

Data for specimens taken between 4 September and 10 September as extracted on 15 September

Trend arrow indicates whether there has been an increase, decrease or no change between this week and last week (specimens taken between 28 August and 3 September)

Some Local Authority areas have been included as part of wider geographical interventions.

† Northampton's increase in incidence is almost solely down relates to a workplace outbreak at the Greencore Factory

‡ Within these Local Authority the interventions have been restricted to some wards

§ Northampton's increase in incidence is almost solely down relates to a workplace outbreak at the Greencore Factory


¶ These Local Authorities are within Norfolk and relate almost solely to a workplace outbreak at Banham Poultry Farm.



As of 09:00 on 15 September 2020, a total of 323,029 have been confirmed positive for COVID-19 in England under Pillar 1 and 2.

Overall case numbers and positivity increased in both Pillar 1 and 2, in week 36, with the majority of cases reported from Pillar 2. The highest case rates continued to be seen in the 20-29 year olds. Positivity was highest in 15-44 year olds in Pillar 1 and in 85+ year olds in Pillar 2. Cases rates and positivity continue to be highest in the North West of England.


Figure 1: Laboratory confirmed COVID-19 cases tested under Pillar 1 and Pillar 2, based on sample week with overall positivity for Pillar 1 and 2 (%)



* For the most recent week, more samples are expected therefore the decrease seen in this graph should be interpreted with caution. The data are shown by the week the specimen was taken from the person being tested. This gives the most accurate analysis of this time progression, but it does mean that the latest days' figures may be incomplete.

Age and sex

Figure 2: Age/sex pyramids for laboratory confirmed COVID-19 cases tested under Pillar 1 and 2 (a) cumulative number since week 27 (n=79,301), and (b) in weeks 36 and 37 (n=30,765)



Age and sex

Figure 3: Weekly laboratory confirmed COVID-19 case rates per 100,000, tested under Pillar 1 and Pillar 2, by sex




Figure 4: Weekly laboratory confirmed COVID-19 case rates per 100,000, tested under Pillar 1 and Pillar 2, by age group




Figure 5: Weekly positivity (%) of laboratory confirmed COVID-19 cases tested overall and by sex under (a) Pillar 1 and (b) Pillar 2, (SGSS and Respiratory DataMart)





Figure 6: Weekly positivity (%) of laboratory confirmed COVID-19 cases tested under Pillar 1, (a) by male and age group and (b) by female and age group and; under Pillar 2, (c) by male and age group and (d) by female and age group, (SGSS and Respiratory DataMart)



Geography

Table 3: Cumulative number of cases under Pillar 1 and 2 (n=313,707) and cumulative number of cases since week 27 under Pillar 1 and 2 (n=78, 770) and total number of people tested under Pillar 1 and 2 (n= 7,439,412) by PHE Centres

PHE Centres	Cumulative Pillar 1 + 2 cases	Cumulative since week 27, Pillar 1 + 2 cases	Total number of people tested (under Pillar 1 + 2)
North East	18,685	3,641	342,825
North West	62,086	19,869	1,114,503
Yorkshire & Humber	40,746	12,047	762,726
West Midlands	34,893	9,756	726,326
East Midlands	29,008	8,361	693,332
East of England	29,493	5,370	835,790
London	43,685	10,028	1,069,749
South East	39,164	6,423	1,172,699
South West	15,947	3,275	721,462

Figure 7: Weekly laboratory confirmed COVID-19 case rates per 100,000 population tested under Pillar 1 and Pillar 2, by PHE Centres and sample week




Figure 8: Weekly positivity of laboratory confirmed COVID-19 cases tested under (a) Pillar 1 (%) and (b) Pillar 2 (%), by PHE Centres and sample week, (SGSS and Respiratory DataMart)





Figure 9: Cumulative rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2, by upper-tier local authority, England (box shows enlarged map of London area)




Figure 10: Cumulative rate (from week 27) of COVID-19 cases per 100,000 population tested under Pillar 1 and 2, by upper-tier local authority, England (box shows enlarged map of London area)




Figure 11: Weekly rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2, by upper-tier local authority, England (box shows enlarged map of London area)






Figure 12: UTLA with the highest weekly rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2*



*The UTLA data presented in this figure, is based on data extracted on Tuesday 15 September, covering the period of 7 August to 13 September 2020 (week 37).

Ethnicity

Figure 13: Weekly incidence per 100,000 population by ethnicity, England



Incidence rates by region

In the regions with the highest overall rates and with most local authorities on the watchlist, the age groups most affected appears to be young working age adults (20-29 years). This is consistent with mixing patterns in this age group who may be more likely to be working away from home, including in public facing roles. In those regions, highest rates are also observed in Asian communities of either Other ethnic background or Pakistani origin, most likely reflecting the ethnic mix in the most affected local areas. In some regions the daily numbers of cases in each ethnic group can be small, so minor variations in rates should be interpreted with caution.

Figure 14: Weekly incidence per 100,000 population by age group and region, weeks 31-37






Figure 15: Weekly incidence per 100,000 population by ethnicity and region, weeks 31-37



*Data presented in Figures 14 & 15 are calculated using Government Office Region denominators

Case rates by Index of Multiple Deprivation (IMD)

Figure 16: Weekly case rate per 100,000 population by IMD quintile (1 being the most deprived and 5 being the least deprived, weeks 27-37



Other respiratory viruses, Datamart

The [Respiratory Datamart system](#) was initiated during the 2009 influenza pandemic to collate all laboratory testing information in England. It is now used as a laboratory surveillance tool, monitoring all major respiratory viruses in England.

Figure 17 and 18 represent weekly positivity of other respiratory viruses in particular rhinovirus. In week 37, the positivity for rhinovirus increased to 22.7% compared to 10.5 in the previous week (Figure 17). The highest positivity was seen in the 5-14 year olds (Figure 18).

Figure 17: Weekly positivity for other respiratory viruses reported through Respiratory Datamart, England





Figure 18: Weekly positivity for rhinovirus by age group, reported through Datamart, England



This section summarises the monitoring of acute respiratory infection incidents and internet based surveillance systems for COVID-19.

Acute respiratory infection incidents, England

Information on acute respiratory infection (ARI) incidents is based on situations reported to PHE Health Protection Teams (HPTs). These include:

- confirmed outbreaks of acute respiratory infections ie two or more laboratory confirmed cases (COVID-19, influenza or other respiratory pathogen) linked to a particular setting
- situations where an outbreak is suspected. All suspected outbreaks are further investigated by the HPT in liaison with local partners and a significant proportion do not meet the criteria of a confirmed outbreak. For example if suspected cases test negative for COVID-19 or other respiratory pathogens, or cases are subsequently found not to have direct links to the setting. Since Pillar 2 testing became open to everyone during week 21 more incidents of mild disease have been detected in settings with healthy young populations.

Processes for reporting ARI incidents vary between PHE Centres.

The number of incidents in each setting with at least one laboratory confirmed case of COVID-19 are reported below.

729 new ARI incidents have been reported in week 37 (Figure 19):

- 313 incidents were from care homes where 228 had at least one linked case that tested positive for SARS-CoV-2
- 18 incidents were from hospitals where 13 had at least one linked case that tested positive for SARS-CoV-2
- 193 incidents were from educational settings where 110 had at least one linked case that tested positive for SARS-CoV-2
- 4 incidents were from prisons where 3 had at least one linked case that tested positive for SARS-CoV-2
- 110 incidents were from workplace settings where 92 had at least one linked case that tested positive for SARS-CoV-2
- 34 incidents were from food outlet/restaurant settings where 25 had at least one linked case that tested positive for SARS-CoV-2
- 57 incidents were from the other settings category where 39 had at least one linked case that tested positive for SARS-CoV-2

Acute respiratory infection incidents, England

Figure 19: Number of acute respiratory infection (ARI) incidents by institution, England




Table 4: Total number of situations/incidents by institution and PHE Centres over the past four weeks with the total number in the last week in brackets

PHE Centres	Cumulative total number of incidents by institution over the past 4 weeks with total number in the last week in brackets							
	Care home	Hospital	Educational settings	Prisons	Workplace settings	Food outlet/restaurant settings	Other settings	Total
East of England	44(23)	1(0)	13(11)	0(0)	9(4)	2(1)	10(1)	79(40)
East Midlands	57(40)	2(1)	21(19)	0(0)	22(6)	8(4)	10(6)	120(76)
London	34(14)	9(6)	25(23)	1(0)	28(12)	8(1)	18(6)	123(62)
North East	22(14)	3(0)	8(7)	1(0)	17(9)	15(4)	16(5)	82(39)
North West	87(42)	7(2)	40(29)	2(2)	56(22)	37(7)	32(3)	261(107)
South East	74(52)	4(3)	17(11)	0(0)	14(9)	13(6)	14(8)	136(89)
South West	66(41)	0(0)	41(36)	0(0)	24(12)	4(3)	16(3)	151(95)
West Midlands	70(41)	12(5)	41(38)	1(0)	36(17)	11(6)	26(11)	197(118)
Yorkshire and Humber	80(46)	1(1)	24(19)	2(2)	46(19)	6(2)	29(14)	188(103)
Total	534(313)	39(18)	230(193)	7(4)	252(110)	104(34)	171(57)	1337(729)

Confirmed COVID-19 clusters or outbreaks in educational settings , England

ARI incidents in educational settings are reviewed to identify confirmed COVID-19 clusters or outbreaks as per the definitions below. This does not include incidents with only one confirmed case identified, incidents where COVID-19 is suspected but confirmation is awaited, incidents where other causative organisms have been identified or incidents where there was no causative organisms identified.


A cluster is defined as two or more test-confirmed cases of COVID-19 among individuals associated with a specific non-residential setting with illness onset dates within a 14-day period (in the absence of detailed information about the type of contact between the cases).

An outbreak is defined as two or more test-confirmed cases of COVID-19 among individuals associated with a specific non-residential setting with illness onset dates within 14 days, and one of:

- (1) Identified direct exposure between at least 2 of the test-confirmed cases in that setting (for example under one metre face to face, or spending more than 15 minutes within 2 metres) during the infectious period of one of the cases
- (2) When there is no sustained local community transmission - absence of an alternative source of infection outside the setting for the initially identified cases

In week 37, there were 108 confirmed COVID-19 clusters or outbreaks in educational settings. The highest number of COVID-19 confirmed clusters or outbreaks were reported through secondary schools (Figure 20).

Figure 20: Number of COVID-19 confirmed clusters or outbreaks by type of educational setting, England



Confirmed COVID-19 clusters or outbreaks in educational settings , England

Table 5: Cumulative number of confirmed COVID-19 clusters or outbreaks by type of educational setting and PHE Centres since week 36, England

PHE Centres	Cumulative number of confirmed COVID-19 clusters or outbreaks by type of educational setting					
	Nursery	Primary school	Secondary school	Special Educational Needs (SEN) schools	College/University	Total
East of England	1	4	4	1	0	10
East Midlands	0	7	2	1	4	14
London	1	4	5	0	2	12
North East	0	1	4	1	0	6
North West	1	7	16	2	3	31
South East	2	2	3	1	0	6
South West	0	0	3	0	1	4
West Midlands	0	13	12	0	0	25
Yorkshire and Humber	0	1	6	3	4	14
Total	5	39	55	9	14	122

Cases by type of residence

Table 4 shows the proportion of confirmed COVID-19 cases according to their type of residence. Property classifications are derived from Ordnance Survey AddressBase and are matched to address details within the laboratory data. Properties are identified by unique property reference number (UPRN) and basic land property unit (BLPU). Cases with poor or no address data which failed the address matching and are classed as 'undetermined'. No fixed abode and overseas addresses identified by recording in the laboratory data.

In week 37, there were small increases in the percentage of cases in residential dwelling (Table 6).

Table 6: Type of residence of confirmed COVID-19 cases by percentage of total weekly cases

Type of residence	week 27	week 28	week 29	week 30	week 31	week 32	week 33	week 34	week 35	week 36	week 37
Residential dwelling (including houses, flats, sheltered accommodation)	74.7	71.6	75.0	73.0	73.6	72.0	71.4	74.1	76.4	77.7	80.4
Undetermined	18.8	20.0	19.3	20.4	19.7	21.0	22.8	21.1	19.0	17.6	15.4
Care/Nursing home	4.9	6.2	4.4	4.5	5.6	5.8	4.2	2.8	2.5	3.1	2.6
Residential institution (including residential education)	0.3	0.2	0.1	0.4	0.1	0.1	0.3	0.4	0.4	0.4	0.5
House in multiple occupancy (HMO)	0.4	0.3	0.3	0.6	0.4	0.4	0.4	0.7	0.7	0.4	0.4
Medical facilities (including hospitals and hospices, and mental health)	0.6	1.4	0.8	0.9	0.3	0.2	0.5	0.5	0.4	0.3	0.4
Other property classifications	0.2	0.2	0.1	0.2	0.2	0.4	0.4	0.4	0.5	0.5	0.3
Prisons, detention centres, secure units	0.2	0.1	0.0	0.1	0.1	0.2	0.1	0.0	0.1	0.0	0.0
Overseas address	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
No fixed abode	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0


Contact tracing

Once a person has a confirmed positive test result for coronavirus, this person is transferred to NHS Test and Trace and a case is opened for them. The NHS Test and Trace service will get in contact via a text, email alert or phone call. People are asked to share details of other people with whom they have had close, recent contact and places they have visited. They can respond online via a secure website or by telephone with a contract tracer. Once contacts have been identified, they will be contacted in turn by the NHS Test and Trace service and advised to self-isolate.

Contacts in Figure 21 are those named by people testing positive and contact traced by NHS Test and Trace. The setting is the potential exposure setting as reported by the person who tested positive, when they had close interaction with the named contact. The most common setting was the household, where 57.2% of all contacts were identified. The next most common setting was visitors to the household of the person who tested positive (13.5%).

The number of contacts excludes those identified as part of management of complex cases: such as those investigated as part of an outbreak, for example, if someone works in or has recently visited a health or care setting such as a hospital or care home, a prison or other secure setting, or a school for people with special needs. For complex cases, contacts are often managed at a situation rather than individual level, with advice being issued to the contact institution (for example in a care home or prison). Therefore information on individual contacts associated with these situations is not available.

Figure 21: Contacts by exposure/activity setting in week 37, England
(Data source: NHS Test and Trace)



Note: categories have been grouped as follows: leisure / community includes eating out, attending events and celebrations, exercising, worship, arts, entertainment or recreation, community activities and attending play groups or organised trips; other workplace includes: retail, manufacturing or construction, hospitality, transport, emergency services or border force, food production and agriculture, prison, financial services, civil service or local government, information and communication, military, critical national infrastructure. Personal services includes hairdressers, barbers, tattooists and nail bars.

NHS 111

The NHS 111 service monitors daily trends in phone calls made to the service in England, to capture trends in infectious diseases such as influenza and norovirus.

Up to 13 September 2020, the daily percentage of NHS 111 ‘potential COVID-19-like’ calls (as a percentage of total NHS 111 calls) and number of online assessments increased. The daily percentage of cold/flu calls (as a percentage of total NHS 111 calls) continued to increase as did cold/flu completed online assessments (Figure 22 and 23).

Please note that NHS 111 callers (from 11 May 2020) and NHS 111 online users (from 11 June 2020), who are assessed as having probable COVID-19 symptoms are now triaged using symptom specific pathways e.g. cold/flu, which are included in routine syndromic indicators.

Further information about these caveats is available from the PHE Remote Health Advice Syndromic Surveillance bulletin.

Figure 22 (a-b): NHS 111 telephony indicators (and 7-day moving average), England

(a) Daily potential COVID-19 calls as a percentage of total calls, all ages

(b) Daily cold/flu calls as a percentage of total calls, all ages







Figure 23 (a-b): NHS 111 completed online assessments (and 7-day moving average), England

(a) Daily ‘potential COVID-19’ online assessments as the number of completed online assessments, all ages

(b) Daily cold/flu online assessments as the number of completed online assessments, all ages



Legend: weekend (light grey), bank holiday (dark grey), indicator (teal line), 7 day mov avg (black line), baseline (dotted black line)

Internet based surveillance

PHE's internet based surveillance systems aim to monitor the volume of people searching for typical symptoms of COVID-19 on the internet as well as tracking self-reported respiratory symptoms and health seeking behaviour patterns related to COVID-19.


Google search queries

This is a web-based syndromic surveillance system which uses daily search query frequency statistics obtained from the Google Health Trends API [1]. This model focuses on search queries about COVID-19 symptoms as well as generic queries about "coronavirus" (eg "covid-19"). The search query frequency time series has been weighted based on symptom frequency as reported in other data sources. Frequency of searches for symptoms is compared with a baseline calculated from historical daily data.

The overall and media-debiasing weighted scores increased during week 37 (Figure 24).

[1] For more information about this model, please see <https://arxiv.org/abs/2003.08086>

Figure 24: Normalised Google search score for COVID-19 symptoms, with weighted score for media-debiasing and historical trend, England




Internet based surveillance

FluSurvey

An internet based surveillance system has been developed based on FluSurvey. FluSurvey is a web tool survey designed to monitor trends of influenza like illness (ILI) in the community using self-reported respiratory symptoms from registered participants. The platform has been adapted to capture respiratory symptoms, exposure risk and healthcare seeking behaviours among registered participants to contribute to national surveillance of COVID-19 activity.

A total of 3,586 participants completed the weekly COVID-19 surveillance survey in week 37, of which 123 (3.4%) reported fever or cough, a slight increase from the previous week. The most commonly reported method of access to healthcare services continue to be through telephoning a GP practice in week 37 (Figure 25).

Figure 25: Rate of contact with different healthcare services among FluSurvey participants reporting fever or cough symptoms, week 27 to 37, England



GP In Hours (GPIH) and GP Out of Hours (GPOOH), Syndromic surveillance

The GP In Hours (GPIH) syndromic surveillance system monitors the number of GP visits during regular hours of known clinical indicators. The GP Out of Hours (GPOOH) syndromic surveillance system monitors the numbers of daily unscheduled visits and calls to GPs during evenings, overnight, on weekends and on public holidays. Both systems cover around 55% of England’s population.

Up to 13 September 2020, GPIH consultations for potential COVID-19-like and ILI consultations remained stable (Figure 26). Please note that the GPIH COVID-19-like indicator presented in this report is derived from a reduced denominator population, compared to ILI. Please also note, week 36 contains a bank holiday and there were also days with a reduced denominator and therefore these recent rates should be interpreted with some caution.

Rates should therefore be treated with caution (baselines are also not available this week). Through GPOOH consultations (up to 13 September 2020), the daily percentage (as a percentage of total contacts with a Read code) for ILI and difficulty breathing/wheeze/asthma contacts have increased (Figure 27).

Please note GP data should be interpreted with caution due to changes in advice regarding accessing GP surgeries due to COVID-19. Further information about these caveats is available from the PHE GP In Hours Syndromic Surveillance bulletin.

Figure 26 (a-b): GPIH clinical indicators, England

(a) potential COVID-19 GP consultations, daily incidence rates per 100,000 population, all ages

(b) Influenza-like illness consultations, daily incidence rates per 100,000 population, all ages





Figure 27 (a-b) : GPOOH contacts indicators, England

(a) Difficulty breathing/wheeze/asthma, daily contacts (%), all ages

(b) Influenza-like illness, daily contacts (%), all ages




RCGP swabbing scheme

This is an extended primary care surveillance system through the RCGP sentinel integrated clinical and virological scheme. The extension of the scheme was initiated on 24 February 2020. A sample of patients presenting to around 300 GP practices with Influenza-like Illness (ILI) and Lower Respiratory Tract Infections (LRTI) (not suspected for COVID-19) will be tested. This enables the week on week monitoring of test “positivity rate” to observe the trend in the proportion of people with confirmed COVID-19.

Up to 15 September 2020, a total of 5,461 patients have been tested of which 622 have tested positive for SARS-CoV-2 through this scheme. The overall positivity was at 7.6% (6/79) in week 37 compared to 1.4% (1/69) in the previous week (Figure 28). This should be interpreted with caution as the overall denominator for patients tested through GPs has decreased due to an increase in patients being tested under Pillar 2. Consultations for LRTI increased and a slight increase was noted in consultations for ILI in week 37 (Figure 28).

Figure 28: Overall weekly positivity (%), ILI and LRTI consultations rates (per 100,000), RCGP, England



*For the most recent week, more samples are expected to be tested therefore the graph in Figures 23-25 should be interpreted with caution

*Positivity (%) is not calculated when the total number tested is less than 10

Emergency Department attendances, Syndromic surveillance

The Emergency Department Syndromic Surveillance System (EDSSS) monitors the daily visits in a network of emergency departments across England.

Up to 12 September 2020, the daily number of ED attendances for all ages as reported by 71 EDs in England during week 37, for COVID-19-like attendances increased (Figure 29). Increases were noted in acute respiratory infection attendances.

Please note: the COVID-19-like ED indicator is an underestimation of the number of COVID-19 attendances as it only includes attendances with a COVID-19-like diagnosis as their primary diagnosis. The EDSSS COVID-19-like indicator should therefore be used to monitor trends in ED attendances and not to estimate actual numbers of COVID-19 ED attendances. Further information about these caveats is available from the PHE Emergency Department Syndromic Surveillance bulletin.

Figure 29: COVID-19-like, daily ED attendances, all ages, England




COVID-19 Hospitalisation in England Surveillance System (CHES)

The CHES surveillance system monitors daily new acute respiratory infections (ARI) and new laboratory confirmed COVID-19 admissions to hospital including critical care (ICU/HDU). Trends in hospital and critical care admission rates need to be interpreted in the context of testing recommendations.

A total of 134 NHS Trusts are now participating, although the number of Trusts reporting varies by day. The weekly rate of new admissions of COVID-19 cases is based on the trust catchment population of those NHS Trusts who made a new return. This may differ from other published figures such as the total number of people currently in hospital with COVID-19.

In week 37, the weekly admission rate for hospital and ICU/HDU admissions increased. The hospitalisation rate was at 1.42 per 100,000 in week 37 compared to 0.77 per 100,000 in the previous week. The ICU/HDU rate was at 0.14 per 100,000 in week 37 compared to 0.07 per 100,000 in the previous week (Figure 30). By NHS regions, the highest hospitalisation and ICU/HDU rates continued to be observed in the North West (Figure 31). By age group, the highest hospitalisation rate was observed in the 85+ year olds and the highest ICU/HDU rate was observed in the 45-64 year olds (Figure 32).

Figure 30: Weekly overall hospital and ICU/HDU admission rates per 100,000 of new COVID-19 positive cases reported through CHES, England



COVID-19 Hospitalisation in England Surveillance System (CHES)

Figure 31: Weekly admission rate for (a) hospital admissions and (b) ICU/HDU admissions by NHS regions of new COVID-19 positive cases reported through CHES




Figure 32: Weekly admission rate for (a) hospital admissions and (b) ICU/HDU admissions by age group of new COVID-19 positive cases reported through CHES





Figure 33: Weekly admission rates for hospitalised laboratory confirmed COVID-19 cases reported through CHES, week 37




COVID-19 Hospitalisation in England Surveillance System (CHES)


Figure 34 and 35 are based on individual patient level data which are provided to CHES from a subset of NHS Acute Trusts, therefore the data should be interpreted with caution as the distribution of age, sex and ethnic group may not be representative of all hospitalised patients.

Figure 34: Age/sex pyramid of new (a) hospital (lower level of care) (n=15,941) and (b) ICU/ HDU (n=6,277) COVID-19 cases reported through CHES, England

(a)




(b)



COVID-19 Hospitalisation in England Surveillance System (CHES)


Figure 35: Ethnic group of new hospitalisations (lower level of care) (n=15,348) and ICU/ HDU (n=5,783) COVID-19 cases reported through CHES, England



UK Severe Respiratory Failure (SRF) centres admissions

Between 3 March and 15 September 2020, a total of 223 laboratory confirmed COVID-19 admissions have been reported from the 5 SRFs in England. There was no new laboratory confirmed COVID-19 admission reported in week 37.

Figure 36: Laboratory confirmed ECMO admissions (COVID-19 and non-COVID-19 confirmed) to SRFs, England



Cumulative deaths

Changes to the definitions of COVID-19 related deaths in England are described in more detail in an [accompanying PHE technical summary](#).

The current definitions used for mortality surveillance of COVID-19 in England are:

- (a) 28 day definition: A death in a person with a laboratory-confirmed positive COVID-19 test and died within (equal to or less than) 28 days of the first positive specimen date
- (b) 60 day definition: A death in a person with a laboratory-confirmed positive COVID-19 test and either: died within 60 days of the first specimen date OR died more than 60 days after the first specimen date only if COVID-19 is mentioned on the death certificate

The introduction of these definitions will affect the numbers which have been presented in past reports and therefore Figure 37 represents these differences by definition.

Figure 37: Cumulative number of deaths since week 27 by week of death and time since laboratory confirmation of COVID-19, England




Figure 38: Age/sex pyramid of laboratory confirmed COVID-19 deaths, since week 27




Table 7: Ethnic group (%) of COVID-19 deaths and time since laboratory confirmation of COVID-19, England


Ethnicity	28 day definition	60 day definition
White	83.7	88.8
Asian / Asian British	12.1	7.4
Black / African / Caribbean / Black British	2.0	1.9
Mixed / Multiple ethnic groups	0.5	0.4
Other ethnic group	1.8	1.5

Table 8: Cumulative number of COVID-19 deaths since week 27 and time since laboratory confirmation of COVID-19 by PHE Centres


PHE Centres	Number of deaths by definition	
	28 day definition	60 day definition
North East	25	84
North West	304	531
Yorkshire & Humber	158	307
West Midlands	103	232
East Midlands	132	253
East of England	150	301
London	68	153
South East	215	427
South West	29	79

Figure 39: Cumulative mortality rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2 since week 27 by (a) 28 day definition and (b) 60 day definition

(a)



(b)




Daily excess all-cause mortality, UK

Deaths occurring from 1 January to 9 September 2020 were assessed to calculate the daily excess above a baseline using age-group and region specific all cause deaths as provided daily by the General Register Office (GRO). The deaths were corrected to allow for delay to registration based on past data on these delays and the baseline was from the same day of the year in the previous 5 years +/- 7 days with an extrapolated time trend, and with 2 and 3 standard deviation (SD) limits shown (Figure 40).

Weeks in which at least 2 days exceeded the 3SD threshold are shown in Table 9 and the daily difference from the baseline by age and region is given in Figure 41. Note that as these data are by date of death with delay corrections, numbers are subject to change each week, particularly for more recent days.

No significant excess all-cause mortality was observed in week 36 overall, by age group or sub-nationally. The excess noted in week 33 coincides with a heat wave (Figure 40, 41 and Table 9).

Figure 40: Daily excess all-cause deaths in all ages, England, 1 January 2020 to 9 September 2020



^ based on same day in previous 5 years +/- 1 week with a linear trend projected

* corrected for delay to registration from death

Daily excess all-cause mortality, UK

Table 9: Excess all-cause deaths by (a) age group and (b) PHE centres , England

(a)

	Excess detected in week 36 2020?	Weeks in excess since week 10 2020
Age group		
All	x	13 to 21, 23, 33
under25	x	None
25 to 44	x	14 to 16, 32
45 to 64	x	12 to 19
65 to 74	x	12 to 19
75 to 84	x	13 to 21, 33
85+	x	13 to 21, 33


(b)

	Excess detected in week 36 2020?	Weeks in excess since week 10 2020
PHE centres		
East of England	x	14 to 19, 21
East Midlands	x	13 to 19
London	x	12 to 19,33
North East	x	14 to 21
North West	x	13 to 21, 33
South East	x	13 to 21, 33
South West	x	14 to 19, 33
West Midlands	x	13 to 20
Yorkshire and Humber	x	14 to 21, 23


Daily excess all-cause mortality, UK

Figure 41: Daily excess all-cause deaths by (a) age group and (b) PHE centres , England, 1 March 2020 to 9 September 2020

(a)



(b)



Sero-prevalence epidemiology, England

In this week's report the results from testing samples from the following sources are included:

- Healthy adult blood donors aged 17 years and older, supplied by the NHS Blood and Transplant (NHS BT collection) between weeks 13 -36. Donor samples from two different geographic regions (approximately 1000 samples per region) in England are tested each week. From week 26, an exclusion of donors aged 70 years and older donating throughout lockdown was lifted, and therefore data from the most recent sampling periods include donors in this older age group.
- Data from samples collected from patients aged 65 years and older attending for routine blood tests at one of the Royal College of General Practitioners Research and Surveillance Centre network of practices (RCGP-RSC).

Seroprevalence in Adults aged 17 years and older (Blood Donors)

The results presented here are based on testing using the Euroimmun assay for blood donor samples collected between weeks 13-36. This week's report includes the results of testing the 7th set of samples from the Midlands (week 35-36) and the 6th set of samples from the North East and Yorkshire region (week 36).

National Prevalence

Overall population weighted prevalence among blood donors aged 17 years and older in England was 5.5% (95% CI 4.9% - 6.1%) (unadjusted) or 5.8% (95% CrI 5.1% - 6.4%) after adjustment for the accuracy of the Euroimmun assay (sensitivity 83.0% and specificity 99.3%) for the period 13 Aug – 6 September (weeks 33-36). Estimates are based on 7899 samples, of which 457 were positive. This compares with 7.8% (95% CI 7.2% - 8.6%) (unadjusted) or 8.3% (95% CrI 7.5% - 9.2%) (adjusted) for the period of 6 – 29 May (weeks 19-22). Declines in prevalence can partially be explained by demographic differences in the donor population, such as later data including donors aged 70 years and older who were previously excluded from donating during lockdown. Waning immunity may also be a contributing factor to the lower prevalence.

Regional Prevalence over Time

Figure 42 shows the overall prevalence in each region over time which has been adjusted for the sensitivity and specificity of the Euroimmun assay. It is important to note that the sensitivity and specificity of assays are subject to change as further data becomes available. Sensitivity for the Euroimmun assay is based on data from testing of convalescent sera taken 3 to 6 weeks after symptom onset.

Adjusted prevalence estimates vary across the country and over time. In London where prevalence estimates are highest, overall adjusted prevalence increased from 2.6% (week 13) to 15.7% (week 21). From week 24 adjusted prevalence was lower and eventually plateaued with estimates at 8.7% in week 31 and 8.2% in week 33. Most recent London data however, show an increase in adjusted prevalence to 12.6% (95% CrI 10.2% - 15.3%). This increase could in part be due to increases in recent infection, although variability in the precise locations of sampling within London and potential changes in the characteristics of the donor population over time are also likely to be contributory factors. Given the current guidance on donation those donating when prevalence was highest in weeks 18-22 are likely to be returning now to donate again.

Prevalence estimates from other regions have been consistently lower than those from London; compatible with the lower incidence of COVID-19 observed in other surveillance systems.

Recent data from the Midlands show a higher adjusted prevalence at 6.8% (95% CrI 4.9%-8.9%) in week 35-36. This compares to 4.6% (95% CrI 3%-6.5%) in week 31-32. This observed increase is likely due to geographical variation of the population sampled, with a lower proportion of samples from Birmingham in week 31-32 compared to other sampling periods.

In the North East and Yorkshire NHS region the adjusted prevalence was 3.9% (95% CrI 2.4%-5.7%) in week 36 compared with 5% (95% CrI 3.3%-6.9%) in week 32. Similar plateauing has been seen across other regions.

Recent data from the North West show the adjusted prevalence was 7.2% (95% CrI 5.4% - 9.4%) in week 31 and more recently at 6.8% in week 35 (95% CrI 4.7-9.2%) showing a continued plateauing.

In the East of England adjusted prevalence amongst donors was 4.4% (95% CrI 2.8% - 6.3%) in the most recent data (weeks 34-35) lower than the prevalence of 6.6% (95% CrI 4.8% - 8.6%) in weeks 30-31.

Adjusted prevalence in the South East region was 3.7% (95% CrI 2.2% - 5.4%) in the latest data (week 34) similar to 3.6% (95% CrI 2.2% - 5.3%) observed in the previous survey in week 30.

The change in prevalence seen in some regions is likely to be largely driven by changes in the precise locations of sample collection. Declines in prevalence can be partially explained by demographic differences in the donor population as lockdown measures are relaxed. Examples include a reduction in attendance of regular donors in August and that donors aged 70 years and above were not allowed to donate during lockdown, but this exclusion was lifted from week 26. Waning immunity may also be a contributing factor to the lower prevalence.

Figure 42: Overall SARS-CoV-2 antibody seroprevalence (%) in blood donors by PHE centres, using Euroimmun test adjusted for sensitivity (83.0%) and specificity (99.3%) and 95% confidence intervals (dashed lines)





Figure 43: Population weighted SARS-CoV-2 antibody seroprevalence in blood donors by age group, weeks 33-36, using Euroimmun test; error bars show 95% confidence intervals



Prevalence by Age Group

Population weighted antibody prevalence (unadjusted) estimates in donors aged 70-84 years are included in the most recent data (weeks 33-36) as this age group, who were advised to shield during lockdown, have been able to return to donor clinics since week 26 (Figure 42). Prevalence is highest in the youngest age group (age 17-29) and lowest in the oldest age group (age 70-84).

Seroprevalence in Adults aged 65 years and older (Royal College of General Practitioners Research and Surveillance Centre (RCGP-RSC) network)


Prevalence was estimated from 8340 samples from patients aged between 65 and 100 years old, who had a routine blood test via the Royal College of General Practitioners Research and Surveillance Centre (RCGP-RSC) network during the period 16 March – 14 August using the Abbott and EuroImmun assays. For the Abbott assay % pos is given as % positive or indeterminate/equivocal, with an indeterminate assay cut-off of 0.8. Using a cut off of 0.8, Abbott sensitivity is estimated to be 95.7% and specificity 99.1%.

Adjusted prevalence estimates were similar using both assays and have increased slightly to between 3.3 and 3.0% during the July and August from 2.5% and 2.7% in June, although confidence intervals are overlapping (Figure 44).

Overall population weighted (by age group (65-69, 70-74, 75-59, 80+) and NHS region, prevalence among RCGP patients aged 65 years and above was 3.1% (95% CrI 2.2%-4.2%) (using the EuroImmun) or 3.4% (95% CrI 2.5%-4.5%) (using the Abbott assay) after adjustment for sensitivity and specificity for the period 1 July to 14th August and 2.8% (95% CrI 1.8-4.1%) (using the EuroImmun) or 2.3% (95% CrI 1.45%-3.3%) (using the Abbott) for the period 1-30 June.

When stratified by age, the population weighted RCGP prevalence estimates in the over 65 year olds suggest that % positive in July/August was no longer highest in individuals aged over 80, as had been observed in May and June. One possible explanation is a higher proportion of samples from care home residents from earlier time periods.


Figure 44: Eurolmmun % positive by age group and periods in RCGP patients aged 65+, 1 May-14 Aug



Global situation


Globally, up to 15 September 2020, a total of 29,099,834 cases of COVID-19 infection have been reported worldwide, including 925,676 COVID-19 related deaths.

Figure 45: Global map of cumulative COVID-19 cases



Global situation

Figure 46: Global map of weekly COVID-19 case incidence rate per 100,000, week 37 2020



PHE has delegated authority, on behalf of the Secretary of State, to process Patient Confidential Data under Regulation 3 The Health Service (Control of Patient Information) Regulations 2002

<http://www.legislation.gov.uk/ukxi/2002/1438/regulation/3/made>. Regulation 3 makes provision for the processing of patient information for the recognition, control and prevention of communicable disease and other risks to public health.