

Department for Environment, Food and Rural Affairs

The Expert Committee on Pesticide Residues in Food (PRiF)

Report on the pesticide residues monitoring programme: Quarter 2 2018

December 2018





© Crown copyright 2018

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gsi.gov.uk

This publication is available at www.gov.uk/government/groups/expert-committee-on-pesticide-residues-in-food-prif

For further details on information contained in this report, previous surveys or information concerning pesticide residues in food, please contact:

Expert Committee on Pesticide Residues in Food
HSE's Chemicals Regulation Division
Mallard House
Kings Pool
Peasholme Green
York
YO1 7PX

prif@hse.gov.uk

Contents

Introduction and summary results	5
Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF).....	5
Chair's summary of results	6
Summary table of all results	7
Summary of samples with residues over the MRL.....	9
Summary of Rapid Alert Notifications sent to FSA	14
Section 1: findings by food.....	18
Animal fats.....	18
Apples.....	20
Aubergine	22
Bananas	23
Beans with pods	25
Beef	28
Berries and small fruits	31
Broccoli.....	34
Cheese (soft).....	36
Chinese cabbage.....	39
Cream.....	41
Curry leaves	44
Eggs	48
Fish (white)	49
Game.....	51
Grapefruit.....	53
Grapes.....	55
Lettuce.....	57
Melon.....	59
Milk	62
Mushrooms.....	64
Mushrooms (speciality).....	65
Okra.....	68
Pears	70
Peas without pods	72
Peppers	75
Pineapple.....	77
Potatoes	79
Soft citrus.....	81
Speciality vegetables.....	85
Section 2: Sample details and supplier responses	87
Sample details	87
Supplier responses.....	89
Section 3: HSE assessment of risk.....	91
Assessment of Risk to Human Health: Short-term intake estimates.....	96
Section 4: issues arising in this report and updates on previous reports	113
Issues arising in this report.....	113
Follow-up from Previous Reports	116

Brand name details of samples where follow-up action is now complete	118
In our next report:	120
Section 5: background and reference	121
Glossary	133
133	

Introduction and summary results

Introduction to the work of the Expert Committee on Pesticide Residues in Food (PRiF)

The PRiF's role is to give Ministers, the Director of the Health and Safety Executive (HSE) and the Chief Executive of the Food Standards Agency (FSA) independent advice on the UK government's national rolling programme of surveys, in particular:

- the planning of surveillance programmes for pesticide residues in the UK food supply and the evaluation of the results;
- Procedures for sampling, sample processing, new methods of analysis, the assessment of variability of pesticide residues in food and related issues.

[More information about PRiF](#)

HSE working under Defra's authority has official responsibility to organise a monitoring programme of UK food for pesticide residues. The programme is made up of a risk-based national rolling programme of surveys and also includes participation in EU-wide monitoring. HSE is also responsible for considering the safety to people who eat the food (in co-operation with the Food Standards Agency if necessary) and following up adverse or unexpected results. They are also responsible for determining whether food is compliant with the law, specifically, whether any pesticide residue found is within the Maximum Residue Level. Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in produce, which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not approved for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory analysis. This provides a mechanism for statutory controls on pesticides in produce which is put into circulation and for monitoring the correct use of these chemicals.

Chair's summary of results

This is our second quarterly report for 2018. During this year's surveillance programme we are looking for a range of up to 371 pesticides in the fruit and vegetable surveys. This quarter's programme surveyed 823 samples of 30 different foods (see contents page for a full list).

55 of the samples contained residues above the legal Maximum Residue Level (the maximum permitted levels by law). These results are in the surveys of beans with pods, beef, berries, broccoli, cheese, cream, curry leaves, game, grapes, melon, mushrooms (speciality), okra, pears, peas without pods and soft citrus. [A summary table of all the results is below.](#)

However, we do not think the findings of chlorate residues in cheese, cream, melon and peas without pods should be treated as breaches of the legislation, and we have not highlighted them as such in the brand name annex. You can read updated information about work currently being done on chlorate residues in [section 4](#).

HSE undertakes a screening risk assessment for every residue found, to determine whether the residues could lead to intakes above the relevant reference (safety) doses. HSE also produces [detailed risk assessments](#) for every case where the actual residue level found could lead to an intake above the safety levels. We have looked carefully at all these findings including the risk assessments. In most cases the presence of the residues found would be unlikely to have had any effect on the health of the people who ate the food.

Full details of suppliers and retailers of the food sampled, and full analytical results, are available on data.gov.uk as ODF (Open Document Format) spreadsheet files. There is no change to the level of detail or data structure previously included in our reports, but we hope the new data format is useful for people wanting to look at the individual results in more detail.

We asked suppliers and the authorities of the exporting countries for an explanation of our findings. Any responses we have received are available in [Section 2](#): Sample details and supplier responses.

Dr Paul Brantom

Chairman of the Expert Committee on Pesticide Residues in Food

Summary table of all results

Food	Analysed	With residues at or below the MRL	With residues above the MRL	With residues of non approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Animal fats	24	0	0	0	0	0	0
Apples	18	11	0	0	9	7	0
Aubergine	12	4	0	0	0	0	0
Banana	18	10	0	0	8	9	1
Beans with pods	28	12	3	1	12	1	0
Beef	24	5	3	0	2	2	1
Berries	48	26	2	2	13	2	0
Broccoli	24	9	1	1	4	3	0
Cheese (soft)	30	4	4	0	2	1	0
Chinese cabbage	22	11	0	3	6	0	0
Cream	36	2	15	0	0	3	1
Curry leaves	24	10	5	0	5	8	4
Eggs	25	0	0	0	0	5	0
Fish (white)	30	6	0	0	0	0	0
Game	19	0	1	0	0	0	0
Grapefruit	24	24	0	0	24	0	0
Grapes	37	36	1	0	34	0	0
Lettuce	15	6	0	0	4	2	0
Melon	30	16	3	0	8	1	0

Food	Analysed	With residues at or below the MRL	With residues above the MRL	With residues of non approved pesticides (UK only)	With multiple residues	Organic samples tested	Organic samples with residues
Milk	72	3	0	0	0	22	0
Mushrooms	18	2	0	0	0	4	0
Mushrooms (speciality)	24	6	2	5	1	2	0
Okra	26	5	3	0	4	0	0
Pears	18	8	1	0	9	8	0
Peas without pods	36	8	9	0	5	0	0
Peppers	19	8	0	0	7	0	0
Pineapple	19	12	0	0	5	0	0
Potatoes	43	22	0	0	7	2	0
Soft citrus	36	34	2	0	35	0	0
Speciality vegetables	24	15	0	2	7	0	0

Summary of samples with residues over the MRL

PRiF Sample ID	Food	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg)	MRL exceedance after allowing for measurement uncertainty
Beans with pods						
0113/2018	Gawar Beans	India	dimethoate (sum)	0.02	0.01*	Yes
0139/2018	Yard Long Beans	Malaysia	dimethoate (sum)	0.06	0.01*	Yes
0640/2018	Hyacinth Beans	Malaysia	carbendazim (sum)	1.7	0.2	Yes
			chlorfenapyr	0.6	0.01*	Yes
			dithiocarbamates	3.9	1	Yes
			Fluopicolide	0.3	0.01*	Yes
			lambda-cyhalothrin	0.3	0.2	No
			methomyl (sum)	0.6	0.1	Yes
			propamocarb (sum)	1.5	0.1	Yes
Beef						
2283/2018	Steak Mince	UK	BAC (sum)	0.2	0.1	No
3433/2018	Top Rump Joint	UK	DDAC (sum)	0.2	0.1	No
3445/2018	Mincd Beef	UK	BAC (sum)	0.3	0.1	Yes
Berries						
1851/2018	Fresh: Blackberries	Mexico	flubendiamide	0.05	0.01*	Yes
			propamocarb (sum)	0.02	0.01*	No
4243/2018	Fresh: Blackberries	Mexico	permethrin (sum)	0.2	0.05*	Yes
Cheese (soft)						
1596/2018	Brie	France	chlorate	0.03	0.01	No
1789/2018	Mozzarella	Italy	chlorate	0.1	0.01	No

PRiF Sample ID	Food	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg)	MRL exceedance after allowing for measurement uncertainty
2961/2018	Mozzarella	Italy	BAC (sum)	1.9	0.1	No
			Chlorate	0.2	0.01	No
2949/2018	Ricotta	Italy	Chlorate	0.04	0.01	No
Cream						
1939/2018	Creme fraiche	UK	chlorate	0.02	0.01	No
2825/2018	Creme fraiche	UK	chlorate	0.04	0.01	No
3436/2018	Creme fraiche	UK	chlorate	0.06	0.01	No
3764/2018	Creme fraiche	UK	chlorate	0.04	0.01	No
2843/2018	Double cream	UK	chlorate	0.03	0.01	No
2948/2018	Double cream	UK	chlorate	0.02	0.01	No
1696/2018	Single cream	UK	chlorate	0.02	0.01	No
1832/2018	Single cream	UK	chlorate	0.02	0.01	No
2962/2018	Single cream	UK	chlorate	0.02	0.01	No
3420/2018	Single cream	UK	chlorate	0.02	0.01	No
4234/2018	Single cream	UK	chlorate	0.02	0.01	No
1742/2018	Sour cream	UK	chlorate	0.03	0.01	No
2505/2018	Sour cream	UK	chlorate	0.03	0.01	No
1791/2018	Whipping cream	UK	chlorate	0.02	0.01	No
1917/2018	Whipping cream	UK	chlorate	0.02	0.01	No
Curry leaves						
2428/2018	Organic Curry Leaves	Sri Lanka	profenofos	0.2	0.13	Yes

PRiF Sample ID	Food	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg)	MRL exceedance after allowing for measurement uncertainty
2430/2018	Dried Curry Leaves	India	acephate	0.06	0.05	No
			bifenthrin	1.4	0.05	Yes
			ethion	0.2	0.03	Yes
			profenofos	2.3	0.13	Yes
			tebuconazole	1.3	0.13	Yes
			triazophos	1.6	0.03	Yes
3952/2018	Dried Curry Leaves	India	acephate	0.06	0.05	No
			bifenthrin	1.4	0.05	Yes
			diphenylamine	0.3	0.13	Yes
			ethion	0.5	0.03	Yes
			profenofos	1.9	0.13	Yes
			tebuconazole	0.4	0.13	Yes
			triazophos	4	0.03	Yes
4053/2018	Curry Leaves	India	profenofos	0.3	0.05	Yes
4206/2018	Curry Leaves	India	profenofos	0.8	0.05	Yes
Game						
2289/2018	Venison	UK	BAC (sum)	0.8	0.1	Yes
Grapes						
0511/2018	Iniagrape One Grapes	Chile	captan (sum)	0.05	0.03*	No
Melon						
2823/2018	Cantaloupe	Honduras	chlorothalonil	1.2	1	No
1859/2018	Frozen Watermelon	Guatemala	chlorate	0.04	0.01	No

PRiF Sample ID	Food	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg)	MRL exceedance after allowing for measurement uncertainty
2387/2018	Frozen Watermelon	Guatemala	chlorate	0.2	0.01	No
Mushrooms (speciality)						
2943/2018	Oyster Mushrooms	UK	chlormequat	4.6	0.9	Yes
4224/2018	Oyster Mushrooms	UK	chlormequat	4.5	0.9	Yes
Okra						
0641/2018	Fresh: Okra	Jordan	indoxacarb	0.03	0.02*	No
			thiacloprid	0.04	0.01*	Yes
1737/2018	Fresh: Okra	India	diuron	0.02	0.01*	Yes
3416/2018	Frozen; Okra	India	amitraz (sum)	0.07	0.05*	No
			diafenthiuron	0.02	0.01*	Yes
			nitenpyram	0.03	0.01*	Yes
Pears						
1935/2018	Conference Pears	the Netherlands	chlormequat	0.1	0.07	No
Peas without pods						
3425/2018	Hand Shelled Peas	Kenya	chlorate	0.4	0.01	No
3947/2018	Hand Shelled Garden Peas	Guatemala	chlorate	0.02	0.01	No
4059/2018	Hand Shelled Garden Peas	Guatemala	chlorate	0.02	0.01	No
4132/2018	Hand Shelled Garden Peas	Guatemala	chlorate	0.03	0.01	No
1886/2018	Garden Peas	UK	chlorate	0.04	0.01	No
1552/2018	Garden Peas in Water	UK	chlorate	0.2	0.01	No
1983/2018	British Garden Peas in Water	UK	chlorate	0.04	0.01	No
2822/2018	Petits pois in Water	France	chlorate	0.02	0.01	No

PRiF Sample ID	Food	Country of Origin	Pesticide Detected	Residue Detected (mg/kg)	MRL (mg/kg)	MRL exceedance after allowing for measurement uncertainty
4246/2018	Petits pois in Water	France	chlorate	0.02	0.01	No
Soft citrus						
0074/2018	Clementines With Leaves	Spain	propiconazole	6	5	No
0578/2018	Satsuma	Peru	thiabendazole	8.1	7	No

* Maximum Residue Levels set at the LOD (LOD MRL): These MRLs are set at a default level, i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. Either insufficient trials data are available, on which to set a maximum residue level or there may be no use of the pesticide on that crop in the EU. However, they may be permitted elsewhere.

Summary of Rapid Alert Notifications sent to FSA

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)	RASFF Number
Beans with pods									
0640/2018	18/06/2018	Hyacinth Beans	Malaysia	Tri-Imp	Units 43-45 St James Market, Essex Street, Bradford BD3 7PN	None stated	Zahid Al Ahad 38200 Ringley Cameron, Highlands Pahang, Malaysia	abamectin (sum) 0.02 (MRL = 0.03) carbendazim (sum) 1.7 (MRL = 0.2) chlorfenapyr 0.6 (MRL = 0.01*) clothianidin 0.03 (MRL = 0.2) difenoconazole 0.01 (MRL = 1) dithiocarbamates 3.9 (MRL = 1) fluopicolide 0.3 (MRL = 0.01*) indoxacarb 0.07 (MRL = 0.5) lambda-cyhalothrin 0.3 (MRL = 0.2) methomyl (sum) 0.6 (MRL = 0.1) myclobutanil 0.3 (MRL = 0.8) propamocarb (sum) 1.5 (MRL = 0.1) tebuconazole 0.2 (MRL = 2) thiamethoxam 0.2 (MRL = 0.3) trifloxystrobin 0.01 (MRL = 1)	2018.2195
Curry leaves									
3952/2018	30/04/2018	Dried Curry Leaves	India		120 Gladstone Street,	Top-op	Top-Op Foods Ltd 5-	acephate 0.06 (MRL = 0.05)	

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)	RASFF Number
				Ribbon Foods Ltd	Darlington DL3 6JZ		7 Garland Road, Stanmore HA7 1QU	acetamiprid 2.6 (MRL = 7.8) azoxystrobin 0.1 (MRL = 182) bifenthrin 1.4 (MRL = 0.05) carbendazim (sum) 0.03 (MRL = 0.26) chlorpyrifos 0.06 (MRL = 0.13) chlorantraniliprole 0.04 (MRL = 52) difenoconazole 0.8 (MRL = 10.4) dimethoate (sum) 0.01 (MRL = 0.05) diphenylamine 0.3 (MRL = 0.13) ethion 0.5 (MRL = 0.03) flonicamid (sum) 0.8 (MRL = 15.6) methamidophos 0.02 (MRL = 0.05) monocrotophos 0.02 (MRL = 0.05) profenofos 1.9 (MRL = 0.13) pendimethalin 0.02 (MRL = 1.56) tebuconazole 0.4 (MRL = 0.13) triazophos 4 (MRL = 0.03)	
2430/2018	16/05/2018	Dried Curry Leaves	India	Amazon.co.uk	Patriot Court, 1-9 The Grovetown, Slough,	Top-op	Top-Op Foods Ltd 5-7 Garland Road,	acephate 0.06 (MRL = 0.05) acetamiprid 1.9 (MRL = 7.8) azoxystrobin 0.2 (MRL = 182)	

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)	RASFF Number
					Berkshire SL1 1QP		Stanmore HA7 1QU	bifenthrin 1.4 (MRL = 0.05) carbofuran (sum) 0.01 (MRL = 0.05) carbendazim (sum) 0.02 (MRL = 0.26) chlorpyrifos 0.01 (MRL = 0.13) chlorantraniliprole 0.07 (MRL = 52) cypermethrin (sum) 1.7 (MRL = 5.2) difenoconazole 0.1 (MRL = 10.4) ethion 0.2 (MRL = 0.03) flonicamid (sum) 0.4 (MRL = 15.6) hexaconazole 0.05 (MRL = 0.05) indoxacarb 0.01 (MRL = 5.2) monocrotophos 0.01 (MRL = 0.05) Novaluron 0.03 (MRL = 0.03) profenofos 2.3 (MRL = 0.13) propargite 0.02 (MRL = 0.05) pendimethalin 0.02 (MRL = 1.56) tebuconazole 1.3 (MRL = 0.13) triazophos 1.6 (MRL = 0.03)	

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. Either insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop in the EU. However, they may be permitted elsewhere.

Section 1: findings by food

Animal fats

Summary of results

In a survey of 24 samples of animal fats collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

No residues were detected at or above the reporting limit

Survey design

The animal fat samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 38 pesticide residues

Dripping

- 9 samples came from the UK

Goose Fat

- 1 sample came from the UK
- 1 sample came from the EU

Lard

- 9 samples came from the UK
- 3 samples came from the EU

Suet

- 1 sample came from the UK

Pesticide residues detected from those sought

- 24 samples contained no residues from those sought

- None of the samples contained residues above the reporting level
- None of the samples were labelled as organic.

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Apples

Summary of results

In a survey of 18 samples of apples collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

One of the samples contained a residue of dithiocarbamates at a level within the MRL. Currently, it is not possible to determine analytically which specific pesticide in the dithiocarbamate group was present. Therefore, HSE undertook a consumer risk assessment based on the assumption that ziram was used. Based on this assumption it was concluded that some people might experience slight loss of appetite after eating or drinking large portions (97.5th percentile consumption) of apple containing the highest levels found in this report. Such effects would be expected to be minor and reversible and we consider the likelihood of an effect on health to be low. In the future it would be valuable if additional cost effective analytical tests could be developed to enable a more specific risk assessment.

Survey design

The apple samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

18 samples were tested for up to 367 pesticide residues

Eating

- 1 sample came from the UK
- 13 samples were imported from outside the EU
- 4 samples came from the EU

Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- 11 samples contained residues above the reporting level
- None of the samples contained residues above the MRL

- 7 samples were labelled as organic. None contained residues from those sought

Multiple residues

9 samples contained residues of more than one pesticide

- 3 samples contained 2 residues
- 3 samples contained 3 residues
- 1 sample contained 4 residues
- 2 samples contained 5 residues

Risk assessments

One sample of apple contained a residue of dithiocarbamate at levels where the effect on health needed to be considered in more detail. The highest level detected was 2.2 mg/kg. HSE's risk assessment was based on the assumption that the residue arose from ziram and concluded that some people might experience slight loss of appetite after eating or drinking large portions (97.5th percentile consumption) of apple containing the highest levels found in this report, but we consider the likelihood of an effect on health to be low. Such effects would be expected to be minor and reversible. Full risk assessment is available at page **Error! Reference source not found.****Error! Reference source not found.****Error! Bookmark not defined.**

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Aubergine

Summary of results

In a survey of 12 samples of aubergines collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year aubergines are being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

None of the residues detected would be expected to have an effect on health

Survey design

The aubergine samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

12 samples were tested for up to 365 pesticide residues

- 1 sample was imported from outside the EU
- 11 samples came from the EU

Pesticide residues detected from those sought

- 8 samples contained no residues from those sought
- 4 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

None of the samples contained residues of more than one pesticide

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Bananas

Summary of results

In a survey of 18 samples of bananas collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year bananas are being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The banana samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

18 samples were tested for up to 366 pesticide residues

- 18 samples were imported from outside the EU

Pesticide residues detected from those sought

- 8 samples contained no residues from those sought
- 10 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- 9 samples were labelled as organic. 1 contained 2 residues from those sought

Multiple residues

8 samples contained residues of more than one pesticide

- 1 sample contained 2 residues
- 3 samples contained 3 residues
- 4 samples contained 4 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Organic sample with a residue

The Secretariat has written to the supplier of the sample of organic bananas from Mexico with residues of boscalid and fenpropidin which are not permitted in organic food production. Defra's Organic Farming branch and the organic certification organisation were also informed.

Beans with pods

Summary of results

In a survey of 28 samples of beans with pods collected between April and June 2018, 3 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

Based on the Chemicals Regulation Division's risk assessment of the residues detected we consider an effect on health to be unlikely.

Survey design

The beans with pod samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

28 samples were tested for up to 366 pesticide residues

Green Beans

- 4 samples came from the UK
- 14 samples were imported from outside the EU

Speciality Beans

- 10 samples were imported from outside the EU

Pesticide residues detected from those sought

- 13 samples contained no residues from those sought
- 15 samples contained residues above the reporting level
- 3 samples contained residues above the MRL
- 1 sample was labelled as organic. It didn't contain any residues from those sought

Multiple residues

12 samples contained residues of more than one pesticide

- 3 samples contained 2 residues
- 5 samples contained 3 residues
- 1 sample contained 5 residues
- 1 sample contained 6 residues
- 1 sample contained 7 residues
- 1 sample contained 15 residues

Residues measured above the MRL

The laboratory detected 9 residues above the MRL in beans with pods

- 1 sample of gawar beans from India contained a residue of dimethoate (sum) at 0.02 mg/kg. The MRL is 0.01** mg/kg.
- 1 sample of yard long beans from Malaysia contained a residue of dimethoate (sum) at 0.06 mg/kg. The MRL is 0.01* mg/kg.
- 1 sample of hyacinth beans from Malaysia contained residues of:
 - carbendazim (sum) at 1.7 mg/kg, the MRL is 0.02 mg/kg
 - chlorfenapyr at 0.6 mg/kg, the MRL is 0.01* mg/kg
 - dithiocarbamates at 3.9 mg/kg, the MRL is 1 mg/kg
 - fluopicolide at 0.3 mg/kg, the MRL is 0.01* mg/kg
 - lambda-cyhalothrin at 0.3 mg/kg, the MRL is 0.2 mg/kg
 - methomyl (sum) at 0.6 mg/kg, the MRL is 0.1 mg/kg
 - propamocarb (sum) at 1.5 mg/kg, the MRL is 0.1 mg/kg

Risk assessments

- 1 sample of beans with pods (Hyacinth beans) contained a residue of methomyl at levels where the effect on health needed to be considered in more detail. The highest level detected was 0.6 mg/kg HSE's risk assessment indicated that if infants ate large portions of beans with pods containing methomyl at 0.6 mg/kg. Their intake of methomyl could be 120% of the Acute Reference dose. This intake is 83 times lower than a dose which caused no observed adverse effects in a study on

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

acute neurotoxicity in rats. We consider the reduced factor of 83 (in comparison to the value of 100 usually used by toxicologists) still enough to make an effect on health unlikely. For full details see the risk assessment at page 99.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

RASFFs issued

HSE sent a draft notification for the following samples to FSA to raise through the EC's Rapid Alert System for Food and Feed (RASFF) (see glossary for more details). RASFF notification 2018.2195 was subsequently issued

- 1 sample from Malaysia containing methomyl at 0.6 mg/kg

Beef

Summary of results

In a survey of 24 samples of beef collected between April and June 2018, 3 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

DDT

One sample contained a residue of DDT. The use of DDT is banned or heavily restricted in many countries because the residues take a long time to breakdown in the environment and can accumulate in fatty tissue.

An interpretation of the analytical results shows that the DDT residue found was in the form of DDE which indicates historical use. More detailed information about DDT residues is on page 115 of this report.

Survey design

The beef samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 104 pesticide residues

- 19 samples came from the UK
- 5 samples came from the EU

The country of origin on the packaging does not necessarily indicate where the animal was from. It may be where the meat was processed or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 16 samples contained no residues from those sought
- 8 samples contained residues above the reporting level
- 3 samples contained residues above the MRL

- 2 samples were labelled as organic. 1 contained residues from those sought

Multiple residues

2 samples contained residues of more than one pesticide

- 2 samples contained 2 residues

Residues measured above the MRL

The laboratory detected 3 residues above the MRL in beef

- 1 sample of steak mince from UK contained a residue of BAC (sum) at 0.2 mg/kg. The MRL is 0.1* mg/kg
- 1 sample of top rump joint from UK contained a residue of DDAC (sum) at 0.2 mg/kg. The MRL is 0.1 mg/kg
- 1 sample of minced beef from UK contained a residue of BAC (sum) at 0.3 mg/kg. The MRL is 0.1 mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

Organic sample with a residue

The Secretariat has written to the supplier of the sample of organic beef mince from the UK with a residue of BAC (sum). This is not permitted for use as a pesticide in organic food production. Defra's Organic Farming branch and the organic certification organisation were also informed.

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Berries and small fruits

Summary of results

In a survey of 48 samples of berries collected between April and June 2018, 2 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The berry samples were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

48 samples were tested for up to 365 pesticide residues

Fresh: Blackberries

- 3 samples were imported from outside the EU
- 4 samples came from the EU

Fresh: Blueberries

- 8 samples were imported from outside the EU
- 20 samples came from the EU

Frozen: Blackberries

- 3 samples were imported from outside the EU
- 1 sample came from the EU

Frozen: Blueberries

- 3 samples came from the UK
- 4 samples were imported from outside the EU
- 2 samples came from the EU

The country of origin on the packaging does not necessarily indicate where the frozen berries were grown. It may be where the berries were processed or where they were packed for consumer purchase.

Pesticide residues detected from those sought

- 20 samples contained no residues from those sought
- 28 samples contained residues above the reporting level
- 2 samples contained residues above the MRL
- 2 samples were labelled as organic. Neither contained residues from those sought

Multiple residues

13 samples contained residues of more than one pesticide

- 9 samples contained 2 residues
- 2 samples contained 3 residues
- 2 samples contained 4 residues

Residues measured above the MRL

The laboratory detected 3 residues above the MRL in berries

- 1 sample of blackberries from Mexico contained residues of:
 - flubendiamide at 0.05 mg/kg, the MRL is 0.01** mg/kg
 - propamocarb (sum) at 0.02 mg/kg, the MRL is 0.01* mg/kg
- 1 sample of blackberries from Mexico contained a residue of permethrin (sum) at 0.2 mg/kg. The MRL is 0.05* mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

Further investigation: Suspected illegal use

We have passed details of 2 samples of frozen blueberries from the UK that contained a residue of captan and phosmet which are not approved for use on blueberries in the UK to HSE. HSE is investigating; brand name details will not be published until the investigations are complete.

Broccoli

Summary of results

In a survey of 24 samples of broccoli collected between April and June 2018, 1 sample contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year broccoli is being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The broccoli samples were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 362 pesticide residues

Fresh

- 1 sample came from the UK
- 15 samples came from the EU

Frozen

- 4 samples came from the UK
- 4 samples came from the EU

Pesticide residues detected from those sought

- 14 samples contained no residues from those sought
- 10 samples contained residues above the reporting level
- 1 sample contained a residue above the MRL
- 3 samples were labelled as organic. None contained residues from those sought

Multiple residues

4 samples contained residues of more than one pesticide

- 3 samples contained 2 residues
- 1 sample contained 3 residues

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in broccoli

- 1 sample of frozen broccoli from UK contained a residue of fluazifop-p (sum) at 0.03 mg/kg. The MRL is 0.01** mg/kg.

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

Further investigation: Suspected illegal use

We have passed details of a sample of frozen broccoli florets from the UK that contained a residue of fluazifop-p (sum) which is not approved for use on broccoli in the UK to HSE. HSE is investigating; brand name details will not be published until the investigations are complete.

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Cheese (soft)

Summary of results

In a survey of 30 samples of cheese collected between April and June 2018, 4 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

DDT

Two samples of feta contained a residue of DDT. The use of DDT is banned or heavily restricted in many countries because the residues take a long time to breakdown in the environment and can accumulate in fatty tissue.

An interpretation of the analytical results shows that the DDT residue found was in the form of DDE which indicates historical use. More detailed information about DDT residues is on page 115 of this report.

Chlorate

We found chlorate over the default MRL in four of the samples. However, we do not think that these findings should be treated as breaches of the legislation, and we have not highlighted them as such in the brand name annex.

We expected to find residues of chlorate above the MRL in dairy products, where disinfection could be used as part of routine processes or where the food is packed in water. We are testing a limited number of foods for chlorate in 2018, as we did in 2017, to provide evidence on consumer safety and confirm that it is necessary to review the existing default MRL in order to take account of non-pesticide sources. Chlorine-based treatments of drinking and irrigation water as well as chlorine-based surface disinfectants which are widely used to ensure microbiological safety. We agree with HSE and the FSA that the current MRL does not take account of these often-unavoidable sources.

Following the HSE's risk assessment, we do not expect any of the residues we found to have an effect on health. We do not think any change in production practice by the brand-owners or manufacturers is needed in response to these findings.

This adds to a growing body of evidence, from both official monitoring across the EU and from the food and farming industries, about the incidence of chlorate residues in food.

More information on work being done on chlorate in the diet is available in [section 4](#)

Survey design

The cheese samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

30 samples were tested for up to 105 pesticide residues

Brie

- 2 samples came from the UK
- 6 samples came from the EU

Camembert

- 6 samples came from the EU

Cottage Cheese

- 4 samples came from the UK

Cream Cheese

- 3 samples came from the EU

Feta

- 3 samples came from the EU

Mozzarella

- 4 samples came from the EU

Ricotta

- 2 samples came from the EU

The country of origin on the packaging does not necessarily indicate where the milk was from. It may be where the cheese was made or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 23 samples contained no residues from those sought
- 7 samples contained residues above the reporting level

- 4 samples contained residues above the MRL
- 1 sample was labelled as organic. It didn't contain any residues from those sought

Multiple residues

1 sample contained residues of more than one pesticide

- 1 sample contained 2 residues

Residues measured above the MRL

The laboratory detected 5 residues above the MRL in soft cheese

- 1 sample of brie from France contained a residue of chlorate at 0.03 mg/kg. The MRL is 0.01* mg/kg.
- 1 sample of mozzarella from Italy contained a residue of chlorate at 0.1 mg/kg. The MRL is 0.01 mg/kg
- 1 sample of mozzarella from Italy contained residues of:
 - BAC (sum) at 1.9 mg/kg, the MRL is 0.1 mg/kg
 - Chlorate at 0.2 mg/kg, the MRL is 0.01 mg/kg
- 1 sample of ricotta from Italy contained a residue of chlorate at 0.04 mg/kg. The MRL is 0.01 mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received can be found below.

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Chinese cabbage

Summary of results

In a survey of 22 samples of Chinese cabbages collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The Chinese cabbage samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

22 samples were tested for up to 363 pesticide residues

Chinese Leaf

- 3 samples came from the UK
- 1 sample came from the EU

Choy Sum

- 2 samples came from the UK
- 2 samples came from the EU

Pak Choi

- 10 samples came from the UK
- 4 samples came from the EU

Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 11 samples contained residues above the reporting level

- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

6 samples contained residues of more than one pesticide

- 4 samples contained 2 residues
- 1 sample contained 3 residues
- 1 sample contained 5 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Further investigation: Suspected illegal use

We have passed to HSE details of 3 samples of UK Chinese cabbage that contained a residue of a pesticide which is not approved for use on that crop in the UK

- 1 sample of choi sum that contained a residue of acetamiprid
- 1 sample of pak choi from the UK that contained a residue of acetamiprid
- 1 sample of pak choi that contained a residue of fluopyram

HSE is investigating; brand name details will not be published until the investigations are complete.

Cream

Summary of results

In a survey of 36 samples of cream collected between April and June 2018, 15 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Chlorate

We found chlorate over the default MRL in seven of the samples. However, we do not think that these findings should be treated as breaches of the legislation, and we have not highlighted them as such in the brand name annex.

We expected to find residues of chlorate above the MRL in dairy products, where disinfection could be used as part of routine processes. We are testing a limited number of foods for chlorate in 2018, as we did in 2017, to provide evidence on consumer safety and confirm that it is necessary to review the existing default MRL in order to take account of non-pesticide sources. In particular, chlorine-based treatments of drinking and irrigation water as well as chlorine-based surface disinfectants which are widely used to ensure microbiological safety. We agree with HSE and the FSA that the current MRL does not take account of these often-unavoidable sources.

Following the HSE's risk assessment, we do not expect any of the residues we found to have an effect on health. We do not think any change in production practice by the brand-owners or manufacturers is needed in response to these findings.

This adds to a growing body of evidence, from both official monitoring across the EU and from the food and farming industries, about the incidence of chlorate residues in food.

More information on work being done on chlorate is on page 113

Survey design

The cream samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

36 samples were tested for up to 106 pesticide residues

Crème fraîche

- 5 samples came from the UK
- 1 sample came from the EU

Double cream

- 18 samples came from the UK

Single cream

- 8 samples came from the UK

Sour cream

- 2 samples came from the UK

Whipping cream

- 2 samples came from the UK

The country of origin on the packaging does not necessarily indicate where the milk was from. It may be where the cream was processed or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 19 samples contained no residues from those sought
- 17 samples contained residues above the reporting level
- 15 samples contained residues above the MRL
- 3 samples were labelled as organic. 1 contained residues from those sought

Multiple residues

None of the samples contained residues of more than one pesticide

Residues measured above the MRL

The laboratory detected 15 residues above the MRL in chlorate

- 1 sample of crème fraîche from UK contained a residue of chlorate at 0.02 mg/kg. The MRL is 0.01* mg/kg.

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

- 2 samples of crème fraiche from UK contained residues of chlorate at 0.04 mg/kg. The MRL is 0.01 mg/kg.
- 1 sample of crème fraiche from UK contained a residue of chlorate at 0.06 mg/kg. The MRL is 0.01 mg/kg.
- 1 sample of double cream from UK contained a residue of chlorate at 0.03 mg/kg. The MRL is 0.01 mg/kg.
- 1 sample of double cream from UK contained a residue of chlorate at 0.02 mg/kg. The MRL is 0.01 mg/kg.
- 5 samples of single cream from UK contained residues of chlorate at 0.02 mg/kg. The MRL is 0.01 mg/kg.
- 2 samples of sour cream from UK contained residues of chlorate at 0.03 mg/kg. The MRL is 0.01 mg/kg.
- 2 samples of whipping cream from UK contained residues of chlorate at 0.02 mg/kg. The MRL is 0.01 mg/kg.

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received can be found below.

Organic sample with a residue

The Secretariat has written to the supplier of the sample of organic crème fraiche from the UK with a residue of chlorate. Chlorate is not permitted for use as a plant protection product in organic food production. However as with other foods we think it is likely that the residue is from use of disinfectants (biocides) on surfaces and equipment to ensure microbiological food safety (see page 113 for more information). Defra's Organic Farming branch and the organic certification organisation were also informed.

Curry leaves

Summary of results

In a survey of 24 samples of curry leaves collected between April and June 2018, 5 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF). The results of this testing are published throughout the year.

Comments by the PRiF

Five samples contained residues above the MRL, including Indian produce which is still subject to increased border control by UK Port Authorities.

One of these samples contained residues of triazophos at a level which although HSE concluded was unlikely to have an effect on health was an exceedance of the acute reference dose. We will ensure these results continue to be notified to the Food Standards Agency (FSA) as part of the border control exercise and plan to test this produce throughout 2018 and 2019.

Two samples contained residues of monocrotophos. Monocrotophos is an insecticide that has not been authorised for use in the EU since 2003. There is uncertainty about the potential for monocrotophos to cause genetic damage; therefore, on a precautionary basis we consider any findings of monocrotophos in food as not desirable. However, considering the very low intakes any risks are likely to be low. A more detailed explanation is with the risk assessments on page 99.

Survey design

The curry leaf samples were bought by a market research company from retail outlets across the UK and online.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 357 pesticide residues

Dried

- 1 sample came from the UK
- 20 samples were imported from outside the EU

Fresh

- 3 samples were imported from outside the EU

Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 15 samples contained residues above the reporting level
- 5 samples contained residues above the MRL
- 8 samples were labelled as organic. 4 contained residues from those sought

Multiple residues

5 samples contained residues of more than one pesticide

- 3 samples contained 3 residues
- 1 sample contained 18 residues
- 1 sample contained 21 residues

Residues measured above the MRL

The laboratory detected 16 residues above the MRL in curry leaves

- 1 sample of organic curry leaves from Sri Lanka contained a residue of profenofos at 0.2 mg/kg. The MRL is 0.13* mg/kg.
- 1 sample of dried curry leaves from India contained residues of:
 - acephate at 0.06 mg/kg, the MRL is 0.05 mg/kg
 - bifenthrin at 1.4 mg/kg, the MRL is 0.05 mg/kg
 - ethion at 0.2 mg/kg, the MRL is 0.03 mg/kg
 - profenofos at 2.3 mg/kg, the MRL is 0.13 mg/kg
 - tebuconazole at 1.3 mg/kg, the MRL is 0.13 mg/kg
 - triazophos at 1.6 mg/kg, the MRL is 0.03 mg/kg
- 1 sample of dried curry leaves from India contained residues of:

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

- acephate at 0.06 mg/kg, the MRL is 0.05 mg/kg
 - bifenthrin at 1.4 mg/kg, the MRL is 0.05 mg/kg
 - diphenylamine at 0.3 mg/kg, the MRL is 0.13 mg/kg
 - ethion at 0.5 mg/kg, the MRL is 0.03 mg/kg
 - profenofos at 1.9 mg/kg, the MRL is 0.13
 - tebuconazole at 0.4 mg/kg, the MRL is 0.13 mg/kg
 - triazophos at 4 mg/kg, the MRL is 0.03 mg/kg
- 1 sample of curry leaves from India contained a residue of profenofos at 0.3 mg/kg. The MRL is 0.05 mg/kg
 - 1 sample of curry leaves from India contained a residue of profenofos at 0.8 mg/kg. The MRL is 0.05 mg/kg

Risk assessments

One sample of curry leaves contained a residue of triazophos at levels where the effect on health needed to be considered in more detail. The highest level detected was 4 mg/kg in dried curry leaves equivalent to 1.54 mg/kg on a fresh leaf basis using a rehydration factor of 2.6), the highest intake (based on consumption data for parsley) for the critical consumer vegetarians, is 184.5 % of the ARfD of 0.001 mg/kg bw/day for triazophos (JMPR, 2002). The HSE assessment indicated that an effect on health was unlikely. Full details of the risk assessment can be found on page 112. The same sample contained 7 other organophosphorus and carbamate pesticides which required a combined risk assessment

Combined risk assessments

Some samples contained residues of more than one pesticide. Some of these residues are from pesticides which belong to similar chemical groups and may have similar toxicological effects. So, the risk assessors needed to consider their possible impacts on human health, both on their own and in combination.

HSE carried out a combined risk assessment of the relevant samples. One sample contained triazophos (as detailed above) plus 7 other organophosphorus and carbamate pesticides. The triazophos component represented the most significant contribution to the exceedance of the ARfD. The presence of the other 7 pesticides contributed minimally to the triazophos exposure already discussed in the detailed risk assessment section, so the conclusion for the triazophos assessment is still considered valid for the combined assessment. This combined acute risk assessment can be found on page 112. Follow up actions

HSE informed the FSA about the sample of curry leaves containing triazophos at 4 mg/kg and 7 other organophosphorus or carbamate pesticides.

Samples of curry leaves from India are subject to increased border control inspection, so all non-compliant samples were notified to the FSA.

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

Organic sample with a residue

The Secretariat has written to the supplier of the sample of organic curry leaves from Sri Lanka with a residue of profenofos which is not permitted in organic food production. Defra's Organic Farming branch and the organic certification organisation were also informed.

RASFFs issued

HSE sent a draft notification for the following samples to FSA to raise through the EC's Rapid Alert System for Food and Feed (RASFF) (see glossary for more details).

- 1 sample from India containing triazophos at 0.4 mg/kg
- 1 sample from India containing triazophos at 1.6 mg/kg

Eggs

Summary of results

In a survey of 25 samples of eggs collected between April and June 2018 none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

No residues were detected at or above the reporting limit

Survey design

The egg samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

25 samples were tested for up to 104 pesticide residues

Duck

- 1 sample came from the UK

Hens

- 24 samples came from the UK

Pesticide residues detected from those sought

- 25 samples contained no residues from those sought
- None of the samples contained residues above the reporting level
- 5 samples were labelled as organic. None contained residues from those sought

Risk assessments

The laboratory did not detect any residues, so we did not carry out a risk assessment.

Fish (white)

Summary of results

In a survey of 30 samples of fish collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

The EU has not at present set MRLs for any pesticide residues in fish. This means it is legal to trade in fish with pesticide residues at any level, so long as no effects on health are expected.

DDT

Two samples of sea bass contained residues of DDT. The use of DDT is banned or heavily restricted in many countries because the residues take a long time to breakdown in the environment and can accumulate in fatty tissue.

An interpretation of the analytical results shows that the DDT residue found was in the form of DDE which indicates historical use. More detailed information about DDT residues is on page 115 of this report.

BAC (sum) and DDAC

5 samples of various fish contained residues of disinfectants used on surfaces and equipment to ensure microbiological food safety at levels above the MRL for foods other than fish. The level of 0.1 mg/kg is not set as a safety level, so these findings are not in themselves a safety risk. We have asked the suppliers or brand owners for comments.

Survey design

The fish samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

30 samples were tested for up to 38 pesticide residues

Basa

- 1 sample was imported from outside the EU

Cod

- 15 samples were imported from outside the EU
- 1 sample came from the EU

Haddock

- 3 samples were imported from outside the EU

Hake

- 2 samples were imported from outside the EU

Plaice

- 2 samples were imported from outside the EU

Pollock

- 3 samples were imported from outside the EU

Sea bass

- 1 sample came from the UK
- 2 samples were imported from outside the EU

The country of origin on the packaging does not necessarily indicate where the fish was caught. It could be where it was landed or processed or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 24 samples contained no residues from those sought
- 6 samples contained residues above the reporting level
- None of the samples were labelled as organic.

Multiple residues

None of the samples contained residues of more than one pesticide

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Game

Summary of results

In a survey of 19 samples of game collected between April and June 2018, 1 sample contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The game samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

19 samples were tested for up to 38 pesticide residues

Duck

- 10 samples came from the UK
- 3 samples came from the EU

Guinea fowl

- 3 samples came from the EU

Partridge

- 1 sample came from the UK

Venison

- 1 sample came from the UK
- 1 sample was imported from outside the EU

The country of origin on the packaging does not necessarily indicate where the animal was from. It may be where the meat was processed or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 18 samples contained no residues from those sought

- 1 sample contained residues above the reporting level
- 1 sample contained a residue above the MRL
- None of the samples were labelled as organic.

Multiple residues

None of the samples contained residues of more than one pesticide

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in game

- 1 sample of venison from UK contained a residue of BAC (sum) at 0.8 mg/kg. The MRL is 0.1* mg/kg.

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Grapefruit

Summary of results

In a survey of 24 samples of grapefruits collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year grapefruit is being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

Several samples contained residues that required a detailed risk assessment. We have presented the risk assessments in full. Based on HSE's risk assessment of the residues detected, we consider an effect on health is not expected.

Survey design

The grapefruit samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 366 pesticide residues

- 21 samples were imported from outside the EU
- 3 samples came from the EU

Pesticide residues detected from those sought

- All samples contained residues
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

24 samples contained residues of more than one pesticide

- 11 samples contained 2 residues
- 4 samples contained 3 residues
- 3 samples contained 5 residues

- 1 sample contained 6 residues
- 3 samples contained 7 residues
- 1 sample contained 8 residues

Risk assessments

Chlorpyrifos

1 sample of grapefruit contained a residue of chlorpyrifos at a level where the effect on health needed to be considered in more detail. The level detected was 0.1 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is not expected. Full risk assessment is available at page 100.

Imazalil

12 samples of grapefruit contained a residue of imazalil at levels where the effect on health needed to be considered in more detail. The highest level detected was 2.7 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is unlikely when it is assumed that all of the peel is consumed. If the peel is not consumed an effect on health is not expected. Full risk assessment is available at page 101.

Thiabendazole

3 samples of grapefruit contained a residue of thiabendazole at levels where the effect on health needed to be considered in more detail. The highest level detected was 2.1 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is unlikely when it is assumed that all of the peel is consumed. If the peel is not consumed an effect on health is not expected. Full risk assessment is available at page 102.

Prochloraz

One sample of grapefruit contained a residue of prochloraz at levels where the effect on health needed to be considered in more detail. The level detected was 0.5mg/kg mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is unlikely when it is assumed that all of the peel is consumed. If the peel is not consumed an effect on health is not expected. Full risk assessment is available at page 103.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Grapes

Summary of results

In a survey of 37 samples of grapes collected between April and June 2018, 1 sample contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The grape samples were collected by the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesale markets, retail depots, ports and import points).

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

37 samples were tested for up to 371 pesticide residues

- 37 samples were imported from outside the EU

Pesticide residues detected from those sought

- All samples contained residues
- 1 sample contained a residue above the MRL
- None of the samples were labelled as organic.

Multiple residues

34 samples contained residues of more than one pesticide

- 7 samples contained 2 residues
- 9 samples contained 3 residues
- 8 samples contained 4 residues
- 2 samples contained 5 residues
- 1 sample contained 6 residues
- 2 samples contained 7 residues

- 2 samples contained 8 residues
- 2 samples contained 9 residues
- 1 sample contained 15 residues

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in grapes

- 1 sample of grapes from Chile contained a residue of captan (sum) at 0.05 mg/kg. The MRL is 0.03* mg/kg.

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any response received is available on page [87](#).

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Lettuce

Summary of results

In a survey of 15 samples of lettuces collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The lettuce samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

15 samples were tested for up to 366 pesticide residues

Iceberg

- 3 samples came from the UK
- 3 samples came from the EU

Little Gem

- 3 samples came from the UK

Other

- 1 sample came from the UK

Romaine

- 2 samples came from the UK

Round

- 3 samples came from the UK

Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 6 samples contained residues above the reporting level

- None of the samples contained residues above the MRL
- 2 samples were labelled as organic. Neither contained residues from those sought

Multiple residues

4 samples contained residues of more than one pesticide

- 1 sample contained 2 residues
- 1 sample contained 4 residues
- 1 sample contained 5 residues
- 1 sample contained 8 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Melon

Summary of results

In a survey of 30 samples of melons collected between April and June 2018, 3 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year melons are being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

Chlorate

We found chlorate over the default MRL in two of the samples. However, we do not think that these finding should be treated as breaches of the legislation, and we have not highlighted them as such in the brand name annex.

We expected to find residues of chlorate above the MRL in food with a high-water content or that has been washed with water. We are testing a limited number of foods for chlorate in 2018, as we did in 2017, to provide evidence on consumer safety and confirm that it is necessary to review the existing default MRL in order to take account of non-pesticide sources. In particular, chlorine-based treatments of drinking and irrigation water as well as chlorine-based surface disinfectants which are widely used to ensure microbiological safety. We agree with HSE and the FSA that the current MRL does not take account of these often-unavoidable sources.

Following the HSE's risk assessment, we do not expect any of the residues we found to have an effect on health. We do not think any change in production practice by the brand-owners or manufacturers is needed in response to these findings.

This adds to a growing body of evidence, from both official monitoring across the EU and from the food and farming industries, about the incidence of chlorate residues in food.

More information on work being done on chlorate in the diet is available in on page 113

Survey design

The melon samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

30 samples were tested for up to 365 pesticide residues

Cantaloupe

- 6 samples were imported from outside the EU

Frozen Watermelon

- 2 samples were imported from outside the EU

Galia

- 1 sample was imported from outside the EU
- 1 sample came from the EU

Honeydew

- 5 samples were imported from outside the EU
- 2 samples came from the EU

Prepared Fresh Watermelon

- 1 sample was imported from outside the EU

Watermelon

- 6 samples were imported from outside the EU
- 6 samples came from the EU

The country of origin on the packaging does not necessarily indicate where the melon was grown. It may be where it was prepared or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 19 samples contained residues above the reporting level
- 3 samples contained residues above the MRL
- 1 sample was labelled as organic. It didn't contain any residues from those sought

Multiple residues

8 samples contained residues of more than one pesticide

- 2 samples contained 2 residues
- 4 samples contained 3 residues
- 2 samples contained 4 residues

Residues measured above the MRL

The laboratory detected 3 residues above the MRL in melon

- 1 sample of cantaloupe from Honduras contained a residue of chlorothalonil at 1.2 mg/kg. The MRL is 1* mg/kg
- 1 sample of frozen watermelon from Guatemala contained a residue of chlorate at 0.04 mg/kg. The MRL is 0.01 mg/kg
- 1 sample of frozen watermelon from Guatemala contained a residue of chlorate at 0.2 mg/kg. The MRL is 0.01 mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Milk

Summary of results

In a survey of 72 samples of milk collected between April and June 2018 none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected have an effect on health

Survey design

The milk samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

72 samples were tested for up to 104 pesticide residues

Cows milk

- 66 samples came from the UK

Goats milk

- 6 samples came from the UK

The country of origin on the packaging does not necessarily indicate where the milk was from. It may be where the milk was processed or where it was packed for consumer purchase.

Pesticide residues detected from those sought

- 69 samples contained no residues from those sought
- 3 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- 22 samples were labelled as organic. None contained residues from those sought

Multiple residues

None of the samples contained residues of more than one pesticide

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Mushrooms

Summary of results

In a survey of 18 samples of cultivated mushrooms collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year mushrooms are being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The mushroom samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

18 samples were tested for up to 368 pesticide residues

Button

- 9 samples came from the UK
- 9 samples came from the EU

Pesticide residues detected from those sought

- 16 samples contained no residues from those sought
- 2 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- 4 samples were labelled as organic. None contained residues from those sought

Multiple residues

None of the samples contained residues of more than one pesticide

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Mushrooms (speciality)

Summary of results

In a survey of 24 samples of speciality mushrooms collected between April and June 2018, 2 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The mushroom samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 369 pesticide residues

Enoki

- 1 sample came from the EU

Oyster

- 2 samples came from the UK

Portobello

- 13 samples came from the UK
- 1 sample came from the EU

Shiitake

- 7 samples came from the UK

Pesticide residues detected from those sought

- 16 samples contained no residues from those sought
- 8 samples contained residues above the reporting level
- 2 samples contained residues above the MRL
- 2 samples were labelled as organic. Neither contained residues from those sought

Multiple residues

1 sample contained residues of more than one pesticide

- 1 sample contained 2 residues

Residues measured above the MRL

The laboratory detected 2 residues above the MRL in speciality mushrooms

- 1 sample of oyster mushrooms from UK contained a residue of chlormequat at 4.6 mg/kg. The MRL is 0.9* mg/kg
- 1 sample of oyster mushrooms from UK contained a residue of chlormequat at 4.5 mg/kg. The MRL is 0.9 mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

Further investigation: Suspected illegal use

We have passed details of 5 samples of speciality mushrooms from the UK that contained residues of chlormequat to HSE:

- 2 samples of shiitake mushrooms both with chlormequat residues of 0.1 mg/kg;
- 1 sample of portobello mushrooms with a chlormequat residue of 0.02 mg/kg; and
- 2 samples of oyster mushrooms with a chlormequat residue of 4.6 mg/kg and 4.5 mg/kg, which are also above the MRL of 0.9 mg/kg

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Chlormequat is not approved for use on mushrooms in UK. We are aware that the MRL was established as a temporary measure to take into account of residues that are incurred from growing mushrooms on cereal straw that could contain chlormequat from legitimate use.

HSE is investigating; brand name details will not be published until the investigations are complete.

Okra

Summary of results

In a survey of 26 samples of okra collected between April and June 2018, 3 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The okra samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

26 samples were tested for up to 366 pesticide residues

Fresh

- 18 samples were imported from outside the EU

Frozen

- 8 samples were imported from outside the EU

Pesticide residues detected from those sought

- 18 samples contained no residues from those sought
- 8 samples contained residues above the reporting level
- 3 samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

4 samples contained residues of more than one pesticide

- 2 samples contained 2 residues
- 1 sample contained 3 residues

- 1 sample contained 4 residues

Residues measured above the MRL

The laboratory detected 6 residues above the MRL in okra

- 1 sample of okra from Jordan contained residues of:
 - indoxacarb at 0.03 mg/kg, the MRL is 0.02* mg/kg
 - thiacloprid at 0.04 mg/kg, the MRL is 0.01* mg/kg
- 1 sample of okra from India contained a residue of diuron at 0.02 mg/kg. The MRL is 0.01* mg/kg
- 1 sample of frozen okra from India contained residues of:
 - amitraz (sum) at 0.07 mg/kg, the MRL is 0.05* mg/kg
 - diafenthiuron at 0.02 mg/kg, the MRL is 0.01* mg/kg
 - nitenpyram at 0.03 mg/kg, the MRL is 0.01*

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

Okra from India is subject to increased border control measure. We have informed the Food Standards Agency about these findings.

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Pears

Summary of results

In a survey of 18 samples of pears collected between April and June 2018, 1 sample contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health

Survey design

The pear samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

18 samples were tested for up to 369 pesticide residues

- 10 samples were imported from outside the EU
- 8 samples came from the EU

Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 9 samples contained residues above the reporting level
- 1 sample contained a residue above the MRL
- 8 samples were labelled as organic. None contained residues from those sought

Multiple residues

9 samples contained residues of more than one pesticide

- 1 sample contained 2 residues
- 2 samples contained 3 residues
- 3 samples contained 4 residues
- 1 sample contained 5 residues

- 2 samples contained 6 residues

Residues measured above the MRL

The laboratory detected 1 residue above the MRL in pears

- 1 sample of conference pears from the Netherlands contained a residue of chlormequat at 0.1 mg/kg. The MRL is 0.07* mg/kg.

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Peas without pods

Summary of results

In a survey of 36 samples of peas without pods collected between April and June 2018, 9 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

We found chlorate over the default MRL in 9 of the samples. However, we do not think that these findings should be treated as breaches of the legislation, and we have not highlighted them as such in the brand name annex.

We expected to find residues of chlorate above the MRL in food that has been processed or has been washed with water. We are testing a limited number of foods for chlorate in 2018, as we did in 2017, to provide evidence on consumer safety and confirm that it is necessary to review the existing default MRL in order to take account of non-pesticide sources. In particular, chlorine-based treatments of drinking and irrigation water as well as chlorine-based surface disinfectants which are widely used to ensure microbiological safety. We agree with HSE and the FSA that the current MRL does not take account of these often-unavoidable sources.

Following the HSE's risk assessment, we do not expect any of the residues we found to have an effect on health. We do not think any change in production practice by the brand-owners or manufacturers is needed in response to these findings.

This adds to a growing body of evidence, from both official monitoring across the EU and from the food and farming industries, about the incidence of chlorate residues in food.

More information on work being done on chlorate in the diet is available on page 113

Survey design

The pea samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

36 samples were tested for up to 363 pesticide residues

Fresh

- 4 samples were imported from outside the EU

Frozen

- 13 samples came from the UK

In Brine

- 15 samples came from the UK
- 2 samples came from the EU

Marrow Fat

- 2 samples came from the UK

The country of origin on the packaging does not necessarily indicate where the peas were grown. It may be where the peas were processed or where they were packed for consumer purchase.

Pesticide residues detected from those sought

- 19 samples contained no residues from those sought
- 17 samples contained residues above the reporting level
- 9 samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

5 samples contained residues of more than one pesticide

- 5 samples contained 2 residues

Residues measured above the MRL

The laboratory detected 9 residues above the MRL in peas without pods

- 1 sample of hand shelled peas from Kenya contained a residue of chlorate at 0.4 mg/kg. The MRL is 0.01* mg/kg
- 2 samples of hand shelled garden peas from Guatemala contained residues of chlorate at 0.02 mg/kg. The MRL is 0.01 mg/kg
- 1 sample of hand shelled garden peas from Guatemala contained a residue of chlorate at 0.03 mg/kg. The MRL is 0.01 mg/kg

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

- 2 samples of garden peas from UK contained residues of chlorate at 0.04 mg/kg. The MRL is 0.01 mg/kg
- 1 sample of garden peas from UK contained a residue of chlorate at 0.2 mg/kg. The MRL is 0.01 mg/kg
- 2 samples of petits pois from France contained residues of chlorate at 0.02 mg/kg. The MRL is 0.01 mg/kg

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page [87](#).

Peppers

Summary of results

In a survey of 19 samples of peppers collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

This year peppers are being surveyed across the EU as part of the EU co-ordinated multi-annual control programme.

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The pepper samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

19 samples were tested for up to 368 pesticide residues

Fresh

- 19 samples came from the EU

Pesticide residues detected from those sought

- 11 samples contained no residues from those sought
- 8 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

7 samples contained residues of more than one pesticide

- 6 samples contained 2 residues
- 1 sample contained 3 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Pineapple

Summary of results

In a survey of 19 samples of pineapples collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The pineapple samples were bought by a market research company from retail outlets across the UK

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

19 samples were tested for up to 366 pesticide residues

Canned

- 4 samples were imported from outside the EU

Fresh

- 2 samples came from the UK
- 7 samples were imported from outside the EU

Frozen

- 1 sample came from the UK
- 5 samples were imported from outside the EU

The country of origin on the packaging does not necessarily indicate where the pineapples were grown. It may be where they were prepared or where they were packed for consumer purchase.

Pesticide residues detected from those sought

- 7 samples contained no residues from those sought
- 12 samples contained residues above the reporting level

- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

5 samples contained residues of more than one pesticide

- 4 samples contained 2 residues
- 1 sample contained 3 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Potatoes

Summary of results

In a survey of 43 samples of potatoes collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

Two samples contained residues of chlorpropham that required a detailed risk assessment. We have presented the risk assessment in full. Based on HSE's risk assessment of the residues detected, we consider an effect on health is unlikely.

We are aware that the renewal of chlorpropham is under consideration at a European level. We are keeping apprised of the situation and any implications for the assessment of risks to consumers from the results of the monitoring programme.

Survey design

The Animal and Plant Health Agency's Plant Health and Seed Inspectors collected the potato samples from a range of points across the supply chain (wholesalers, potato processors, ports and import points).

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

43 samples were tested for up to 366 pesticide residues

Maincrop

- 18 samples came from the UK
- 1 sample came from the EU

New

- 17 samples came from the UK
- 7 samples were imported from outside the EU

Pesticide residues detected from those sought

- 21 samples contained no residues from those sought
- 22 samples contained residues above the reporting level
- None of the samples contained residues above the MRL

- 2 samples were labelled as organic. Neither contained residues from those sought

Multiple residues

7 samples contained residues of more than one pesticide

- 7 samples contained 2 residues

Risk assessments

Two samples of potato contained a residue of chlorpropham at levels where the effect on health needed to be considered in more detail. The highest level detected was 4.6 mg/kg. If infants ate large portions of potatoes containing chlorpropham at 4.6mg/kg, their intake of chlorpropham could be 142% of the EU Acute Reference Dose. This intake is 70 times lower than the single dose given to dogs without any adverse effects. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 70 still enough to make an effect on health unlikely. This estimate assumes that potatoes are eaten unpeeled (for example as jacket potato); much of the residue is expected to be associated with the peel.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Soft citrus

Summary of results

In a survey of 36 samples of soft citrus collected between April and June 2018, 2 samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

Several samples contained residues that required a detailed risk assessment. We have presented the risk assessments in full. Based on HSE's risk assessment of the residues detected, we consider an effect on health is not expected.

One sample contained prochloraz at 5.7 mg/kg. Some people eating large portions of soft citrus (including all of the peel) containing this level may experience gastro intestinal disturbances after eating large portions of soft citrus. We consider the likelihood of an effect to be low and the effects to be minor, short lived and reversible. If all of the peel is not consumed then an effect on health is not expected.

One sample contained thiabendazole at 8.1 mg/kg. Based on HSE's risk assessment of the residues detected we consider that some people might experience loss of appetite after eating large portions of soft citrus including all of the peel. Such effects would be expected to be minor, short-lived and reversible. If none of the peel is consumed then an effect on health is not expected.

Survey design

The soft citrus samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

36 samples were tested for up to 366 pesticide residues

Clementine

- 6 samples were imported from outside the EU
- 1 sample came from the EU

Mandarin

- 14 samples were imported from outside the EU

- 4 samples came from the EU

Satsuma

- 8 samples were imported from outside the EU
- 3 samples came from the EU

Pesticide residues detected from those sought

- All samples contained residues
- 2 samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

35 samples contained residues of more than one pesticide

- 1 sample contained 2 residues
- 2 samples contained 3 residues
- 10 samples contained 4 residues
- 10 samples contained 5 residues
- 7 samples contained 6 residues
- 2 samples contained 7 residues
- 3 samples contained 8 residues

Residues measured above the MRL

The laboratory detected 2 residues above the MRL in soft citrus

- 1 sample of clementines from Spain contained a residue of propiconazole at 6 mg/kg. The MRL is 5* mg/kg
- 1 sample of satsumas from Peru contained a residue of thiabendazole at 8.1 mg/kg. The MRL is 7 mg/kg

* **Maximum Residue Levels set at the LOD (LOD MRL):** These MRLs are set at a default level, i.e. at the limit of determination (LOD) as specified in EC Regulation 396/2005.

Risk assessments

Chlorpyrifos

1 sample of soft citrus contained a residue of chlorpyrifos at a level where the effect on health needed to be considered in more detail. The level detected was 0.3 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is not expected. Full risk assessment is available at page 105.

Imazalil

14 samples of soft citrus contained a residue of imazalil at levels where the effect on health needed to be considered in more detail. The highest level detected was 4.0 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is unlikely when it is assumed that all of the peel is consumed. If the peel is not consumed an effect on health is not expected. Full risk assessment is available at page 109.

Propiconazole

1 sample of soft citrus contained a residue of propiconazole at a level where the effect on health needed to be considered in more detail. The level detected was 6 mg/kg. Based on HSE's risk assessment of the residues detected an effect on health is not expected. Full risk assessment is available at page 109.

Prochloraz

1 sample of clementine contained a residue of prochloraz at levels where the effect on health needed to be considered in more detail. The highest level detected was 5.7 mg/kg. If toddlers ate large portions of soft citrus (including all of the peel) containing prochloraz at 5.7 mg/kg their intake could be 1269 % of the ARfD. This intake is 8 times lower than a dose which caused no observed adverse effect in a 90-day dog study, a multi-generation rat study and 14-day dog study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider this significant reduction in the factor of 100 to a level of 8 undesirable.

In conclusion we consider that some people might experience gastrointestinal disturbance (salivation, soft faeces, vomiting) after eating large portions (97.5th percentile consumption) of soft citrus containing the highest levels found in this report. Such effects would be expected to be minor, short-lived and reversible.

This assessment assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected. Full risk assessment is available at page 104.

Thiabendazole

4 samples of soft citrus contained a residue of thiabendazole at levels where the effect on health needed to be considered in more detail. The highest level detected was 8.1 mg/kg. Based on HSE's risk assessment of the residues detected we consider that some people might experience loss of appetite after eating large portions of soft citrus including all of the peel. Such effects would be expected to be minor, short-lived and reversible. If the peel is not consumed an effect on health is not expected. Full risk assessment is available at page 107.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Letters sent

The secretariat has written to the suppliers of the samples with residues above the MRL. Any responses received are available on page 87.

Speciality vegetables

Summary of results

In a survey of 24 samples of speciality vegetables collected between April and June 2018, none of the samples contained a pesticide residue above the MRL. These results were reviewed by the Expert Committee on Pesticide Residues in Food (PRiF).

Comments by the PRiF

None of the residues detected would be expected to have an effect on health.

Survey design

The speciality vegetable samples were collected by either the Rural Payment Agency's Horticultural Marketing Inspectors from a range of points in the supply chain (wholesalers, retail depots, ports and import points) or they were bought by a market research company from retail outlets across the UK.

Full sample details, including brand name information, pesticides sought and residues found are available in an accessible format at <https://data.gov.uk/dataset/pesticide-residues-in-food>

Samples tested

24 samples were tested for up to 366 pesticide residues

Celeriac

- 7 samples came from the UK
- 1 sample came from the EU

Eddoes

- 3 samples were imported from outside the EU

Jerusalem artichoke

- 1 sample came from the EU

Mooli

- 10 samples came from the EU

Turmeric

- 1 sample was imported from outside the EU

Yam

- 1 sample was imported from outside the EU

Pesticide residues detected from those sought

- 9 samples contained no residues from those sought
- 15 samples contained residues above the reporting level
- None of the samples contained residues above the MRL
- None of the samples were labelled as organic.

Multiple residues

7 samples contained residues of more than one pesticide

- 1 sample contained 2 residues
- 4 samples contained 3 residues
- 2 samples contained 4 residues

Risk assessments

None of the individual residues detected by the laboratory would be expected to have an effect on health.

Combined risk assessments

Some samples contained residues of more than one pesticide. The pesticide residues found are not chemically related to each other and do not have the same toxicological effects. Following the Health and Safety Executive (HSE)'s risk assessment, we do not expect these residues to have an effect on health, either separately or in combination.

Follow up actions

Further investigation: Suspected illegal use

We have passed details of 2 samples of celeriac from the UK that contained a residue of chlorpropham, which is not approved for use on celeriac in the UK to HSE. HSE is investigating; brand name details will not be published until the investigations are complete.

Section 2: Sample details and supplier responses

Sample details

The sample details are published on data.gov.uk as a dataset in ODS format.

About sample information

The following information is available on each sample collected this quarter:

- Date and place of collection
- Description (e.g. 'runner bean', organic milk);
- Country of origin or manufacture;
- Brand name and packer/manufacturer; and
- Residues detected (results shown in green indicate residues above the MRL).
- Where the brand name of a sample is given the produce involved may have been on sale in other retail premises at the same time.

The description and country of origin are taken from labelling on the food or at the point of sale. The country of origin of processed food may not be the country where the unprocessed produce was produced. This is true even of food that has undergone minimal processing, such as meat that has been butchered or frozen vegetables.

Samples with residues above the MRL are in bold, green text.

UK samples with residues of pesticides not approved for use on that food in the UK in bold, blue text.

Organic samples with residues of pesticides not permitted for use in organic food production are in bold text.

Some brand name details have been withheld – these will be published once enquiries are complete.

The Government's 'brand naming' policy

The Government has decided that brand name information should be published as part of the Government food chemical surveillance programme. Brand names have been published for most pesticide residue surveys since 1998. Certain samples are excluded from the release of brand name information. These include samples taken as part of any pesticide residues enforcement programme and those taken as part of surveys to study

individual people/farms. This policy was reviewed in 2000/1, when Ministers agreed to its continuation.

Where we find residues above an MRL or the presence of non-approved pesticides brand owners/retailers/ growers are notified of the result in advance of publication of reports and given four weeks to comment.

Interpreting brand name information

There is no ready definition of what constitutes a brand in all cases. For clearly branded produce like breakfast cereals or biscuits the “brand owner” is shown. In the case of “own brand” goods this may be one of the multiple retailers. For fruit and vegetables, the retailer is generally shown. For meat, milk and most other animal products the retailer is also generally shown. Finally, for all commodities the country of origin is shown where this was displayed either on the produce or in the store.

Our programme takes samples of produce in approximate proportion to the market share of the main retailers. This has been done to ensure we obtain an accurate representation of a sector (e.g. fruit and vegetables).

Individual programmes are not capable of generating statistically valid information on residues in particular crops from particular retailers. This would require the collection of a much larger number of samples: either substantially increasing costs or greatly reducing the range of different foods sampled in any one year. Therefore, results from an individual survey cannot be taken as a fair representation of the residues status of any particular brand.

However, we do collect samples from a variety of outlets in a range of locations, over a period of years. Successive programmes should therefore help generate information on the typical residues profile of particular types of produce and on major trends in the incidence and levels of pesticides. It should be noted that this quarterly report is not intended to give a comprehensive comparison with previous surveys of the same commodities.

A particular issue arises in relation to the country of origin of fruit and vegetables. The origins included in the reports are those recorded either on the produce or in the store. However, it is not uncommon for mixing to occur on shop shelves. We have responded by increasing the proportion of pre-packed goods sampled. However, pre-packed samples are not available for some produce in some stores and it could also introduce bias to surveys if loose produce were not sampled. Loose produce is therefore sampled but the origin of the sample should be interpreted with a degree of caution.

Action taken by HSE

HSE wrote to:

- The suppliers of all samples containing residues above the MRL
- The authorities of the exporting countries of all samples containing residues above the MRL
- The suppliers of UK samples that contained residues that were not approved for that crop.
- The Organics branch of Defra about samples that were labelled as organic and contained residues of pesticides not approved for organic production
- The suppliers and certification organisation of all organic samples containing residues of pesticides not approved for organic production.

Recipients of the letters are given 4 weeks to provide a statement for inclusion in the report. The Expert Committee on Pesticide Residues in Food reviews any replies received.

Supplier responses

Sample numbers 2948/2018 and 3420/2018: Cream with residues of chlorate above the MRL

Response from Asda

“We are concerned that the residue was detected but re-assured that as you state that this does not pose any risk to the consumer.

Our investigation has confirmed the following:

- Our supplier confirmed that chlorate residues are likely to originate from a combination of chlorine based chemicals used on farms to disinfect milk plants and bulk tanks, cleaning chemicals used in manufacture of products and packaging, and through treated mains water.
- We are working with our suppliers to understand the scope and extent of these issues and what options are open to us to resolve.
- We work and continue to understand the source of chlorate contamination.
- Our UK packing operation has confirmed that they do not use any QAC.

I would like to emphasise that ASDA is committed to selling creams that have been produced in line with good manufacturing, good hygiene practice and residue legislation”.

Sample number 2949/2018: Cheese (soft) with residue of chlorate above the MRL

Response from Asda

“We are concerned that the residue was detected but re-assured that as you state that this does not pose any risk to the consumer.

Our investigation has confirmed the following:

- Our supplier confirmed that chlorate residues are likely to originate from a combination of chlorine based chemicals used on farms to disinfect milk plants and bulk tanks, cleaning chemicals used in manufacture of products and packaging, and through treated mains water.
- We are working with our suppliers to understand the scope and extent of these issues and what options are open to us to resolve.
- We work and continue to understand the source of chlorate contamination.

I would like to emphasise that ASDA is committed to selling cheese that have been produced in line with good manufacturing, good hygiene practice and residue legislation”.

Section 3: HSE assessment of risk

The surveillance programme is designed to enable the regulatory authorities to check that:

- specified pesticide MRLs are being respected;
- users of pesticides are complying with conditions of use specified in the authorisation;
- Dietary intakes of residues are within acceptable limits.

This section details how risks from dietary intakes are assessed.

When assessments are carried out

A screening assessment is done for each residue and commodity combination to identify residue levels that would lead to intakes above the relevant reference doses. Further information on this screening approach is available on request from HSE. Detailed assessments are then produced for every case where the actual residue level found could lead to an intake by any group above the reference dose.

Assessing Dietary intakes

Assessing the acceptability of dietary intakes is complicated. Consumer risk assessments are carried out for both short-term (peak) and long-term intakes. These assessments use information on food consumption collected in UK dietary surveys in conjunction with the residue levels we find. Occasionally, additional pesticide specific information on the losses of residues that occur during preparation and/or cooking of food is also used.

How the assessment is carried out

Short-term intakes (also called NESTIs) are calculated using consumption data for high-level consumers, based on single-day consumption values and the highest residue found in a food commodity. The residue found is multiplied by a variability factor to take account of the fact that residues may vary between individual items that make up the sample analysed. The estimated intake is compared to the Acute Reference Dose (ARfD). This is done for ten consumer groups; adults, infants, toddlers, 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, vegetarians, elderly living in residential homes and elderly living in their own homes.

Long-term intakes (NEDI) are also calculated for high-level consumers, but in this case the consumption data are high-level long-term values rather than peak single-day events, and similarly the residue values used reflect long-term average levels rather than occasional high values. Again, these estimates are made for the ten consumer groups. In this case the estimated intake is compared to the Acceptable Daily Intake (ADI). More information on intake assessments is available on HSE's website:

www.pesticides.gov.uk/guidance/industries/pesticides/topics/pesticide-approvals/pesticides-registration/data-requirements-handbook/consumer-intake-assessments-new-intake-calculation-models.

The reference doses (ADI, ARfD) are set by the Advisory Committee on Pesticides (ACP), or agreed within the EC (an increasing proportion of UK pesticide authorisations are now carried out in accordance with harmonised EU processes). However, where neither the UK nor the EC has set a reference dose, levels set by regulatory authorities in other countries may be used. For a small number of pesticides the reference doses used have been determined by HSE. These have not been independently peer-reviewed and should therefore be regarded as provisional. Reference dose values are available on the EU website:

http://ec.europa.eu/sanco_pesticides/public/index.cfm?event=activesubstance.selection.

Although MRLs are not safety levels, an MRL would not be established if the residue concentrations measured in the supervised trials used to support the MRL would give rise to health concerns. In most cases residues present at the MRL result in intakes below the ARfD and the ADI. So even if the MRL is exceeded this does not always lead to an intake above the ARfD or ADI.

In addition, an estimated intake that exceeds the ADI or ARfD does not automatically result in concerns for consumer health, because a protective approach is used in setting the ADI and ARfD. In the unusual circumstance of an intake exceeding the ADI or ARfD, an evaluation of the toxicological data is made, and details of this assessment would be presented.

Most consumer intake assessments are for short-term exposure rather than chronic exposure. This is because in most cases the monitoring data show the majority of samples to contain residues below the reporting limit and so chronic exposure would not present a concern. Long-term risk assessments have been carried out on a case-by-case basis but are not routinely reported. Long-term exposure assessments are done using median residue levels, rather than using the highest residues found. Therefore, long-term risk assessments would only need to be carried out where data indicated a high proportion of samples contained residues above the MRL (this would result in a higher median residue level than that previously assessed when setting the MRL), or where there is no MRL and acute toxicology is not considered relevant for the particular pesticide concerned.

Where intakes exceed a reference dose, it is necessary for the underlying toxicological studies (animal studies) to be considered to enable the significance of such an exceedance to be understood. Toxicological studies are conducted using different doses to determine the nature of any ill health effects as well as the levels at which such effects can be expected to occur.

Toxicological studies are conducted using test animals to identify the highest experimental dose that causes no detectable adverse effects (the NOAEL). Where there is more than one relevant toxicological study, the lowest appropriate NOAEL for the most sensitive adverse effect is typically used. There is some uncertainty in extrapolating between animals and people and it is therefore important to use a 'safety factor' to account for sources of variation. This safety factor is incorporated (by dividing the NOAEL by the safety factor) in deriving a reference dose, either an ADI or an ARfD, to which consumer intakes are compared. A safety factor therefore extrapolates from the animal testing to the general

population. Factors in the order of x100 are commonly used, x 10 for animal to man, and x10 for within human population differences in sensitivity. However, toxicologists may propose different values (e.g. from 5 to 1000) based on scientific reasoning in accordance with study designs and the quality of the data that has been generated from the studies.

In order to ensure exposures to pesticides do not pose unacceptable risk to humans a wide range of investigations are performed. Most of these are performed on experimental animals because the only end-points that can be examined in human volunteers are those involving observation or blood and urine sampling. Human volunteer studies involving pesticides are not generated in current regulatory work. There is debate at the international level as to whether human studies that have been generated should be used for risk assessment purposes. In the EU, the policy is not to use these data in assessments; the JMPR chose to apply judgement in the appropriate use of these data if available. The HSE risk assessments will usually refer to test animal species, such as dog, rat, and rabbit. All toxicological work is undertaken based on principles of minimising animal distress. Where scientifically valid human data are available the risk assessments will refer to these as they reduce the uncertainty in the assessment. Therefore, human data is only referred to in more limited circumstances.

Acute (short term) toxicology is not a concern for all pesticides, as some are not acutely toxic. In terms of the pesticides that have been found in fruit and vegetables through the surveillance programme an acute risk assessment would not be necessary on the following: tecnazene, maleic hydrazide, diphenylamine, furalaxyl, iprodione, kresoxim-methyl, pendimethalin, propargite, propyzamide, quintozone and tolclofos-methyl.

As the surveillance programme monitors residues in all types of food, from raw commodities (e.g. potatoes) to processed (e.g. wine), dried (e.g. dried fruit) and composite foods (e.g. fruit bread), consumer risk assessments are specifically tailored to address processed and mixed food products. MRLs are generally set for raw commodities, although when MRLs are established the assessment of dietary intakes takes into account the potential for residues to remain in processed foods produced from the raw agricultural commodities. MRLs have been set for processed infant foods, and in future may be extended to other processed food products.

Residues are usually reduced during food processing and occasionally may concentrate. The alteration of residues can be considered in consumer risk assessments, for example, in oil seed rape a fat-soluble pesticide may result in higher residues in the oil compared to residues in the raw seed. Consumption data are available for many major processed food items such as boiled potatoes, crisps, fruit juice, sugar, bread, and wine. Where such consumption data are not available, the intake estimates are based on the total consumption of the raw commodity, which would represent the worst-case (for example, breakfast cereals consumption would be based on total cereal products consumption). In the case of composite products a suitable worst-case alternative would be used, for example total bread consumption for fruit bread consumption.

Dithiocarbamate residues

Dithiocarbamate residues are determined as carbon disulphide which is a common product from different dithiocarbamate pesticides; for the risk assessment a precautionary approach is taken: the worst case dithiocarbamate residue is calculated by assuming the residue is derived from ziram ((a molecular weight conversion is applied to estimate the level of residue based on ziram) and this is compared to the ARfD for ziram. Where it can be confirmed that a specific dithiocarbamate was applied the equivalent residue of the specific active substance is estimated and the intake compared to the appropriate reference dose. We only present a detailed risk assessment when either the worst-case assessment of intake (based on ziram) leads to an exceedance of the ziram ARfD and it has not been possible to further identify the dithiocarbamate source of the residues, or, when further refined assessments based on a specific knowledge of the dithiocarbamate pesticide applied in practice still lead to an exceedance of the ARfD for the known dithiocarbamate pesticide.

Probabilistic Modelling

The standard calculations of consumer exposure use realistic consumption data and residue levels. However, they tend to overestimate intakes in most circumstances. This is due to the assumptions used; fruit and vegetables would contain high levels of residue in an individual unit and that these would be consumed by high-level consumers. They do not take into account the possible range of residue levels and consumption distributions that may occur in reality. These possible combinations of residues and consumption levels can be taken into account using modelling/simulation techniques to produce probability distributions of residue intake levels to indicate the range of consumer intakes, presented as a probabilistic assessment of consumer exposure. These techniques are not yet routinely used to estimate dietary intakes of pesticide residues in the EC.

Multiple residues

The risk assessment process is not standing still. We are aware that some consumers are concerned by the 'cocktail effect'- the possible implications of residues of more than one chemical occurring in, say, a single portion of fruit or vegetables or the interaction between mixtures of pesticides and veterinary medicines at residue levels.

Where more than one pesticide residue is found in a sample, we produce a separate table which identifies each sample and what was found (see Appendix D). If more than one organophosphate/carbamate is found we will undertake an additional risk assessment. If the combination of pesticides found is either unusual or gives cause for concern then this will be detailed in the report.

The Food Standards Agency (FSA) asked the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment to assess these concerns. Their report "2Risk Assessment of Mixtures of Pesticides and Veterinary Medicines" was published in 2002. The Committee concluded that the probability of any health hazard from exposures to mixtures is likely to be small. Nonetheless, it identified areas of uncertainty in the risk assessment process and made recommendations for further work. These fell under the broad headings of regulatory, surveillance, research and public information issues. An

action plan to take forward the recommendations has been published on the FSA website at: <http://www.food.gov.uk/safereating/chemsafe/pesticides/pestmixbranch/>. A number of research projects have been commissioned by the FSA to help progress the action plan; details can be found at <http://www.food.gov.uk/multimedia/pdfs/ressurprilistsep07> and <http://www.food.gov.uk/science/research/researchinfo/researchportfolio/>

Scientific methodologies have yet to be developed to deal with mixtures from groups of pesticides identified by the Committee. However, the Advisory Committee on Pesticides (ACP) has developed an approach for the anticholinesterase compounds. They have also recommended an approach for assessing compounds that might have combined toxicity. This includes a consideration of the proportion of the respective reference doses taken up by the predicted exposures to each active substance. If this is only a small proportion (e.g. <50% if there are two components; <33% for 3 etc.) then assuming simple additivity the risks would still be acceptable. However if exposures to each active substance represent a high proportion of the respective reference doses and the total exceeds 100% a more detailed consideration is needed

(www.pesticides.gov.uk/guidance/industries/pesticides/topics/pesticide-approvals/pesticides-registration/data-requirements-handbook/toxicity-assessment-of-combinations-of-2-or-more-compounds-in-a-formulation).

We are keen to ensure our reports reflect consumer concerns. We therefore now regularly assess findings showing multiple residues of organophosphate and carbamate pesticides. Combined assessment is a new development in risk assessment, which is being taken forward at the international level, e.g. the European Food Safety Authority (EFSA) held a colloquium in 2006 and has set-up two working groups to help develop the methodology (<http://www.efsa.europa.eu/en/events/event/colloque061128.htm>; <http://www.efsa.europa.eu/en/supporting/pub/117e.htm>; <http://www.efsa.europa.eu/en/efsajournal/pub/705.htm>; <http://www.efsa.europa.eu/en/efsajournal/pub/1167.htm>). Further advances in risk assessment methodology will be taken into account in developing the approach to multiple risk assessments in the future.

Assessment of Risk to Human Health: Short-term intake estimates

Short-term intake estimates

Screening assessments have been done for all acutely toxic and potentially acutely toxic pesticides to check that predicted intakes are within the ARfD (or ADI, as appropriate, where an ARfD is not available). An acute exposure assessment is not done for pesticides which are not acutely toxic where it has been established that an ARfD is not required. Toxicological endpoints can be found in the DG Sanco EU Pesticides database which is available at http://ec.europa.eu/food/plant/protection/evaluation/database_act_subs_en.htm

The screening assessment uses the internationally agreed approach to short-term (acute) consumer exposure assessment with UK food consumption data as detailed within the UK NESTI model which is available on the PSD website at <http://www.pesticides.gov.uk/approvals.asp?id=1687>.

For the Q2 (2018) assessments, the following approaches have been taken to refine the NESTI according to case-by-case issues and to ensure that appropriate consumption values are used for less frequently consumed commodities where available food consumption data may be limited:

- Data on beans with pods were used for okra and all forms of speciality green beans.
- Data on both blackberries and raspberries were considered for the screening assessment for blackberries and blueberries as there are low numbers of consumers in some groups. Data on raspberry alone were used for adults, toddlers, vegetarians, and elderly in own home, where the numbers are sufficient, with data from both commodities used for the remaining groups. Although there are low numbers of consumers in the infant and 4-6 year old children groups for both commodities, use of these data was considered reasonable after comparison with alternative data.
- Data on broccoli with the usual variability factor of 5 were used for all forms of broccoli (fresh and frozen broccoli florets; for the latter this represents a worst case screen as a variability factor is not needed).
- Data on broccoli with a variability factor of 7 were used for choy sum and pak choy. This was previously considered in 2016 after the move of choy sum MRL to within the broccoli MRL, when it was considered appropriate to use broccoli data with a variability factor of 7, taking account of the unit weights for choy sum and pak choy, rather than 5 (as in the UK model for broccoli). Chinese cabbage data would be the most appropriate however the number of consumers is too low for robust risk assessments. The EU PRIMo data indicate that broccoli is precautionary when compared to EU chinese cabbage data.

- Data on parsley were used for fresh curry leaves. Residues in dried curry leaves were expressed on a fresh product basis to perform the risk assessment using a rehydration factor of 2.6.
- Consumption values were provided by the Food Standards Agency for grapefruit to identify the various contributions of grapefruit in the diet. The consumption values used here (and as in Q1 2018, Q2 2016 and Q4, 2016) cover consumption for fresh grapefruit excluding juice, canned grapefruit and grapefruit consumed in other recipe forms. For the current assessments, the consumer groups covered represent infants (4 to 18 months old), toddlers (1.5 to 3 years), 4-6 year olds, 7-10 year olds, 11-14 year olds, 15-18 year olds, and elderly. These are slightly different to the groups used previously. For the new survey data vegetarians, elderly residential and elderly in their own home are not reported separately. For some of the consumer subgroups, the number of grapefruit consumers in the surveys (from years 2008 to 2012) are very low e.g. infants which is not unexpected. The highest number of young consumers out of all the young consumer groups was three persons for infants (3 out of 2863 infant consumers). These data are used as they represent the best up to date data. Adults and elderly consume fresh grapefruit at a higher frequency (a higher number of consumers) than the other consumer groups, and of these adults is the most critical consumer group.
- Data on melon were used for all forms of melon and watermelon
- For large speciality mushrooms (large portobello mushrooms), data on mushrooms with a variability factor of 7 and a unit weight of 85 g were used. As per ordinary mushrooms, no variability factor was used for the smaller forms of speciality mushrooms (shiitake, oyster and mini portobello types).
- Data on peas without pods, were used for all forms of peas
- For potato/chlorpropham, as per Q1 2018, the default variability factor of 7 was used, from the EFSA Conclusion (EFSA, 2017). This represents an altered approach to the variability factor which was previously at a lower specific level of 3. The default factor of 7 is applied pending further data that could support a lower variability factor. As an additional change, EFSA have concluded not to further refine risk assessments for chlorpropham according to losses over cooking, since a metabolite might form over cooking that would counterbalance this (EFSA, 2017). The risk assessment below also uses this altered approach whereas in previous PRiF assessments some losses due to cooking of potatoes were previously assumed.
- Data on meat (excluding poultry and offal) were used for all forms of beef and game (venison).
- Data on cheese were used for all forms of soft cheese.
- Data on milk were used for all forms of cream and milk.

- Data on fish were used for all forms of white fish

Monocrotophos residues:

Monocrotophos was found in dried curry leaves at a level of 0.01 and 0.02 mg/kg. The highest residue (representing 0.077 mg/kg in fresh curry leaves) gives a highest estimated short term intake (using parsley consumption data) of 0.00008 mg/kg bw/day for 7-10 year old children. Authorisation for use in the EU were withdrawn in 2003 and EU reference values have not been set. The EFSA use JMPR reference values, set in 1995, to assess risks from monocrotophos residues. This intake is less than both the ARfD of 0.002 mg/kg bw/day and ADI of 0.0006 mg/kg bw/day. However, studies in laboratory animals at doses orders of magnitude higher which were toxic to the animals have indicated that monocrotophos can damage genetic material. It is not known if lower doses which are not toxic also have this effect. Monocrotophos did not increase cancer incidence in long term feeding studies in rats or mice or cause dominant lethal mutations in mice and these findings provide some reassurance that any risks from exposure are likely to be small. Nevertheless, because of uncertainty about the potential for genetic damage at low doses, on a precautionary basis any findings of monocrotophos in food are not desirable.

Apple risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Apple	Dithiocarbamates (ziram)	2.2*	0.033	0.22 (infants) 0.16 (toddlers) 0.12 (4-6 year olds) 0.090 (7-10 year olds)	0.08	EU, 2004

Comment on risk assessment

The usual non-specific approach for dithiocarbamates indicated a potential intake above the ARfD for ziram. The intakes for infants, toddler, 4-6 year old children, and 7-10- year old children exceeded the ARfD. The highest intake was for infants.

If infants ate or drunk large portions of apple containing ziram at 2.2 mg/kg their intake would be 269% of the Acute Reference Dose. This intake is 36 times lower than a dose which caused no observed adverse effects in a ten day rat developmental study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the likelihood of an effect on health to be low, given the remaining factor of 36. This is because an adverse effect on health would rely on

- 1) a susceptible individual eating a large quantity of the product which in turn had the highest levels of residue (i.e. 7 times the maximum value found in monitoring); and
- 2) the actual difference in susceptibility between that individual and rats, being higher than the factor we are left with in this situation; and
- 3) the critical NOAEL being close to the actual doses needed to produce an adverse effect in the animals studied.

In conclusion we consider that some people might experience slight loss of appetite after eating or drinking large portions (97.5th percentile consumption) of apple containing the highest levels found in this report, but we consider the likelihood of an effect on health to be low. Such effects would be expected to be minor and reversible.

Beans with pods risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Beans with pods (Hyacinth beans)	Methomyl	0.6	0.0014	0.0030 (infants) 0.0030 (toddlers)	0.0025	EFSA, 2006

Comment on risk assessment

The intakes for toddlers and infants exceeded the ARfD. The highest intake was for infants.

If infants ate large portions of beans with pods containing methomyl at 0.6 mg/kg, their intake of methomyl could be 120 % of the Acute Reference Dose. This intake is 83 times lower than a dose which caused no observed adverse effect in a study on acute neurotoxicity in rats. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 83 still enough to make an effect on health unlikely.

Curry leaves risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Curry leaves	Triazophos	1.54 (4 [£])	0.0010	0.0018 (vegetarian) 0.0017 (7-10 year olds)	0.001	JMPR, 2002

Comment on risk assessment

The intakes for vegetarians and 7-10 year olds exceeded the ARfD. The highest intake was for vegetarians.

If vegetarians ate large portions of curry leaves containing triazophos at 1.54 mg/kg, their intake of triazophos could be 184 % of the Acute Reference Dose. An EU ARfD has not been set as triazophos has not been assessed at an EU level. However, the JMPR (Joint FAO/WHO meetings on pesticides) has recommended an Acute Reference Dose (ARfD) of 0.001 mg/kg bw/d using human data. It allows an appropriate factor to account for possible differences in susceptibility between people. In such cases a factor of 10 might be used but in this case the factor was larger (12.5) as there was additional rounding. The intake for vegetarians is 7 times lower than a dose of 0.0125 mg/kg bw/d, which caused no observed adverse effect in a three week human volunteer study. The ARfD was also based on a repeated dose study rather than a single dose, which on balance is likely to be more conservative. Taking all the above into account we consider the reduced factor of 7 (from 12.5) still sufficient to make any effect on health unlikely.

Grapefruit risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Grapefruit	Chlorpyrifos	0.1	0.0014	0.0055 (infants)	0.005	EU, 2015

Comment on risk assessment

The risk assessments detailed below refer to the EU acute Reference Dose 2015 value but also consider the risks based on the existing JMPR value which was based on data which examined impacts upon humans. HSE accepts that relevant human toxicology data can be used to calculate the possible impacts of residues in food on humans and based on this assessment do not expect an effect on health.

Assessment A using the ARfD set in the EU

Grapefruit flesh after peeling

EU MRL risk assessment usually assumes that grapefruits are peeled before consumption. After peeling only 2% of the residue remains (EFSA, 2017), the highest intake is below 0.005 mg/kg bw/day, and there are no exceedances of the ARfD.

However, assuming that consumers eat all the peel, intakes for infants exceed the acute reference dose of 0.005 mg/kg bw/day.

Whole grapefruit, including all the peel

The intakes for infants exceeded the EU ARfD. If infants ate large portions of grapefruit containing chlorpyrifos at 0.1 mg/kg, their intake of chlorpyrifos could be 109% of the EU Acute Reference Dose. This intake is 91 times lower than a dose which caused no observed adverse effects in a single dose rat study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 91 still enough to make an effect on health unlikely.

Assessment B with reference to the ARfD set by the JMPR

The intakes for infants exceeded the EU ARfD. If infants ate large portions of grapefruit, including all of the peel, containing chlorpyrifos at 0.1 mg/kg, their intake of chlorpyrifos could be 109% of the Acute Reference Dose. However, the EU ARfD was set without taking into account scientifically valid human data. The JMPR (Joint FAO/WHO meetings on pesticides) has recommended a higher Acute Reference Dose (ARfD) of 0.1 mg/kg bw/day using that human data. It allows an appropriate factor (10) to account for possible differences in susceptibility between people. Intakes in all groups are within the JMPR ARfD. Based on this assessment we do not expect an effect on health.

Conclusion

HSE accept that relevant human toxicology data can be used to calculate the possible impacts of residues in food on humans and based on this assessment do not expect an effect on health.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Grapefruit	Imazalil	2.7	0.039	0.15 (infants)	General population 0.1 Pregnant and nursing females 0.05	EFSA, 2007

Comment on risk assessment

Grapefruit flesh after peeling

The EU MRL risk assessment assumes that grapefruits are peeled before consumption. After peeling only 7% of the residue remains (EU, 2010), the highest intake is below 0.05 mg/kg bw/d, and there are no exceedances of either ARfD.

However, assuming that consumers eat all the peel, intakes for infants exceed the acute reference dose of 0.1 mg/kg bw/day (for the general population excluding pregnant and nursing women).

Whole grapefruit, including all the peel

Pregnant and nursing women

The intakes for adults, (elderly) and 11-14 year old children are all below the ARfD of 0.05 mg/kg bw/d for pregnant and nursing females.

General population

The intakes for infants exceed the ARfD of 0.1 mg/kg bw/d for the general population.

If infants ate large portions of grapefruit containing imazalil at 2.7 mg/kg their intake could be 150% of the Acute Reference Dose of 0.1 mg/kg bw/day. This intake is 67 times lower than a dose which caused no observed adverse effects in a rabbit developmental study, used as the basis of the ARfD. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. Also, it is noted that an ARfD based on maternal toxicity in a developmental study with repeated dosing (13 days) might be over-protective for the general population. Based on this assessment we consider the reduced factor of 67 still enough to make an effect on health unlikely.

This estimate assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within both ARfDs and an effect on health is not expected.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Grapefruit	Thiabendazole	2.1	0.030	0.11 (infants)	0.1	EU, 2017

Comment on risk assessment

Grapefruit flesh after peeling

The EU MRL risk assessment assumes that grapefruits are peeled before consumption. After peeling only 2% of the residue remains (EFSA, 2016), the highest intake is below 0.1 mg/kg bw/d, and there are no exceedances of the ARfD.

Whole grapefruit, including all the peel

However, assuming that consumers eat all the peel, intakes for infants exceed the acute reference dose of 0.1 mg/kg bw/day.

If infants ate large portions of grapefruit containing thiabendazole at 2.1 mg/kg, their intake of thiabendazole could be 115% of the EU Acute Reference Dose. This intake is 87 times lower than a dose which caused no observed adverse effects in a developmental study in rats over 11 days. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 87 still enough to make an effect on health unlikely.

This estimate assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Grapefruit	Prochloraz	0.5	0.0072	0.027 (infants)	0.025	EFSA, 2011

Comment on risk assessment

Grapefruit flesh after peeling

The EU MRL risk assessment assumes that grapefruits are peeled before consumption. After peeling only 10% of the residue remains (JMPR, 2004), the highest intake is below 0.025 mg/kg bw/d, and there are no exceedances of the ARfD.

However, assuming that consumers eat all the peel, intakes for infants exceeded the ARfD.

Whole grapefruits, including all the peel

If infants ate large portions of grapefruit containing prochloraz at 0.5 mg/kg, their intake of prochloraz could be 109% of the Acute Reference Dose. This intake is 92 times lower than a dose which caused no observed adverse effect in a 90-day dog study, a multigeneration rat study and 14-day dog study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 92 still enough to make an effect on health unlikely.

This estimate assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected.

Potato risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Potato	Chlorpropham	4.6	0.11	0.71 (infants)	0.5	EU, 2004

Comment on risk assessment

The intakes for infants exceeded the EU ARfD. If infants ate large portions of potatoes containing chlorpropham at 4.6mg/kg, their intake of chlorpropham could be 142% of the EU Acute Reference Dose. This intake is 70 times lower than the single dose given to dogs without any adverse effects. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 70 still enough to make an effect on health unlikely. This estimate assumes that potatoes are eaten unpeeled (for example as jacket potato); much of the residue is expected to be associated with the peel.

Soft citrus risk assessment

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Soft Citrus	Prochloraz	5.7	0.063	0.32 (toddlers) 0.22 (4-6 year olds) 0.16 (7-10 year olds) 0.11 (infants) 0.10 (11-14 year olds) 0.082 (15-18 year olds) 0.077 (vegetarians) 0.074 (elderly – in own home) 0.063 (adults)	0.025	EFSA, 2011

				0.045 (elderly – in residential setting)		
--	--	--	--	--	--	--

Comment on risk assessment

Soft citrus flesh after peeling

EU MRL risk assessment usually assumes that soft citrus are peeled before consumption. The EU MRL risk assessment assumes that soft citrus are peeled before consumption. After peeling only 5% of the residue remains (JMPR, 2004), the highest intake is below 0.025 mg/kg bw/d, and there are no exceedances of the Acute Reference Dose (ARfD).

However, assuming that consumers eat all the peel, intakes for all the consumer groups exceed the ARfD. The highest intake is for toddlers.

Whole soft citrus, including all the peel

If toddlers ate large portions of soft citrus containing prochloraz at 5.7 mg/kg their intake could be 1269 % of the ARfD. This intake is 8 times lower than a dose which caused no observed adverse effect in a 90-day dog study, a multigeneration rat study and 14-day dog study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider this significant reduction in the factor of 100 to a level of 8 undesirable.

In conclusion we consider that some people might experience gastrointestinal disturbance (salivation, soft faeces, vomiting) after eating large portions (97.5th percentile consumption) of soft citrus containing the highest levels found in this report. Such effects would be expected to be minor, short-lived and reversible.

This assessment assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Soft Citrus	Chlorpyrifos	0.3	0.0033	0.017 (toddlers) 0.011 (4-6 year olds)	0.005	EU, 2015

				0.0086 (7-10 year olds) 0.0060 (infants) 0.0054 (11-14 year olds)		
--	--	--	--	---	--	--

Comment on risk assessment

The risk assessments detailed below refer to the EU acute Reference Dose 2015 value but also consider the risks based on the existing JMPR value which was based on data which examined impacts upon humans. HSE accept that relevant human toxicology data can be used to calculate the possible impacts of residues in food on humans and based on this assessment do not expect an effect on health.

Assessment A using the ARfD set in the EU

Soft citrus flesh after peeling

EU MRL risk assessment usually assumes that soft citrus are peeled before consumption. After peeling only 2% of the residue remains (EFSA, 2017), the highest intake is below 0.005 mg/kg bw/d, and there are no exceedances of the ARfD.

However, assuming that consumers eat all the peel, intakes for toddlers, 4-6 year old children, 7-10 year old children, infants and 11-14 year old children exceed the acute reference dose of 0.005 mg/kg bw/day.

Whole soft citrus, including all the peel

The intakes for toddlers, 4-6 year old children, 7-10 year old children, infants and 11-14 year old children exceed the EU acute reference dose of 0.005 mg/kg bw/day. The highest intake was for toddlers.

If toddlers ate large portions of soft citrus containing chlorpyrifos at 0.3 mg/kg, their intake of chlorpyrifos could be 334% of the EU Acute Reference Dose. This intake is 30 times lower than a dose which caused no observed adverse effects in a single dose rat study. The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the likelihood of an effect on health to be low, given the remaining factor of 30.

This is because an adverse effect on health would rely on

- 1) a susceptible individual eating a large quantity of the product which in turn had the highest levels of residue (i.e. 7 times the maximum value found in monitoring); and
- 2) the actual difference in susceptibility between that individual and rats being higher than the factor we are left with in this situation; and
- 3) the critical NOAEL being close to the actual doses needed to produce an adverse effect in the animals studied.

Furthermore, HSE consider that the EU ARfD was derived using a particularly sensitive approach since red blood cell cholinesterase inhibition was used as the end-point. This is a sensitive way to assess adverse effects due to cholinesterase inhibition.

Based on this assessment, we consider that some people might experience salivation, intestinal disturbances or sweating after eating large portions (97.5th percentile consumption) of soft citrus containing the highest levels found in this report, but we consider the likelihood of an effect on health to be low. Such effects would be expected to be minor, short-lived, and reversible.

Assessment B with reference to the ARfD set by the JMPR

The intakes for toddlers, 4-6 year old children, 7-10 year old children, infants and 11-14 year old children exceeded the EU ARfD. The highest intake was for toddlers.

If toddlers ate large portions of soft citrus, including all of the peel, containing chlorpyrifos at 0.3 mg/kg, their intake of chlorpyrifos could be 334% of the Acute Reference Dose. However, the EU ARfD was set without taking into account scientifically valid data from studies using human volunteers. The JMPR (Joint FAO/WHO meetings on pesticides) has recommended a higher Acute Reference Dose (ARfD) of 0.1 mg/kg bw/d using that human data. This value allows an appropriate factor (10) to account for possible differences in susceptibility between people. Intakes in all groups are within the JMPR ARfD. Based on this assessment we do not expect an effect on health.

Conclusion

HSE accept that relevant human toxicology data can be used to calculate the possible impacts of residues in food on humans and based on this assessment do not expect an effect on health.

Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Soft Citrus	Thiabendazole	8.1	0.090	0.45 (toddlers) 0.31 (4-6 year olds)	0.1	EU, 2017

				0.23 (7-10 year olds) 0.16 (infants) 0.15 (11-14 year olds) 0.12 (15-18 year olds) 0.11 (vegetarians) 0.11 (elderly in own home)		
Comment on risk assessment						
Soft citrus flesh after peeling						
<p>The EU MRL risk assessment assumes that soft citrus are peeled before consumption. After peeling only 2% of the residue remains (EFSA, 2016), the highest intake is below 0.1 mg/kg bw/d, and there are no exceedances of the Acute Reference Dose (ARfD).</p> <p>However, assuming that consumers eat all the peel, intakes for toddlers, 4-6 year old children, 7-10 year old children, infants, 11-14 year old children, 15-18 year old children, vegetarians and elderly (living in their own home) exceed the ARfD. The highest intake is for toddlers.</p> <p>Whole soft citrus, including all the peel</p> <p>If toddlers ate large portions of soft citrus containing thiabendazole at 8.1 mg/kg their intake could be 451 % of the ARfD. This intake is 22 times lower than a dose which caused no observed adverse effects in a developmental study in rats over 11 days. The European Food Safety Authority used this study as the basis of the ARfD.</p> <p>Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider this significant reduction in the factor of 100 to 22 undesirable.</p> <p>In conclusion, we consider that some people might experience loss of appetite after eating large portions (97.5th percentile consumption) of soft citrus including all the peel containing the highest levels found in this report. Such effects would be expected to be minor, short-lived, and reversible.</p> <p>This assessment assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected.</p>						
Crop	Pesticide		Intake (mg/kg bw/day)	ARfD		

		Highest residue (mg/kg)	Adult	Critical group [†]	(mg/kg bw/day)	Source
Soft Citrus	Propiconazole	6	0.066	0.33 (toddlers)	0.3	EU, 2003
Comment on risk assessment						
Soft citrus flesh after peeling						
<p>The EU MRL risk assessment assumes that soft citrus are peeled before consumption. After peeling only 1% of the residue remains (EFSA, 2015), the highest intake is below 0.3 mg/kg bw/d, and there are no exceedances of the ARfD.</p> <p>However, assuming that consumers eat all the peel, intakes for toddlers exceed the ARfD.</p>						
Whole soft citrus, including all the peel						
<p>If toddlers ate large portions of soft citrus containing propiconazole at 6 mg/kg, their intake of propiconazole could be 111 % of the EU Acute Reference Dose. This intake is 91 times lower than a dose which caused no observed adverse effects in a developmental study in rats dosed over a ten day period. The European Food Safety Authority used this study as the basis of the ARfD.</p> <p>Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 91 still enough to make an effect on health unlikely.</p> <p>This estimate assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within the ARfD and an effect on health is not expected.</p>						
Crop	Pesticide	Highest residue (mg/kg)	Intake (mg/kg bw/day)		ARfD (mg/kg bw/day)	Source
			Adult	Critical group [†]		
Soft Citrus	Imazalil	4.0	0.044	0.22 (toddlers) 0.15 (4-6 year olds) 0.11 (7-10 year olds) 0.072 (11-14 year olds) 0.057 (15-18 year olds) 0.054 (vegetarians) 0.051 (elderly in own home)	General population 0.1 Pregnant and nursing females 0.05	EFSA, 2007

Comment on risk assessment

Soft citrus flesh after peeling

The EU MRL risk assessment assumes that soft citrus are peeled before consumption. After peeling only 7% of the residue remains (EFSA, 2017), the highest intake is below 0.05 mg/kg bw/d, and there are no exceedances of either ARfD.

Whole soft citrus, including all the peel

However, assuming that consumers eat all the peel, intakes for toddlers, 4-6 year old children and 7-10 year old children exceed the acute reference dose of 0.1 mg/kg bw/day (for the general population excluding pregnant and nursing women). It is noted that in consumer groups aged over 11 years intakes for 11-14 year old children, 15-18 year old children, and vegetarians exceed the acute reference dose of 0.05 mg/kg bw/day (for pregnant and nursing women).

Pregnant and nursing women

The intakes for 11-14 year old children, 15-18 year old children, and vegetarians exceeded the ARfD of 0.05 mg/kg bw/d for pregnant and nursing females. The highest intake was for 11-14 year old children.

If 11-14 year old children ate large portions of soft citrus containing imazalil at 4.0 mg/kg their intake could be 144% of the Acute Reference Dose of 0.05 mg/kg bw/day. This intake is 70 times lower than a dose which caused no observed adverse effects in a 13 day repeat dose rabbit developmental study (the ARfD is based on a NOAEL of 5 mg/kg bw/day for fetal toxicity (increased resorptions). The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account the uncertainties caused by using animal data and possible differences in susceptibility between people. We consider the reduced factor of 70 still enough to make an effect on health unlikely.

General population

The intakes for toddlers, 4-6 year old children, 7-10 year old children and infants exceed the ARfD of 0.1 mg/kg bw/d for the general population. The highest intake was for toddlers.

If toddlers ate large portions of soft citrus containing imazalil at 4.0 mg/kg their intake could be 223% of the Acute Reference Dose of 0.1 mg/kg bw/day. This intake is 45 times lower than a dose which caused no observed adverse effects in a rabbit developmental study, used as the basis of the ARfD (the ARfD is based on a NOAEL of 10 mg/kg bw/day for reduced bodyweight gain and food consumption in dams). The European Food Safety Authority used this study as the basis of the ARfD.

Toxicologists usually apply a factor of 100 to this dose to take into account uncertainties caused by using animal data and possible differences in susceptibility between people. Also, it is noted that an ARfD based on maternal toxicity in a developmental study with repeated dosing (13 days) might be over-protective for the general population. Based on this assessment we consider the reduced factor of 45 still enough to make an effect on health unlikely.

This estimate assumes that peel of the fruit is consumed. However, if the peel is not consumed then the risk assessment that is the basis for the MRL applies (see the first paragraph of this assessment) and intakes in all groups are within both ARfDs and an effect on health is not expected.

[†]Highest intake of all ten consumer groups, or intakes for all consumer groups that exceed the ARfD.

[£]The residue in the dried curry leaves was 4 mg/kg triazophos. Expressed on a fresh leaf product basis, the residue was recalculated as 1.54 mg/kg using a re-hydration factor of 2.6.

Dithiocarbamate residues are determined as carbon disulphide which is a common product from different dithiocarbamate pesticides; for the risk assessment a precautionary approach is taken: the worse case dithiocarbamate residue is calculated by assuming the residue is derived from ziram and this is compared to the ARfD for ziram. Where it can be confirmed that a specific dithiocarbamate was applied the equivalent residue of the specific active substance is estimated and the intake compared to the appropriate reference dose.

* Dithiocarbamate residue calculated as 2.2 mg/kg (using a molecular weight conversion of 2.007 for ziram) based on a carbon disulphide residue of 1.1 mg/kg.

Acute risk assessments for samples containing more than one triazole fungicide, organophosphorus/carbamate, carbendazim/thiophanate-methyl, clothianidin/thiamethoxam or captan/folpet following screening assessment

Curry leaves combined risk assessment

Crop/Critical group	Pesticide	Residue mg/kg [§]	Intake			ARfD	Source
			mg/kg bw	%ARfD			
Curry Leaves [§] (vegetarian)	acephate	0.023	0.000028	0.1	} Total 187.5	0.025	EU, 2007
	chlorpyrifos	0.023	0.000028	0.6		0.005	EU, 2015
	dimethoate	0.0038	0.000005	0.05		0.01	EU, 2014
	ethion	0.19	0.00023	1.5		0.015	ACP, 1999
	methamidophos	0.0077	0.000009	0.3		0.003	EU, 2006
	monocrotophos	0.0077	0.000009	0.5		0.002	JMPR, 1995
	profenofos	0.73	0.00087	0.1		1	JMPR, 2007
	triazophos	1.54	0.00185	184.5		0.001	JMPR, 2002

Comment on risk assessment:

Triazophos represents a within two fold exceedance of its acute reference dose (see the risk assessment in the table above for the detailed risk assessment for triazophos in curry leaves). The presence of the other organophosphorus and carbamate pesticides in the same sample as listed above adds minimally to the triazophos exposure. As such, the conclusion above for triazophos is still considered valid for this combined assessment, and that despite a reduction in the safety factor, an effect on health is unlikely.

[§]The residue levels have been recalculated and expressed on a fresh leaf basis (using a rehydration factor of 2.6) in order to conduct the assessment using fresh parsley consumption data. This sample was a dried curry leaves sample.

Section 4: issues arising in this report and updates on previous reports

Issues arising in this report

Chlorate

Since 2016 we have been testing a limited number of foods for chlorate. The pesticide sodium chlorate is a residual broad action weed killer that is not authorised for use in the EU. However, we are confident that the residues we are detecting come from use of chlorine-based disinfectants used to maintain microbiological safety (control microorganisms that cause food poisoning). Because these residues are unavoidable, and important for the maintaining of microbiological control vital for food safety, we are not treating these results as breaches of the MRL. We are not advising that food companies change their existing practices as a result of our findings but they should be aware about the ongoing discussion in this area.

We are only part of the work going on across government and beyond to consider what to do about chlorate residues in food and water.

The Food Standards Agency is working with the food industry to develop and promote best practice in the use of sanitisers. This is important because the presence of low level residues of chlorate in food results from measures taken by the food and water industries to protect food safety by reducing microbiological contamination of food and drink (including drinking water, which is a significant source of chlorate in food). Chlorate itself is not used as a disinfectant, but chlorine-based sanitisers can contain small amounts of chlorate.

The Health and Safety Executive is leading UK work in the EU to establish more meaningful statutory levels for chlorate in food to provide reassurance to consumers and allow the continued use of disinfectants that are themselves important for safeguarding human health. Chlorate was historically used as a pesticide and residues of chlorate in food fall under EU legislation on plant protection products. Since it is no longer authorised for use as a pesticide, chlorate is subject to a Maximum Residue Limit (MRL) of 0.01 mg/kg. This level was, in line with normal practice for pesticides that are not currently used, set at the default limit of detection rather than on the basis of an assessment of health risks. Our findings are adding to the evidence that that current legal limits are not sufficient to allow for the essential use of disinfectants to protect food and water hygiene. We have asked to be kept informed on developments on setting appropriate MRLs, in particular when any formal proposals are made.

Defra is also working on the EU recast of its Drinking Water Directive. Discussions are underway about the possible future monitoring of chlorate and the level to be achieved. In national legislation throughout the UK it is already a requirement to keep disinfection by-

products as low as possible. This is usually achieved through management of disinfectant dosing and storage.

Departments have an approach to enforcement, which reflects an agreement within the EU that, while the default MRL for chlorate remains in place, enforcement should be left to the discretion of Member States. The UK approach, in line with that normally taken for environmental or process contaminants, is to require that levels in food are as low as reasonably achievable to ensure the protection of human health.

Beginning in 2018 the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) will be considering chlorate as part of its on-going work looking at the chemicals in the diet of infants and young children (up to 5 years). The European Food Safety Authority's 2015 opinion on chlorate¹ establishes appropriate health-based guidance values for chlorate exposure to protect against acute and chronic risks to health.

Residues below the MRL that exceed the ARfD

When MRLs are agreed at the EU level they are set at levels that are compatible with consumer safety. Occasionally, assessment of PRiF monitoring samples containing residues below or at the MRL will show consumer intakes could potentially be above the ARfD. This situation typically arises because of one of three reasons:

- the ARfD may have been lowered because of new information but there is a delay before MRLs have been reassessed or new MRLs are put in place;
- during the MRLs process the risk assessments are currently based on the highest residue level observed in residues trials used to support the MRL which will often be less than the actual MRL (it is expected that most residues found will be below the MRL, and if for this reason there are later samples which give intakes above the ARfD the numbers are expected to be low);
- the agreed EU approach might assume the commodity is peeled and data are used to reduce the intake in the risk assessment at the time of setting MRLs, whereas in the PRiF work risk assessments for the whole commodity are presented as routine and, if information showing the effects of processing on residues level is available to PRiF, a refined assessment is presented.

The first two of these reasons are common to EU assessments and the third represents a difference between the approach used by HSE for the risk assessment and that used at the time the MRL is set. We will highlight how our assessments differ from that done at the EU level so that readers are aware of the basis of the evaluation.

¹ [EFSA Journal 2015;13\(6\):4135 \[103 pp.\]](http://efsa.europa.eu/food/plant/standing_committees/sc_phytopharmaceuticals/index_en.htm)
http://ec.europa.eu/food/plant/standing_committees/sc_phytopharmaceuticals/index_en.htm

DDT

The use of DDT is banned or heavily restricted in many countries. It isn't allowed for use on food crops any more but it is still used in some countries outside the EU as a public health insecticide. Residues of DDT take a long time to break down in the environment and can accumulate in fatty tissue which is a major reason that it has been banned in the EU and many other countries.

Due to the bans and restrictions on use, the levels in food have decreased substantially since the 1960s and 1970s. Even so, because it takes a long time to breakdown we do expect, and do see, occasional DDT residues in our monitoring results. Overall, the incidence and the size of residues have fallen steadily over time, which is what we would expect. In recent years none of our findings were unusual, unexpected or of concern. We can tell from the chemical form that we detect whether the residues we have found are from historic use (which is what we usually find). We explain this every time we publish DDT results to try to make it as clear as we can that the results show food producers are not using DDT today. However, there are occasional media stories about DDT and various links and associations, which do not make this distinction.

The residues we find nowadays are at levels that would not be expected to have any effect on health, either in the short term or in the long term, when checked against today's understanding of the effect of DDT on health. As a committee, we take care to ensure we look thoroughly at this, and the Food Standards Agency is also actively involved in our considerations.

Follow-up from Previous Reports

Quarter 3 2017

Kale

Linuron: Sample numbers 2101/2017, 2108/2017, 5310/2017, 2471/2017 and 3653/2017

We passed details of five samples of kale from the UK that contained residues of linuron to HSE. HSE's enquiries are not yet complete; an update will appear in a future report.

Raspberries

Chlorpyrifos: Sample number 1855/2017

We passed details of a sample of raspberries from the UK that contained a residue of chlorpyrifos to HSE. HSE's enquiries are not yet complete; an update will appear in a future report.

Quarter 4 2017

Cauliflower

Triallate: Sample number 3698/2017

We passed details of a cauliflower from the UK that contained triallate to HSE. HSE's enquiries are not yet complete; an update will appear in a future report.

Cucumber

Propamocarb (sum): Sample number 3334/2017

We passed details of a cucumber from the UK that contained propamocarb to HSE. HSE's enquiries are not yet complete; an update will appear in a future report.

Quarter 1 2018

Apples

Myclobutanil: Sample number 2973/2018 and 2833/2018

We passed details of two samples of apples from the UK that contained myclobutanil to HSE. Myclobutanil was authorised for use on apples until 30 September 2017. HSE have found no evidence of use after that date, and not been able to rule out that these residues came from authorised treatments before that date

Lettuce

Inorganic Bromide: Sample number 4787/2018

We passed details of a lettuce sample from the UK that contained inorganic bromide to HSE. HSE's enquiries are not yet complete; any update will appear in a future report.

Brand name details of samples where follow-up action is now complete

Sample ID	Date of Sampling	Description	Country of Origin	Retail Outlet	Address	Brand Name	Packer / Manufacturer	Pesticide residues found in mg/kg (MRL)
2973/2018	05/02/2018	Royal Gala	UK	Aldi	13 Adams Walk, Kingston Upon Thames KT1 1DF	Aldi	Aldi Stores Ltd PO Box 26 Atherstone, Warwickshire CV9 2SQ	boscalid 0.2 (MRL = 2) flonicamid (sum) 0.04 (MRL = 0.3) myclobutanil 0.05 (MRL = 0.6) pyraclostrobin 0.05 (MRL = 0.5)
	05/02/2018	Cox	UK	Morrisons		Morrisons		boscalid 0.09 (MRL = 2)

2833/2018					1 Roman Way, Malvern, Worcester WR14 1PZ		W M Morrisons Supermarkets Plc 71 Gain Lane, Bradford, West Yorkshire BD3 7DL	captan (sum) 0.09 (MRL = 10)
								flonicamid (sum) 0.01 (MRL = 0.3)
								fludioxonil 0.02 (MRL = 5)
								myclobutanil 0.05 (MRL = 0.6)
								pyraclostrobin 0.03 (MRL = 0.5)

In our next report:

In Quarter 3 of 2018 we will look at results for:

- Apples
- Aubergines
- Bananas
- Beans with pods
- Beef
- Berries
- Bread
- Broccoli
- Cheese (soft)
- Chinese cabbage
- Eggs
- Fish (white)
- Frozen fruits and smoothie mixes
- Game
- Ginger
- Grapefruit
- Grapes
- Infant food (cereal based)
- Lentils
- Melon
- Milk
- Mushrooms
- Okra
- Pears
- Peppers
- Pineapple
- Potatoes
- Soft citrus
- Vine leaves

Section 5: background and reference

Reasons for pesticide residue testing

Food safety is important. Modern food production processes have given us plentiful supplies of a wide range of good quality affordable produce.

In the food industry of today the production environment can be managed from the preparation of seeds used for crops, through to growth, harvesting and storage of the produce.

One of the ways the food industry controls the environment in which foodstuffs are produced is by applying pesticides. They help farmers and growers maximise the production of food stuffs by, for example, preventing weeds inhibiting the growth of the crop, or insects destroying or infesting them. Pesticides can also be used to help protect seeds or prolong the life of crops after they have been harvested. Biological and physical (cultural) controls are also used to protect crops or as part of an integrated system.

As pesticides are used to control unwanted pests, weeds and diseases, they can potentially also harm people, wildlife and the environment. This is why the UK, in common with most other countries, imposes legally enforceable conditions as to how and when pesticides can be used. No pesticide can be supplied or used on a food or ornamental crops in the UK without Government authorisation. To obtain this authorisation the manufacturer of the pesticide must show that it does not present a concern for people's health or the environment. Naturally derived and synthetic pesticides are subject to the same regulation.

Once the authorisation has been granted Government authorities carry out follow up checks to ensure that the authorisation is providing the necessary degree of protection to users, consumers and the environment and that those who use pesticides are complying with conditions specified within it.

The Government authority responsible for checking pesticide residues in foodstuffs is the Health and Safety Executive. Defra's Expert Committee on Pesticide Residues in Food (PRiF) oversees and provides an independent check on this work. We know that the use of pesticides on crops may lead to traces (residues) of these chemicals in food and we expect to find these in our monitoring programme.

Defra's Expert Committee on Pesticide Residues in Food (PRiF)

The Expert Committee on Pesticide Residues in Food was established in 2011. Our members have a broad range of expertise relating to the food supply industry. The main function of the Committee is to oversee Government's £2 million pesticide residues surveillance programme. Previously this work was carried out by the Pesticide Residues Committee.

Our Chairman, Dr Paul Brantom is an independent consultant in toxicological risk assessment. The Committee also includes members with expertise in toxicology, food production and supply as well as two public interest experts.

Information on the membership of the PRiF is also available on the PRiF's website:

<https://www.gov.uk/government/groups/expert-committee-on-pesticide-residues-in-food-prif>

Our role is to advise Ministers, the Director of the Health and Safety Executive (HSE) and the Chief Executive of the Food Standards Agency (FSA) on:

- the planning of surveillance programmes for pesticide residues in the UK food supply and the evaluation of the results;
- Procedures for sampling, sample processing, new methods of analysis, the assessment of variability of pesticide residues in food and related issues.

Detail of reporting practice

Results by food commodity

We include information about the survey (for instance where samples came from) for each commodity

Detailed tabulated results are at the back of this report - these tables are also available for download from our website

We summarise our findings and any follow-up action taken.

Risk assessments – single residues

All results are screened by HSE to check for intakes above the Acute Reference Dose (ARfD). HSE assumes a relatively high level of intake and also assumes that most produce is eaten whole including peel/skin even when these are rarely consumed

Where intakes above the ARfD are identified, we consider a detailed risk assessment prepared by HSE (at Section II of this report).

Our observations and the follow-up action taken are summarised in the section for that food.

Risk assessments – multiple combined residues

- Residues of more than one pesticide from the same category/class of particular categories of pesticides, which have a similar toxicological mode of action, are screened by HSE to check for intakes above the combined Acute Reference Dose (ARfD).

- Where combined intakes above the combined ARfD are identified, we consider a detailed combined risk assessment prepared by HSE (at Section II of this report).
- Our observations and any follow-up action taken are summarised in the section for that food commodity.

Risk assessment - conclusions

Where, in the light of current knowledge and considering the usual level of scientific uncertainty (or precaution) the intake will not cause ill health the conclusion will say no effect on health is expected.

Where, in the light of current knowledge and considering a slightly higher level of scientific uncertainty (or less precaution) the intake is not likely to cause ill health, the conclusion will be less definite and state that an effect on health is unlikely.

Where scientific uncertainty is greater more information is provided.

Residues in UK produce of pesticides which are not approved for use on that crop in the UK.

- All residues found in UK-produced foods are checked by HSE to make sure the pesticide is approved for use.
- Where no UK approval is identified, details of the sample are referred to HSE's Enforcement Section for follow up.
- Our observations and any follow-up action taken to date are summarised in the section for that food commodity. We may have to withhold details of samples while investigations are underway, in which case the details will be published in a later report.

Residues above the MRL, after taking into account measurement uncertainty

- Samples containing residues above the MRL are listed at Appendix B, and those which are clearly above the MRL after taking into account measurement uncertainty of plus or minus 50% are highlighted.
- Our observations and any follow-up action taken are summarised in the section for that food commodity.

The results in our reports are rounded for publication but not adjusted for measurement uncertainty.

We apply measurement uncertainty only to decide whether to highlight a result as over the MRL in the brand name annex. To do this we use the actual value reported by the laboratory before rounding. If after taking measurement uncertainty into account that value is found to be over the MRL the result will be highlighted in the brand name annex.

For example:

- The lab reports the results of duplicate analysis of a residue above an MRL at 0.023 mg/kg and 0.025 mg/kg giving an average value of 0.024mg/kg. For reporting purpose this value would be 0.02 mg/kg.
- If measurement uncertainty is then applied to the reported value of 0.02 mg/kg it could take the value to between 0.01 - 0.03 mg/kg. If the MRL is 0.01 mg/kg the lower value would be at the MRL and there is no exceedance.
- However, if measurement uncertainty is applied to the measured result, e.g. 0.024 mg/kg the value could then be in the range of 0.012 – 0.036 mg/kg. In this case the lower value is above the MRL and so will be treated as an exceedance.

Residues in organic food

- We monitor pesticide residues in all the UK food supply, including organic food.
- We are not responsible for checking compliance with the rules associated with organic production. However, when we do detect residues in an organic food we explain whether or not those residues indicate a breach of the rules and inform Defra's Organic Farming Branch.

Brand Name Annex

- Full brand name details for samples included in this report are published in a brand name annex. Within this annex, samples with results of interest are highlighted.
- Brand name details are only published when enough follow-up work is completed for us to be reasonably sure whether a breach of the law or good practice has occurred. Therefore, sometimes brand name details are withheld pending completion of this work but are published in a later report.

Pesticides analysed as multi-component analytes and their reporting limits

Why some results cover more than one substance

Both the legal controls and our analytical tests are aimed at checking food for the presence of residues of specific pesticides. Residues are the chemical traces left behind after pesticides are used. In most cases the residue of a pesticide is measured by first identifying the pesticide and then measuring the quantity of that pesticide in the food itself. But for some pesticides the residue remaining in the food is known to be chemically different from the original pesticide and so the laboratory needs to look for more than one component. There are various reasons why this happens, for example:

- the animal or plant can change the pesticide into related chemicals
- the pesticide can change in the environment into related chemicals
- some pesticides are mixtures of chemicals, so the relevant components of the mixture need to be checked for
- in the laboratory sample preparation and/or analysis may change pesticides into related chemicals
- related chemicals may be pesticides in their own right

The MRL setting process takes account of all these issues. The EU may set a complex residue definition to ensure that the identity and quantity of the residue found is representative of the pesticide present. A complex residue definition may be set where it is necessary for safety reasons or to be able to accurately identify the pesticide residue present in the food. This definition usually includes the actual pesticide, plus other related chemicals. These residues are usually reported together as a “sum”. Sometimes different foods need different definitions because different pesticide residues are known to occur in that food. For instances, plants and animals may metabolise a pesticide differently, which forms different residues.

The full definitions of pesticides that we have found in our surveys are described in the table below. If you would like more detail about a particular residue definition, please get in touch. You can email us at prif@hse.gov.uk and other contact details are on the back cover.

Where the detailed individual analysis results tell us something useful, we mention that in our conclusions.

How we calculate sums

Unless the definition says otherwise, the summed result is a simple addition. For individual components that are not detected that result is treated as a zero.

Where a residue definition says, “expressed as”, that means that the individual component results are adjusted by molecular weight before being added together. The residue definition is set this way so that the final calculated result for the whole definition is an expression of the level

of the most toxic component, and so that value can be used directly in consumer risk assessment without further adjustment.

The EU Reference Laboratories for pesticide residues have an e-learning package aimed at analytical chemists on this very technical subject at <http://www.eupt.es/e-learning/>.

Complex residue definitions used in our reports

There are a large number of pesticides used and types of food in the world. So other complex residue definitions may apply to food/pesticide combinations not yet considered by PRiF. You can look up all the EU MRL definitions for pesticide residues at the European Commission's pesticide database at http://ec.europa.eu/food/plant/pesticides/pesticides_database/index_en.htm

Short name we use in our reports	Legal residue definition – These definitions apply to all foods unless otherwise stated
2,4-D (sum)	2,4-D (sum of 2,4-D and its esters expressed as 2,4-D)
abamectin (sum)	Abamectin (sum of Avermectin B1a, AvermectinB1b and delta-8,9 isomer of Avermectin B1a)
aldicarb (sum)	Aldicarb (sum of Aldicarb, its sulfoxide and its sulfone, expressed as Aldicarb)
aldrin and dieldrin	Aldrin and Dieldrin (Aldrin and dieldrin combined expressed as dieldrin), aka dieldrin (sum)
Amitraz	Amitraz (amitraz including the metabolites containing the 2,4 - dimethylaniline moiety expressed as amitraz)
BAC (sum)	Benzalkonium chloride (mixture of alkylbenzyltrimethylammonium chlorides with alkyl chain lengths of C ₈ , C ₁₀ , C ₁₂ , C ₁₄ , C ₁₆ and C ₁₈)
benthiavalicarb (sum)	Benthiavalicarb (Benthiavalicarb-isopropyl (KIF-230 R-L) and its enantiomer (KIF-230 S-D) and diastereomers (KIF-230 R-L and KIF-230 S-D))
bixan (animal products)	Sum of bixafen and desmethyl bixafen expressed as bixafen This definition applies to animal products only
captan and folpet	Sum of captan and folpet aka captan/folpet This definition applies only to pome fruit (fruits such as apples and pears), strawberries, raspberries, currants, tomatoes and beans. For all other foods there are separate MRLs for captan only and for folpet only.
carbendazim (animal products)	Carbendazim and thiophanate-methyl, expressed as carbendazim

Carbendazim (sum)	Carbendazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim)
carbofuran (sum)	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)
chlordane (animal products)	Chlordane (sum of cis- and trans-isomers and oxychlordane expressed as chlordane) This definition applies to animal products only
chlordane (sum)	Chlordane (sum of cis- and trans- isomers) This definition applies to all foods except animal products
chlorpropham (potatoes)	Chlorpropham only This definition applies only to potatoes
chlorpropham (sum for animal products)	Chlorpropham and 4-hydroxychlorpropham-O-sulphonic acid (4-HSA), expressed as chlorpropham This definition applies only to animal products
chlorpropham (sum)	Chlorpropham (Chlorpropham and 3-chloroaniline, expressed as Chlorpropham) This definition applies to all foods except potatoes and animal products
DDAC (sum)	Didecyldimethylammonium chloride (mixture of alkyl-quaternary ammonium salts with alkyl chain lengths of C ₈ , C ₁₀ and C ₁₂)
DDT (sum)	DDT (sum of p,p'-DDT, o,p'-DDT, p-p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Dichlorprop	Sum of Dichlorprop, including dichlorprop-p and its conjugates, expressed as dichlorprop
dicofol (sum)	Dicofol (sum of p, p' and o,p' isomers)
Dimethenamid	Dimethenamid-p (Dimethenamid-p including other mixtures of constituent isomers (sum of isomers))

dimethoate (sum)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
disulfoton (sum)	Disulfoton (sum of disulfoton, disulfoton sulfoxide and disulfoton sulfone expressed as disulfoton)
dithiocarbamates	Dithiocarbamates are a group of pesticides that are chemically similar. Testing for them individually in routine analysis is not possible, so MRLs are set for a test for the group.
endosulfan (sum)	Endosulfan (sum of alpha- and beta-isomers and endosulfan-sulphate expressed as endosulfan)
fenamiphos (sum)	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
fenchlorphos (sum)	Fenchlorphos (sum of fenchlorphos and fenchlorphos oxon expressed as fenchlorphos)
fensulfothion (sum)	Fensulfothion (sum of fensulfothion, its oxygen analogue and their sulfones, expressed as fensulfothion).
fenthion (sum)	Fenthion (fenthion and its oxygen analogue, their sulfoxides and sulfone expressed as parent)
fenvalerate & esfenvalerate (all isomers)	Fenvalerate (any ratio of constituent isomers (RR, SS, RS & SR) including esfenvalerate)
fipronil (infant food)	Sum of fipronil and fipronil-desulfinyl, expressed as fipronil This definition applies to foods for babies only
fipronil (sum)	Fipronil (sum Fipronil and sulfone metabolite (MB46136) expressed as Fipronil) This definition applies to all foods except foods for babies
flonicamid (sum)	Flonicamid (sum of flonicamid, TNFG and TNFA) This definition applies to all food except animal products
fluazifop-p-butyl (sum)	Fluazifop-P-butyl (fluazifop acid (free and conjugate))

haloxyfop (sum)	Haloxyfop including haloxyfop-R (Haloxyfop-R methyl ester, haloxyfop-R and conjugates of haloxyfop-R expressed as haloxyfop-R)
Heptachlor (infant food)	Sum of heptachlor and trans heptachlor epoxide This definition applies to foods for babies only
Heptachlor (sum)	Heptachlor (sum of heptachlor and heptachlor epoxide expressed as heptachlor) This definition applies to all foods except infant foods
hexachlorocyclohexane (sum)	Hexachlorocyclohexane (HCH), sum of isomers, except the gamma isomer This definition applies to all foods except animal products (For animal products the alpha and beta isomers have separate MRLs)
Malathion	Malathion (sum of malathion and malaoxon expressed as malathion)
MCPA (animal products)	[Residue definition, animal products] MCPA, MCPB and MCPA thioethyl expressed as MCPA This definition applies to animal products only
MCPA (sum)	MCPA and MCPB (MCPA, MCPB including their salts, esters and conjugates expressed as MCPA) This definition applies to all foods except animal products
mepanipyrim (sum)	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine) expressed as mepanipyrim
methiocarb (sum)	Methiocarb (sum of methiocarb and methiocarb sulfoxide and sulfone, expressed as methiocarb)
methomyl (sum)	Sum of methomyl and thiodicarb expressed as methomyl
oxydemeton-methyl (sum)	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)

parathion-methyl (sum)	Parathion-methyl (sum of Parathion-methyl and paraoxon-methyl expressed as Parathion-methyl)
Permethrin	Permethrin (sum of isomers)
phorate (sum)	Phorate (sum of phorate, its oxygen analogue and their sulfones expressed as phorate)
phosmet (sum)	Phosmet (phosmet and phosmet oxon expressed as phosmet) This definition applies to all foods except animal products
pirimicarb (sum)	Pirimicarb (sum of Pirimicarb and Desmethyl pirimicarb expressed as Pirimicarb)
Prothioconazole (sum)	Prothioconazole (sum of prothioconazole-desthio and its glucuronide conjugate, expressed as prothioconazoledesthio) This definition applies to animal products only
PTU & propineb	Sum of PTU and propineb This definition applies to food for babies only
quintozene (sum)	Quintozene (sum of quintozene and pentachloro-aniline expressed as quintozene)
Prochloraz (sum)	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)
Terbufos (sum)	Terbufos (sum of terbufos, its sulfoxide and sulfone) This definition applies only to foods for babies
thiametoxam (sum)	Thiametoxam (sum of thiametoxam and clothianidin expressed as thiametoxam) There are <u>also</u> separate clothianidin MRLs
tolyfluanid (sum)	Tolyfluanid (Sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)

triadimefon & triadimenol	Triadimefon and triademenol
vinclozolin (animal products)	Vinclozolin, iprodione, procymidone, sum of compounds and all metabolites containing the 3,5-dichloroaniline moiety expressed as 3,5-dichloroaniline This definition applies to animal products only
vinclozolin (sum)	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5-dichloroaniline moiety, expressed as vinclozolin) This definition applies to all foods except animal products

Glossary

This is a 'standard' glossary which defines the key terms used in the PRiF reports. Not all the terms listed here are used in this particular report.

Acceptable Daily Intake (ADI): This is the amount of a chemical which can be consumed every day for a lifetime in the practical certainty, on the basis of all known facts, that no harm will result. It is expressed in milligrams of the chemical per kilogram of body weight of the consumer. The starting point for the derivation of the ADI is usually the 'no observed adverse effect level' (NOAEL) that has been observed in animal studies for toxicity. This is then divided by an uncertainty factor (most often 100) to allow for the possibility that animals may be less sensitive than humans and also to account for possible variation in sensitivity between individuals. The studies from which NOAELs and hence ADIs are derived take into account any impurities in the pesticide active substance as manufactured, and also any toxic breakdown products of the pesticide.

Acetylcholine: Acetylcholine is a neurotransmitter, a chemical that carries signals through the nervous system. *See cholinergic*

Acetylcholinesterase: This is an enzyme which degrades acetylcholine and is involved in the regulation of nerve impulses.

Acute Reference Dose (ARfD): The definition of the ARfD is similar to that of the ADI, but it relates to the amount of a chemical that can be taken in at one meal or on one day without appreciable health risk to the consumer. It is normally derived by applying an appropriate uncertainty factor to the lowest NOAEL in studies that assess acute toxicity or developmental toxicity.

As a matter of policy, the EU does not use NOAELs from tests that involve deliberate administration of pesticides to humans to determine ADIs and ARfDs. However, where such data have been ethically and scientifically derived some authorities, e.g. the World Health Organization, do consider such data. Where human data are used there is usually less uncertainty in the resulting reference value compared to extrapolating from animal tests to humans, and a lower uncertainty factor (most often 10) is used to account for the variation in sensitivity between individuals.

The initial risk assessments in PRiF reports use the agreed EU reference values. However, where intakes are above the EU value and a reference value based on acceptable human data is available a refined assessment, which is a more appropriate indicator of the risk, is also reported.

Analyte: This is the name for the substance that the PRiF surveys look for and measure if present; it could be a pesticide itself or a product from a pesticide when it is degraded, or metabolised.

COLEACP (Europe-Africa-Caribbean-Pacific Liaison Committee): It aims to promote the competitive export of fresh fruit, vegetables, flowers and ornamental plants from the ACP. Its specialised information and advisory services are open to all ACP companies in the horticultural export sector and are financed by the European Commission. It has two overriding objectives to enable ACP companies to comply with European food safety and traceability requirements and to consolidate the position of small-scale producers in the ACP horticultural export sector.

Cholinergic: In relation to the animal nervous system, processes and structures are cholinergic if they release or use acetylcholine.

Cryogenic Milling: Processing of commodities at very low temperatures can be achieved by milling/grinding pre-frozen samples in the presence of dry ice, a procedure known as 'cryogenic milling'.

Extensions of Authorisations for Minor Use (EAMUs): Users and authorisation holders of agricultural Plant Protection Products (PPP) may apply to have the authorisation of specific PPP's extended to cover uses additional to those authorised and shown on the manufacturer's product label. For many reasons, label recommendations of approved pesticides do not cover the control of every problem which may arise. This is particularly true for crops that are grown on a comparatively small scale in the UK as well as for pests and diseases that occur less often or which are new to the UK. As part of the process evidence on residues that would arise from the use is required, and consumer safety is evaluated and if necessary a specific MRL set. EAMU is pronounced "emu" these types of authorisations are also informally called "off labels".

Good Agricultural Practice in the Use of Pesticides (GAP): The nationally authorised safe uses of pesticides under conditions necessary for effective and reliable pest control (the way products should be used according to the statutory conditions of approval which are stated on the label). GAP encompasses a range of pesticide applications up to the highest authorised rates of use, applied in a manner which leaves a residue which is the smallest practicable. Authorised safe uses are determined at the national level and include nationally registered recommended uses, which take into account public and occupational health and environmental safety considerations. Actual conditions include any stage in the production, storage, transport, distribution and processing of food commodities and animal feed.

High-level Consumer: A term used in UK risk assessment calculations to describe the amount of food consumed by a person. In line with internationally agreed approaches, the PRiF uses the 97.5th percentile value, which is generally about three times the average amount consumed. This takes account of different eating patterns that may occur throughout the population.

Human Data: See under Acute Reference Dose

Import Tolerance: an MRL set for imported products where the use of the active substance in a plant protection product on a commodity is not authorised in the European Community (EC) or an existing EC MRL is not sufficient to meet the needs of international trade. All import tolerances are assessed for consumer safety.

Imported: The tables in the reports record whether the sample was of UK origin, or imported. This can mean different things depending on the commodity. See also 'Origin'. The PRiF report the country from where the produce has been imported only if this is clear from the packaging or labelling.

JMPR: Joint FAO/WHO Meeting on Pesticide Residues, which conducts scientific evaluations of pesticide residues in food.

LOD (Limit of Determination) and LOD MRLs: The Limit of Determination (LOD) is the lowest concentration of a pesticide residue or contaminant that can be routinely identified and quantitatively measured in a specified food, agricultural commodity or animal feed with an acceptable degree of certainty by the method of analysis.

LOD MRL (Maximum Residue Levels set at the LOD): For some pesticides and commodities insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level i.e.; at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).**

Off Label: See EMAUs

Maximum Residue Level (MRL): The maximum concentration of a pesticide residue (expressed as mg/kg) legally permitted in or on food commodities and animal feeds. MRLs are based on good agricultural practice data and residues in foods derived from commodities that comply with the respective MRLs are intended to be toxicologically acceptable.

MRLs are intended primarily as a check that GAP is being followed and to assist international trade in produce treated with pesticides. **MRLs are not in themselves 'safety limits'**, and exposure to residues in excess of an MRL does not automatically imply a hazard to health.

The MRLs applicable in the UK are now largely set under EC legislation.

Maximum Residue Levels (MRLs) reflect levels of pesticides that could occur in produce, which has been treated in accordance with good agricultural practice. Where pesticides do not give rise to readily detectable residues, or are not approved for use on particular commodities, MRLs are set at the lowest level which can be identified in routine laboratory analysis. Thus, they provide a mechanism for statutory controls on pesticides in produce which is put into circulation and for monitoring correct use of these chemicals.

If no use of a pesticide on a crop is identified when MRLs are set the tolerance for that pesticide/crop combination is set at the limit of determination (effectively zero). Limit of determination MRL are marked by a '*'

MRLs are established under the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) (England and Wales) Regulations 1999 (as amended), the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) (Scotland) Regulations 2000 and the Pesticides (Maximum Residue Levels in Crops, Food and Feeding Stuffs) Regulations (Northern Ireland) 2002. These Regulations list all statutory MRLs established under UK national or EC

procedures. Today, virtually all these MRLs are set under an ongoing EC programme and the Regulations are amended periodically as levels are set for increasing numbers of pesticides.

There are a number of pesticides which do not yet have statutory MRLs. In the absence of such MRLs we advise suppliers to adhere to any appropriate levels established by the Codex Alimentarius Commission (CAC) a United Nations body established to promote global trading standards. Codex MRLs are not statutory but have been risk-assessed when set and provide a suitable standard in the absence of a statutory MRL.

MRLs may be extended to composite and processed products but levels are not specifically laid down in legislation. They are derived by calculation on an individual basis.

Maximum Residue Levels set at the LOD (LOD MRL): See LOD MRL. For some pesticides and commodities, insufficient trials data are available on which to set a maximum residue level or there may be no use of the pesticide on that crop. In these cases, the MRL may be set at a default level, i.e. at the limit of determination (LOD) where analytical methods can reasonably detect the presence of the pesticide. **These MRLs are not based on Good Agricultural Practice (GAP).**

MRL exceedances: When a residue is found at a level higher than that set for the MRL.

MRL Exceedances and Relationship with the Acceptable Daily Intake (ADI): Before permitting any use of a pesticide, a detailed assessment is made to ensure that residues in foods derived from commodities comply with MRLs and will not give rise to unacceptable risks to consumers. MRLs do take account of consumer safety aspects and, in effect, are set at levels below safety limits. However, MRLs must not be confused with safety limits, which are expressed in terms of the acceptable daily intake (ADI) of a particular pesticide residue from all sources. The ADI (expressed as mg/kg bw/day) is the amount of chemical that can be consumed every day of an individual's entire lifetime in the practical certainty, on the basis of all known facts, that no harm will result. See ADI for further information.

Whenever unexpectedly high or unusual residues occur during monitoring, the risk to consumers, from exposure to residues at the highest levels found, is assessed by comparison of predicted intakes with the ADI or ARfD as appropriate.

No MRL: For certain pesticides an MRL may not have been set.

Metabolite: A degradation or conversion product from a pesticide when it is metabolised.

Multiple Residues: In this report this term is used to describe when more than one pesticide is found in an individual food sample. It may have arisen because the crop was treated at different times with pesticides applied singularly, or when pesticides are applied as mixtures (several pesticides mixed in the spray tank at the same time) or the marketed pesticide product contains more than one pesticide or any combination of these three situations. Mixtures may be used in response to specific pest pressures and also as part of strategies to minimise pesticide resistance building up on pest populations.

NEDI: National Estimate of Daily Intake. An estimate of intake of pesticide in the diet over the long-term to compare to the ADI. The NEDI is based on median or mean residue levels and a high level consumption (97.5th percentile value) for the daily amounts of the food item consumed over the long term. For further details on the calculation of NEDIs please refer to section 3 of the data requirements handbook: www.pesticides.gov.uk/guidance/industries/pesticides/topics/pesticide-approvals/pesticides-registration/applicant-guide/the-applicant-guide-contents.

NESTI: National Estimate of Short Term Intake. An estimate of peak intake of pesticide in the diet to compare to the ARfD. The NESTI is based on the highest residue found multiplied by a variability factor (see glossary description) and a high level consumption (97.5th percentile value) for the amount of the food item consumed over a single day. For further details on the calculation of NESTIs please refer to section 3 of the data requirements handbook: www.pesticides.gov.uk/guidance/industries/pesticides/topics/pesticide-approvals/pesticides-registration/applicant-guide/the-applicant-guide-contents.

Neurotoxicity: Neurotoxicity is the effect of substances (called neurotoxins) which alter the normal working of an animal's nervous systems and/or damage the nervous tissue.

No Observed Adverse Effect Level (NOAEL): The greatest concentration or amount of a substance, found by experiment or observation, which causes no detectable adverse alteration of morphology, functional capacity, growth, development or life span of the target organism under defined conditions of exposure.

Origin: The brand name annex reports the origins of the samples tested. This can mean different things depending on the commodity. For example, butter is often labelled as 'UK origin'; however, the majority of it comes in bulk from New Zealand and is split into smaller blocks and packaged in the UK. Lettuce is a fresh produce and 'UK origin' usually means that it has been grown and packaged in the UK. Processed commodities such as cereal bars often contain multiple raw ingredients, each of which may come from a different source/origin. Therefore, the origin of the produce usually reflects the place where it was manufactured. The PRiF report the origin as stated on the packaging or labelling of the commodity concerned, unless other more accurate information is available to indicate that the origin is from elsewhere. Some products are listed as 'unknown origin' because the labelling does not give this information.

Parent: The chemical form of a pesticide as applied to plants, as opposed to metabolites and breakdown products.

Percentile: A percentile is a value that divides a sample of measurements at a specific point when they are listed in ascending order of magnitude. For example, the 97.5th percentile from a food consumption survey is a value that is equal to or more than 97.5% of the measurements and equal to or less than 2.5% of the measurements. So, in a sample of 40 daily food consumption values, the 97.5th percentile is equal to or more than 39 of the measurements. Such high percentile estimates of food consumption are used in risk assessments as they are more protective than using average consumption levels.

Permitted Level (PL): The permitted levels (expressed as mg/kg), in specific commodities, of some substances which can be classified as pesticides but are controlled under the Miscellaneous Food Additives Regulations 1995 (S.I. 1995 No. 3187).

Pesticide: A pesticide is any substance, preparation or organism prepared or used for destroying any pest. The majority of pesticides sought by the PRiF in its monitoring are those used to control pests in agricultural crops, although non-agricultural products may be included where there is a specific reason for doing so, e.g. where there are implications in terms of possible intakes of residues.

Probabilistic Modelling: The usual estimates of consumer exposure use single high values for both consumption amounts and residue levels. Whilst these are based on realistic UK dietary survey data and residue levels, they tend to overestimate most representative intakes. This is because they do not take into account actual variations in both amounts consumed and residue levels. Probabilistic modelling is a technique that considers all the possible different combinations of consumption and residue levels. This provides information on the probability of particular intakes occurring.

Rapid Alert System for Food and Feed (RASFF): The European Commission's Rapid Alert System for Food and Feed (RASFF) allows member authorities (EU and EFTA member States) to quickly exchange information about measures taken when responding to risks detected in food or feed. This exchange of information helps authorities in countries inside the European single market to act more rapidly and in a coordinated way in response to a possible health threats caused by food or feed.

RASFFs notifications about pesticide residues are sent when a residue is over the MRL taking into account measurement uncertainty and a potential consumer risk has been identified. For pesticide residues in food traded in the single market this means when a risk assessment has identified that risk to people eating the food cannot be ruled out.

More information is available on the European Commission website at https://ec.europa.eu/food/safety/rasff_en

Relationship between GAP and MRLs: The MRL can be defined as the maximum concentration of a pesticide residue (expressed as mg/kg) likely to **occur** in or on food commodities and animal feeds, after the use of the pesticide according to the GAP.

Reporting Limit: The reporting limit is the lowest calibrated level employed during analysis to detect residues. The reporting limit may vary slightly from laboratory to laboratory depending on the equipment available and operating procedures used.

'None were detected above the Set RL': This term is used in the Brand Name Annex, where no residues were found above their reporting limit.

Residue: Residues may be present in vegetable and animal products following the application(s) of a pesticide(s). They may not only include the pesticide that was applied but other degradation or reaction products and metabolites that may be of toxicological significance.

The levels or amounts of residues present are expressed in milligrams of the chemical in a kilogram of crop/food/commodity (mg/kg), or parts per million.

Risk Assessment: A risk assessment is carried out when residues are found in foods to determine whether, at the levels found, they present a concern for consumer health or not. Consumer risk assessments are routinely conducted as part of the approval process for pesticides and are based on residue trials. Approval of a pesticide is only recommended when the consumer risk is acceptable.

Safety Factor: Values used in extrapolation from experimental studies in animals (usually 100) or humans (usually 10) to the population: for PRiF assessments this represents a value by which the NOAEL is divided to derive an ADI or ARfD. The value depends on the nature of the effect, the dose-response relationship, and the quality of the toxicological information available. The use of such a factor accounts for possible differences in susceptibility between the animal species tested and humans, and for variation between different individuals in the population. The terms 'uncertainty factor' and 'assessment factor' are also sometimes used for this factor; the PRiF will use 'safety factor'.

Sample: The nature of all samples is as designated in the EC's 'sampling' Directive – 2002/63/EC. Examples are: apple – at least 10 apples weighing at least 1 kg; grapes – at least 5 bunches, weighing at least 2 kg.

Technical Exceedances: When an MRL has been set at the LOD because there have been no data to support a higher level. In the context of this report, 'technical exceedances' always relate to produce from third countries.

Variability Factor: A value that describes the variation in residue levels between the highest unit level and the average level in samples made up of many units. Internationally this is agreed to be the 97.5th percentile unit residue level divided by the average of the sum. The variability factor multiplied by the measured residue level from a composite sample (i.e. a sample made up by mixing several units before analysis) gives an estimate of the likely higher residue levels that may have occurred in individual units. These estimated higher levels are used in short-term risk assessments involving fruit and vegetables where consumers eat only a portion of a single item, e.g. melon, or a small number of units e.g. apples and potatoes.

Ware: Ware potatoes, sometimes referred to as main crop potatoes, are harvested between August and November, and are available throughout the period August to June because they are stored under controlled temperature after October.