

This is a paper for discussion. It does not represent the views of the Committee and must not be quoted, cited or reproduced.

MUT/2020/17

**COMMITTEE ON MUTAGENICITY OF CHEMICALS IN FOOD, CONSUMER
PRODUCTS AND THE ENVIRONMENT (COM)**

Guidance on the genotoxicity testing strategies for germ cell mutagens

Continued consideration and comments of the updated COM Guidance document on the genotoxic testing strategies for germ cell mutagens.

Members are asked to complete review of this latest draft as attached and consider the following questions:

1. The OECD Test Guidelines have been listed in paragraphs 8 – 15 in ascending numerical order. Could members clarify whether each of these is still in use and a note to that effect will be added to the text before finalisation.
2. Could members confirm whether the statement regarding TG 488 in paragraph 12 is accurate “However, at the current time the chemical database for this test in germ cells is limited”.
3. Do members have an update to the comet assay validation exercise cited in paragraph 34.
4. Do members have an update to the SCSA and TUNEL assay validation exercise cited in paragraph 37.
5. Could members provide additional references where indicated in the text.

Secretariat

November 2020

24

25 Background

26 1. The Committee on Mutagenicity of Chemicals in Food, Consumer Products and
27 the Environment (COM) has a remit to provide UK Government Departments and
28 Agencies with advice on the most suitable approaches to testing chemical substances
29 for genotoxicity. The COM views regarding the most appropriate strategy for
30 genotoxicity testing are outlined in full in the COM (202x) “Guidance On A Strategy
31 For Genotoxicity Testing Of Chemical Substances”.

32 2. In brief, the COM recommend a staged approach to genotoxicity testing. **Stage**
33 **0**, in the absence of test data from adequately designed and conducted genotoxicity
34 tests, consists of preliminary considerations of the test chemical substance, including,
35 physico-chemical properties, Structure Activity Relationships (SAR), and information
36 from screening tests. **Stage 1** consists of *in vitro* genotoxicity tests that provide
37 information on three types of genetic damage (namely, gene mutation, chromosomal
38 damage and aneuploidy) and gives appropriate sensitivity to detect chemical
39 genotoxins. **Stage 2** consists of *in vivo* genotoxicity tests which are chosen on a case-
40 by-case basis to address any genotoxic endpoints identified in Stage1; investigate
41 genotoxicity in tumour target tissue(s) and/or site of contact tissues; investigate
42 potential for germ cell genotoxicity; and investigate potential genotoxicity for chemicals
43 where high/moderate and prolonged exposure is anticipated, even if negative in Stage
44 1.

45 3. A mutation in the germ cells of sexually-reproducing organisms may be
46 transmitted to the offspring, whereas a mutation that occurs in somatic cells may be
47 transferred only to descendant daughter cells. Mutagenic chemicals may present a
48 hazard to health since exposure to a mutagen carries the risk of inducing germ-line
49 mutations with the possibility of inherited disorders, and the risk of somatic mutations
50 including those leading to cancer.

51 4. The COM affirms that a chemical considered a positive *in vivo* somatic cell
52 mutagen should also be considered as a possible germ cell mutagen unless data can
53 be provided to the contrary, as most, if not all, germ cell mutagens are also genotoxic
54 in somatic cells. It has been noted that some rare examples have been reported that
55 are contrary to this statement (e.g. sodium orthovanadate, (Attia et al., 2005).
56 However, the data on such compounds are conflicting and it is not known, for example,
57 whether somatic mutations or DNA strand breaks would have been identified if other
58 test systems (e.g. transgenic assays and the comet assay) had been used and other
59 tissues sampled (Attia et al., 2005; Ciranni et al., 1995; Witt et al., 2003).

60
61 5. There are also examples of germ cell mutagens which affect specific stages of
62 gametogenesis in males (Adler, 2008) and where there are differences between male
63 and female germ cell genotoxicity (Bishop, 2003). Currently, the focus for germ cell

mutagenicity assays is on male germ cells due to the accessibility of sperm. However, the gap relating to female germ cell assays in regulatory testing is recognised. The male germ cell assays described in this discussion document differ in the specificity, sensitivity and the endpoint detected. It should be noted that all such assays must ensure that the most appropriate phases of spermatogenesis are being tested through specified sample collection timings (Yauk et al., 2015).

6. The development of testing strategies for germ cell mutagens is a rapidly evolving field. Therefore the COM considered it appropriate to prepare a supplementary document on the topic, to support the COM “*Guidance On A Strategy For Genotoxicity Testing Of Chemical Substances*” which can be updated at regular intervals as new information becomes available (COM, 202x). This discussion paper seeks to provide a brief summary of test methodologies that are currently used or under development and/or validation, to assess germ cell mutagenicity. The strategy detailed here closely aligns with the views expressed by The International Workshops on Genotoxicity Testing (IWGT), with findings from workshops in 2013 and 2017 included in this document.

OECD Test Guidelines

7. Classification of a substance as a germ cell mutagen should be based on the findings from well conducted, scientifically validated tests in a weight of evidence approach. Where germ cell testing is indicated, there are a number of OECD test guidelines to assess germ cell mutations. These are briefly described below and, for ease of reference, are presented in ascending numerical order.

Dominant lethal test (OECD TG 478)

8. The dominant lethal test (DLT, OECD TG 478) (OECD, 2016a) has been the most widely used of the germ cell mutagenicity assays with only minor changes being introduced since its development in 1984. The DLT is usually conducted in male rats or mice and provides information on unstable chromosome changes in gametes that lead to fetal death after fertilisation in non-exposed mated females; indications on the stage of gametogenesis affected can also be determined (COM, 2011). Pre- and post-implantation embryonic losses are considered to be due to severe structural or numerical chromosomal changes inherited from the father (Brewen et al., 1975; Marchetti et al., 2004). The limitations of the assay are that cytotoxicity cannot be excluded as a cause of embryonic death and that the endpoint is not truly heritable. However, the DLT has been well standardised and used to assess many chemicals; some of these have also been tested in the HTT assay with a good correlation of positive results being seen between the two assays (Yauk et al., 2015).

Cytogenetic analysis of spermatogonia or embryos (OECD TG 483)

9. The cytogenetic analysis of spermatogonial metaphases (OECD TG 483) (OECD, 2016b) is a standardised method to detect chromosomal aberrations in male germ cells of mice and rats (Yauk et al., 2015). Chromosome painting techniques have been applied to the method which allows stable balanced aberrations (e.g. reciprocal translocations) to be distinguished from unstable aberrations (e.g. acentric fragments, dicentric chromosomes) (Marchetti and Wyrobek, 2003). Technically the method is challenging and so not widely used. The main limitation however, is that transmission of mutagenic effects to mature gametes and offspring is not demonstrated, as any possible mutagenicity is observed at the beginning of germ cell differentiation (Marchetti and Wyrobek, 2005).

Heritable translocation (OECD TG 485) and specific locus tests

10. The mouse heritable translocation test (HTT; OECD TG 485) (OECD, 1986) was previously viewed as the gold standard assay for determining the transmission of germ cell mutations to the offspring of exposed parents. The mouse HTT is defined by the COM as detecting '*heritable structural chromosome changes (i.e. translocations) in mammalian germ cells as recovered in first-generation progeny*'. The mouse specific locus test (SLT) is described by the COM as '*a technique used to detect recessive induced mutations in diploid organisms; a strain that carries several known recessive mutants in a homozygous condition is crossed with a non-mutant strain that has been treated to induce mutations in its germ cells; induced recessive mutations allelic with those of the test strain will be expressed in the progeny*'.

11. Following the development of molecular cytogenetics and genomics technologies, these assays are **now viewed negatively** ^[RB1] as requiring large numbers of animals (including the use of a mutant mouse strain in the SLT) and as being labour intensive. As a result, these assays are no longer performed in the UK and Europe, **however TG 485 is maintained by OECD as some countries have included this TG in their legislative guidance.**

Transgenic rodent somatic and germ cell mutation assay (OECD TG 488)

12. The transgenic rodent mutation assays (TGR; OECD TG 488) (OECD, 2020) are based on the detection of a mutation in a transgenic sequence that can be isolated from most rodent tissues and expressed in a bacterial system (Yauk et al., 2015). The assays can be used to assess gene mutations in a wide range of rodent tissues (including germ cells) using all routes of administration (Lambert et al., 2005; Kirkland et al., 2019a) and is particularly valuable when investigating gene mutation as the genotoxic endpoint. Determination of the mutation spectrum (base substitutions, insertions/deletions, frameshifts) following chemical exposure of testicular cells and epididymal sperm has been described (Lambert et al., 2005). However, at the current

time the chemical database for this test in germ cells is limited. ~~and its ability to detect changes during different spermatogenic phases is still unknown.~~

~~42.13.~~ Limitations of TG 488 include the use of some mutation reporter genes that are limited in use and availability. There are sufficient data to assess the performance of the LacZ bacteriophage mouse (MutaTMmouse), LacZ plasmid mouse, lacI mouse and rat (BigBlue[®]) (including use of λ cII transgene), , and the gpt delta (gpt and Spi-) mouse and rat models, although it is noted that the *gpt* models are not widely used and are less well validated (COM, 202x). In addition, the TGR assay only infers potential inheritance of mutations, however the cII positive selection assay can be used for assessing mutations in the BigBlue[®] and MutaTMmouse models which does directly assess heritability (Singer et al., 2006). It has also been reported that the assays may not detect some types of mutations, including large deletions/insertions for some TGR loci, and rearrangements or copy number variants (CNVs) (Yauk et al., 2015). Molecular analysis (sequencing) of the mutations can provide additional information here (OECD, 2020).

~~43.14.~~ The sampling time is a critical variable in the TGR assay as it is determined by the period needed for mutations to be fixed. This period is tissue-specific and appears to be related to the turnover time of the cell population. Both somatic and germ cells can be sampled in the TGR assay using a 28 day administration period followed by a 28 day sampling time. This allows significant reductions in animal usage, in line with the 3Rs principles¹, cost, and time and enables a quantitative comparison of the same mutagenic endpoints between somatic and germ cell tissues (Yauk et al., 2015).

~~44.15.~~ Development of the TGR assay for detecting female germ cell mutations is not considered possible due to the low numbers of oocytes available per female for analysis, and therefore is not considered further here (Yauk et al., 2015; OECD, 2020).

Detection of genotoxic and mutational changes in sperm

~~45.16.~~ Genotoxicity tests in sperm can be applied in the same way to humans and animals, providing a direct comparison between biomonitoring and experimental data. There are a number of assay systems that detect different types of pre-mutational and mutational changes in sperm; these are outlined below. Importantly, these could offer quick, higher throughput pre-screening tools for detecting germ cell mutagens, even though they do not assess heritable effects. Many of these have not currently undergone standardisation and harmonisation processes and are discussed more fully in paragraphs 28 - 37.

- Comet - detects DNA strand breaks and abasic sites;

- TUNEL - Terminal deoxynucleotidyl transferase dUTP nick end labelling detects DNA fragmentation;
- SCSA – sperm chromatin structure assay detects chromatin packing alterations;
- FISH – fluorescent in situ hybridisation detects numerical and structural chromosome changes.

Other toxicity assays providing evidence of potential germ cell genotoxicity

~~46-17.~~ Standard repeat dose and reproductive toxicity studies are a potential source of information that may indicate germ cell genotoxicity. In both types of study, cytotoxic and reproductive endpoints can indicate that a substance has been delivered to particular organs, including gonadal tissue and associated male and female germ cells. These are described further in paragraphs 18 - 23. It should be noted that these studies do not assess mutagenicity specifically and further studies would need to be carried out to confirm this as a mechanism of action.

Segmented reproductive toxicity tests

~~47-18.~~ Segmented studies assess adverse effects following exposure at particular time periods of development rather than the entire life cycle. There are a number of segmented designs within guidance from The International Conference on Harmonization (ICH) and the OECD:

- ICH guideline S5(R2) describes three segmented phases for the testing of pharmaceuticals; a fertility and early embryonic development study with exposure of males for 4 weeks prior to mating and of females for 2 weeks prior to mating, through to implantation; exposure of the pregnant dam from implantation through fetal development (assessing organogenesis); pre- and post-natal developmental (PPND) with exposure of the dam from implantation, through lactation until pup weaning.
- OECD TG 421 Reproduction/Developmental Toxicity Screening Test (OECD, 2016c) has a similar pre-mating exposure in males and females, with continuous exposure of females to post-natal day (PND) 4. TG 421 is designed to provide limited information regarding the effects of exposure of the test chemical on fertility (male and female reproductive performance such as gonadal function, mating behaviour, conception) and development of the conceptus and parturition. Although this test provides an assessment of transferred effects from exposed males (which may include mutagenic effects), any effects may also be due to exposure *in utero*. No specific assessment of mutagenicity is carried out.
- OECD TG 414 Prenatal Developmental Toxicity Study (OECD, 2018a) assesses the effects of *in utero* exposure to a test chemical from the implantation phase through to parturition. TG 414 is designed to provide general information on the effects of prenatal exposure to a test chemical on

the developing organism. The parameters assessed in TG 414 include maternal effects (including death), structural abnormalities and/or altered growth in the fetus. Although it is possible that some effects seen will be due to mutagenicity, no specific assessment of this is carried out.

[18-19](#). Within these studies, adverse effects on fertility and litter size are determined from the pregnant female, and developmental outcomes can be assessed in fetal tissue which can be examined to assess morphologic changes and through functional tests in pups, including reproductive performance testing (Yauk et al., 2015).

Continuous cycle reproductive toxicity tests

[19-20](#). Continuous cycle study designs assess all the different stages of the reproductive life cycle from germ cell through fetal development to adulthood and are often multigenerational. There are two main approaches for continuous study designs:

- The National Toxicology Program's (NTP) reproductive assessment by continuous breeding (RACB) (Gulati et al., 1991);
- The OECD Two-Generation Reproduction Toxicity Study (OECD TG 416) (OECD, 2001).

[20-21](#). As described for the segmented reproductive toxicity test (paragraphs 18 – 19), the continuous cycle reproductive studies indirectly assess potential effects of germ cell mutagenicity through histopathological analysis of the reproductive and endocrine systems; however, mutagenicity *per se* is not assessed. Effects on fertility and fecundity are assessed through mating of the F0 animals. However, limitations arise in that for any effects arising in the F1 generation, it is not possible to distinguish between those passed on from the F0 generation or those due to *in utero* exposure.

One generation reproduction toxicity study

[21-22](#). The extended one-generation study design (enhanced pre and postnatal study) (OECD TG 443) (OECD, 2018b) has been developed from the one generation reproduction toxicity study (OECD TG 415) (no longer an active Test Guideline) and multigenerational reproductive studies. In the extended study rodents are dosed before mating through gestation with exposure being stopped at various times, with either necroscopy or mating to produce an F1/F2 generation.

Repeat dose toxicity studies

[22-23](#). Short-term and long-term repeat dose toxicity studies (e.g. 90 day studies) can be combined with reproduction/development toxicity screening tests (e.g. OECD TG 408 Repeated Dose 90-day Oral Toxicity Study and 422 Combined Repeated Dose Toxicity Study with the Reproduction/Developmental Toxicity Screening Test) (OECD, 2018c; 2016c). As previously discussed (paragraphs 17 – 23), the assessment of

ovarian and testicular histopathology, sperm count, motility and morphology, might be used to indicate potential germ cell effects.

Assays under development and/or validation

[23-24](#). A number of new assays to assess germ cell mutagens are currently being developed and validated. In addition, modifications to current OCED TGs are being explored.

Expanded simple tandem repeat (ESTR) assays

[24-25](#). ESTRs are long homogenous arrays of relatively short repeats (4–9bp) which have a very high spontaneous mutation rate of length changes both in germline and somatic cells (Bois et al., 1998). ESTR loci are considered to be a class of expanded microsatellites, with the spontaneous mutation being replication-driven (Hardwick et al., 2009; Shanks et al., 2008). The analysis of length change mutations occurring at ESTR loci has been utilised for assessing male germ cell mutagenicity in mice (Barber et al., 2009; Dubrova et al., 2000; Vilariño-Güell et al., 2003; Marchetti et al., 2011).

[25-26](#). The sensitivity of the assay has been increased through the use of single-molecule PCR to detect ESTR mutations, which has also decreased the numbers of animals required and assay duration. In addition, this approach is applicable to human studies; for example, mutation induction has been measured in mice using human clinically-relevant doses of anticancer drugs (Glen et al., 2008).

[26-27](#). There are some limitations currently reported for the ESTR assay. The mutations detected occur in a very specific genomic context of tandem repeats. In addition, the mechanism underlying ESTR mutation induction is not fully defined. One hypothesis is that non-targeted events cause mutagen-related DNA damage elsewhere in the genome, which leads to an increased mutation rate at the ESTR loci.

[27-28](#). The ESTR assay can be integrated with standard genetic toxicology tests in mice, however it is currently not known whether ESTR mutations can be assayed in testicular cells sampled under these protocols. Further, integration may require additional animals to be used specifically for the ESTR assay, with an additional appropriate sampling time. From a methodological perspective, the assay is also technically challenging which can lead to variable inter-laboratory results.

Spermatid micronucleus (MN) assay

[28-29](#). OECD TGs currently exist for the analysis of MN formed as a consequence of chromosome damage and/or spindle malfunction, in *in vitro* (OECD TG 487 In Vitro Mammalian Cell Micronucleus Test) (OECD, 2016d) and *in vivo* (OECD TG 474 Mammalian Erythrocyte Micronucleus Test) (OECD, 2016e) somatic cells. These are widely used and predominant assays for somatic cell testing, with high sensitivity

facilitated by flow cytometric analysis. Attempts are being made to develop an equivalent germ cell assay.

[29-30](#). The spermatid MN assay detects MN originating during meiosis. It was originally developed in rats and subsequently adapted for mouse spermatids. The assay is able to be combined with other genotoxicity tests, including the transgenic rodent assay, and potentially analysis in erythrocytes (Yauk et al., 2015).

[30-31](#). Some current limitations of the spermatid MN assay include its labour-intensive nature, which limits the number of cells that can be scored and hence the sensitivity. This is being addressed through development of automated detection of MN by flow cytometry, as exists for somatic cells. In addition, although it is not known what the fate of a sperm cell carrying MN is, it is considered unlikely that the micronuclei would be inherited (Yauk et al., 2015).

Sperm Comet assay

[31-32](#). OECD TG 489 (In Vivo Mammalian Alkaline Comet Assay) (OECD, 2016f) describes the *in vivo* alkaline Comet assay for the measurement of DNA strand breaks in single cells. The Comet assay has been used to assess genotoxic hazard for a large number of chemical and physical genotoxicants both *in vivo* and *in vitro*. Although the main use of the assay has been for the assessment of somatic cells, the assay has been conducted both on mature sperm and on germ cells isolated from the seminiferous tubules (Speit et al., 2009; Haines et al., 2001; Haines et al., 2002). When applied to germ cells, the assay does not show heritability but does indicate genotoxicity.

[32-33](#). There are, however, a number of limitations that need to be addressed before the assay could potentially be applied to assess germ cell DNA damage for regulatory purposes (Kirkland et al., 2019b). The exposure protocol outlined in TG 489 would result in only fully mature sperm being exposed, which have a high resistance to DNA damage. Although the analysis of germ cells collected from the seminiferous tubules is not fully validated, it is known that two different germ cell populations (spermatocytes and elongating spermatids) are present. For both cell populations, DNA double strand breaks are part of the normal process of development (meiotic recombination in spermatocytes and chromatin compaction in elongating spermatids) which may lead to false positive findings. Mature sperm also require a pre-digestion step before analysis in the Comet assay, which can lead to poorly reproducible results (Yauk et al., 2015).

[33-34](#). There are currently initiatives underway to standardise the Comet assay, with 10 laboratories worldwide developing fully validated protocols to ensure data reproducibility (Yauk et al., 2015).

Sperm chromatin quality assays

[34.35.](#) Two other commonly used assays to assess the integrity of sperm DNA include the sperm chromatin structure assay (SCSA) and the terminal deoxynucleotidyl transferase-mediated (TdT) deoxyuridine triphosphate (dUTP) nick end labeling assay (TUNEL). Both assays were developed around 30 years ago and validation is more advanced in humans than in animals.

[35.36.](#) SCSA uses flow cytometry methods to assess the susceptibility of sperm DNA to acid-induced denaturation, as denaturation is linked to the presence of single stranded DNA, an indicator of potential genotoxicity (Sills et al., 2004). The TUNEL assay measures DNA breaks *in situ* as assessed by the incorporation of dUTP at the sites of breaks (Gorczyca et al., 1993). Both assays, therefore, measure different aspects of DNA integrity.

[36.37.](#) There are currently initiatives underway to standardise the SCSA and TUNEL assays, with 10 laboratories worldwide developing fully validated protocols to ensure data reproducibility. It is hoped that validation of the assays in humans will allow rapid transfer to animal models (Yauk et al., 2015).

[37.38.](#) The main limitation with using sperm DNA integrity as an endpoint for genotoxicity testing is that currently we do not understand the mechanisms and consequences of sperm chromatin damage. The integrity of sperm chromatin has been identified as a contributing factor to a healthy pregnancy and offspring (Aitken et al., 2013; Aitken et al., 2009; Lewis and Simon, 2010; Robinson et al., 2012) however, clinically relevant parameters that would allow chromatin integrity to be assessed have not currently been defined.

Whole genome sequencing

[38.39.](#) Advancements in genome sequencing technologies allow detection of the effects of mutagens on heritable germ cells. These technologies have been applied to the full genomic sequencing of 78 individuals and findings suggested that the father's age is a dominant factor in determining the number of *de novo* mutations in their offspring (Kong, 2012). If genome-wide mutation spectra and frequencies in rodent models are shown to be comparable to humans, this technology has the potential to determine phenotypic consequences to an organism as a whole (Yauk et al., 2015).

[39.40.](#) However, as the methodology is still in development it has not been applied from a toxicological basis, and extensive validation will be needed. Other limitations include the high costs and long analysis time, which are expected to be reduced with improved data handling (Yauk et al., 2015).

Copy number variants (CNVs)

~~40.41.~~ CNVs comprise a structural variation of DNA ranging in size from 50 base pairs to megabases, which alters or rearranges the number of copies of specific DNA segments. CNVs account for around 12 % of genetic variation in humans and are considered to be related to a broad range of human genetic disorders (Stankiewicz and Lupski, 2010; Campbell, 2013; Girirajan and Eichler, 2010; Sebat et al., 2004; Lupski, 2007). CNVs are not detected using currently available genotoxicity testing assays and require high- resolution array comparative genomic hybridization (or aCGH) and SNP (single nucleotide polymorphism) microarray technologies (Yauk et al., 2015).

~~41.42.~~ Both technologies have been applied in the clinic to identify the sources of idiopathic diseases (Dittwald et al., 2013; Wiszniewska et al., 2014; Cheung et al., 2005; Boone et al., 2013; Pham et al., 2014) but only limited assessments of the effects of mutagens on CNV formation have been reported (Arlt et al., 2009; Arlt, 2011; Arlt et al., 2014). It has been shown *in vitro* that replication stress (for example through exposure to hydroxyurea or low doses of ionising radiation) can lead to the formation of CNVs. Increases in *de novo* CNVs is also associated with increasing paternal age (Sun et al., 2012).

~~42.43.~~ The major current limitation of this technology is the lack of evidence to show its application *in vivo* and extensive development and validation is therefore required (Yauk et al., 2015).

High-throughput analysis of egg aneuploidy in nematode *C. elegans*

~~43.44.~~ High throughput screening (HTS) tools for chemical testing is a rapidly developing field, which is aimed at increasing chemical testing capacity whilst reducing animal use. A major gap exists in HTS assays for the detection of mutagens and aneugens (Knight et al., 2009). Existing assays focused on the initiation of a DNA damage response have low sensitivity and do not consider effects on germ cells. A new screening tool, which is currently under development, utilises the nematode *C. elegans* to measure chromosome segregation errors occurring in eggs and has been proposed as an HTS assay for Tier 1 screening of female germ cells (Yauk et al., 2015). Preliminary validation using 50 chemicals showed an accuracy of 69 % (average of sensitivity and specificity) in predicting the ability of chemicals that cause reproductive toxicity in rodents (Allard et al., 2013).

~~44.45.~~ *C. elegans* is an established model system in genetics as there is a good degree of conservation with humans in key meiotic pathways. Limitations concerning the applicability of the relationship of aneuploidy in *C. elegans* to the same potential outcome in humans, has been raised (Yauk et al., 2015).

420 **Summary of assays under development or undergoing validation**

421

422 **Table 1 - Advantages and disadvantages of assays under development or**
423 **undergoing validation**

Endpoint	Advantages	Disadvantages
Transgenic rodent mutation	Can be performed on a wide range of tissues; Allows a comparison of somatic and germ cell sensitivity/specificity; Neutral gene; Scores gene mutation; OECD guideline; Can be integrated into multiple test strategies.	Transgenic rodent model used; Scores mutations in a non-transcribed exogenous gene; Performed on germ cells so inheritance is assumed; May miss some types of mutations.
Tandem repeat assays	Endogenous loci; High spontaneous mutation rate; Adaption to any species possible; Links shown between some markers and disease; Sensitive at low doses; Integration into multiple test strategies possible (requires validation).	Indirect mechanism of mutation with unknown mode; Non-coding markers; Relevance of tandem repeat mutation to gene mutations is unclear; Small dynamic range; Technically challenging.
Spermatid micronucleus (MN)	Can be integrated into transgene mutation reporter assay and other toxicity tests; Can be performed in any species; Directly comparable to somatic MN to assess germ cell specificity/sensitivity.	Methodology is laborious (but potential for flow cytometry modifications); Small database; Performed on germ cells so inheritance is assumed.
Sperm comet assays	Can be performed in any species; Technically simple; Directly comparable to most somatic cell types; Detects a variety of DNA damage.	Difficult to integrate with other tests; High inter-laboratory and inter-study variability; Biological relevance of endpoint unclear; Technically challenging; Pre-mutational damage only detected.
Sperm chromatin structure	Rapid technique (flow cytometry approach); Can be performed in any species including humans; Major validation exercises are underway.	Performed on germ cells so inheritance is assumed; Pre-mutagenic lesion detected; Mechanisms causing changes in chromatin are not known; Technically challenging giving high inter-laboratory and inter-study variability.

424 Source: adapted from (Yauk et al., 2015)

Do the available assays reflect human relevant endpoints?

[45-46.](#) There is a spectrum of mutational events occurring *in vivo* that have the potential to impact on human health, and new genomics tools allow for the quantitation of genome-wide mutation rates. Single nucleotide variants (SNVs) and CNVs may affect coding and non-coding DNA sequences; for example, it has been reported that 76 % of SNVs originate in the paternal lineage (Campbell et al., 2012; Conrad et al., 2011; Roach et al., 2010).

[46-47.](#) The mutation rate (per locus and per total nucleotide number affected) is higher for CNVs than SNVs. It has been estimated that one large *de novo* CNV (>100 kbp) occurs per 42 births in humans, compared to an average of 61 new SNVs per birth; however, the average number of base pairs affected by large CNVs is 8–25 kbp per gamete versus 30.5 bp per gamete for SNVs (Yauk et al., 2015). CNVs are caused by chromothripsis events whereby multiple *de novo* rearrangements in a single event (Stankiewicz and Lupski, 2010).

[47-48.](#) A number of additional functional genomic changes also arise, including:

- small insertions and deletions;
- mobile element insertions;
- tandem repeat mutations;
- translocations; and
- aneuploidies

[48-49.](#) Proportionally higher *de novo* mutation rates are reported for microsatellites than for SNVs, which is considered an important source of genetic variation (Sun et al., 2012). An inverse relationship has been reported between mutation size and frequency, meaning that although more rare, the number of nucleotides affected by large genomic changes, including CNVs and aneuploidies, is orders of magnitude greater (Yauk et al., 2015).

[49-50.](#) In humans, epidemiological studies look to measure the phenotypic effects of induced dominant mutations occurring in the descendants of exposed parents. Importantly, such studies have shown that as many mutations occurring in humans are recessive, phenotypic changes are not apparent for several generations until conception occurs with a complementary mutation or the mutation occurs in a somatic cell.

[50-51.](#) Some of these potentially important genomic changes may therefore not be effectively captured by both the existing battery of genotoxicity testing assays nor by those under development.

What is the current status of regulatory requirements for germ cell testing?

[51-52.](#) The testing of chemicals for germ cell mutagenicity is a regulatory requirement for many organisations worldwide, including the World Health Organisation / International Programme on Chemical Safety (WHO/IPCS), Globally Harmonized System of Classification and Labelling of Chemicals (GHS) and the regulatory agencies in the US, Canada, Japan, UK and the EU. Although many other countries follow the approach taken by the US, India and Australia do not require germ cell mutation tests for regulatory purposes. Genetic toxicity tests used across organisations comprise three tiers, with Tier 1 containing *in vitro* and somatic *in vivo* tests and Tiers 2 and 3 the supporting germ cell studies that can be requested by regulatory bodies under certain conditions. For example, Tier 2 contains DNA damage assays in the testes or spermatogonia and Tier 3 the gene cell mutation tests.

[52-53.](#) The testing of pharmaceuticals for registration under the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) does not stipulate that germ cell assays should be carried out, rather it is assumed that *in vivo* somatic tests and carcinogenicity data will provide sufficient predictivity/protection for germ cell effects.

[53-54.](#) The WHO/IPCS Harmonised Scheme states that a positive *in vivo* somatic cell mutagen can trigger testing for germ cell mutagenicity, but this is not required. Optional recommended tests include transgenic mouse models, the ESTR assay, the spermatogonial chromosome aberration assay, chromosome aberration analysis by FISH, the Comet assay, and assays for DNA adducts. The WHO/IPCS tests in offspring include the ESTR assay, the DLT, the HTT, and the SLT (Yauk et al., 2015).

[54-55.](#) The GHS, together with OECD, ECHA and many other countries categorise mutagens according to three criteria:

- Category 1A – chemicals known to induce heritable mutations in germ cells of humans (based largely on human evidence);
- Category 1B – chemicals that should be regarded as if they induce heritable mutations in germ cells of humans (based largely on experimental animal data);
- Category 2 – chemicals that cause concern for induction of heritable mutations in germ cells of humans.

[55-56.](#) In the EU, for example, under the REACH regulations, any genotoxic agent in somatic cells is evaluated for germ cell mutagenicity using bioavailability and *in vivo* data. Where no data are available, the chemical can be further tested using relevant germ cell assays. Issues around the types of studies able to provide data suitable for distinguishing between mutagen categories 2 and 1B were discussed at a joint workshop between the Member State Committee (MSC) and Committee for Risk Assessment (RAC). Workshop participants agreed that refinement of the current MSC

approach was possible with regards to follow-up testing of positive somatic cell mutagens, including testing for mutagenic potential in both somatic and germ cells in the same study (ECHA, 2019).

~~56-57.~~ Germ cell mutagenicity is an established regulatory endpoint and existing assays have identified >50 substances as germ cell mutagens in rodents. It has been noted by Yauk and colleagues that no agent has currently been regulated solely as a germ cell mutagen or evaluated to be a human germ cell mutagen (Yauk et al., 2015) i.e. there are no known Cat 1A substances.

NCET at WRc/IEH-C under contract supporting the PHE COT Secretariat
November 2020

References

- Adler, I.-D. 2008. Mutagenicity Tests *in vivo*. In: GREIN, H., SNYDER, R. (ed.) *Toxicology and Risk Assessment* Wiley and Sons, UK
- Aitken, R. J., Bronson, R., Smith, T. B., et al. 2013. The source and significance of DNA damage in human spermatozoa; a commentary on diagnostic strategies and straw man fallacies. *Mol Hum Reprod*, 19, 475-85.
- Aitken, R. J., De Iuliis, G. N. & Mclachlan, R. I. 2009. Biological and clinical significance of DNA damage in the male germ line. *Int J Androl*, 32, 46-56.
- Allard, P., Kleinstreuer, N. C., Knudsen, T. B., et al. 2013. A C. elegans screening platform for the rapid assessment of chemical disruption of germline function. *Environ Health Perspect*, 121, 717-24.
- Arlt, M. F., Mülle, J. G., Schaibley, V. M., et al. 2009. Replication stress induces genome-wide copy number changes in human cells that resemble polymorphic and pathogenic variants. *Am J Hum Genet*, 84, 339-50.
- Arlt, M. F., Ozdemir, A.C. Birkeland, S.R. Et Al. 2011. Hydroxyurea induces de novo copy number variants in human cells. *Proc. Natl. Acad. Sci. U. S. A*, 108, 17360–17365.
- Arlt, M. F., Rajendran, S., Birkeland, S. R., et al. 2014. Copy number variants are produced in response to low-dose ionizing radiation in cultured cells. *Environ Mol Mutagen*, 55, 103-13.
- Attia, S. M., Badary, O. A., Hamada, F. M., et al. 2005. Orthovanadate increased the frequency of aneuploid mouse sperm without micronucleus induction in mouse bone marrow erythrocytes at the same dose level. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 583, 158-167.
- Barber, R. C., Hardwick, R. J., Shanks, M. E., et al. 2009. The effects of in utero irradiation on mutation induction and transgenerational instability in mice. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 664, 6-12.
- Bishop, J. B. 2003. Female-specific reproductive toxicities following preconception exposure to xenobiotics. *Adv Exp Med Biol*, 518, 1-9.
- Bois, P., Williamson, J., Brown, J., et al. 1998. A Novel Unstable Mouse VNTR Family Expanded from SINE B1 Elements. *Genomics*, 49, 122-128.
- Boone, P. M., Soens, Z. T., Campbell, I. M., et al. 2013. Incidental copy-number variants identified by routine genome testing in a clinical population. *Genet Med*, 15, 45-54.
- Brewen, J. G., Payne, H. S., Jones, K. P., et al. 1975. Studies on chemically induced dominant lethality. I. The cytogenetic basis of MMS-induced dominant lethality in post-meiotic male germ cells. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 33, 239-249.

- Campbell, C. D., Chong, J. X., Malig, M., et al. 2012. Estimating the human mutation rate using autozygosity in a founder population. *Nat Genet*, 44, 1277-81.
- Campbell, C. D., Eichler, E.E. 2013. Properties and rates of germline mutations in humans. *Trends Genet.*, 29, 575-584.
- Cheung, S. W., Shaw, C. A., Yu, W., et al. 2005. Development and validation of a CGH microarray for clinical cytogenetic diagnosis. *Genet Med*, 7, 422-32.
- COM 2011. Guidance On A Strategy For Genotoxicity Testing Of Chemical Substances
- Conrad, D. F., Keebler, J. E., Depristo, M. A., et al. 2011. Variation in genome-wide mutation rates within and between human families. *Nat Genet*, 43, 712-4.
- Dittwald, P., Gambin, T., Szafranski, P., et al. 2013. NAHR-mediated copy-number variants in a clinical population: mechanistic insights into both genomic disorders and Mendelizing traits. *Genome Res*, 23, 1395-409.
- Dubrova, Y. E., Plumb, M., Brown, J., et al. 2000. Induction of minisatellite mutations in the mouse germline by low-dose chronic exposure to γ -radiation and fission neutrons. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 453, 17-24.
- Girirajan, S. & Eichler, E. E. 2010. Phenotypic variability and genetic susceptibility to genomic disorders. *Hum Mol Genet*, 19, R176-87.
- Glen, C. D., Smith, A. G. & Dubrova, Y. E. 2008. Single-molecule PCR analysis of germ line mutation induction by anticancer drugs in mice. *Cancer Res*, 68, 3630-6.
- Gorczyca, W., Gong, J. & Darzynkiewicz, Z. 1993. Detection of DNA strand breaks in individual apoptotic cells by the in situ terminal deoxynucleotidyl transferase and nick translation assays. *Cancer Res*, 53, 1945-51.
- Gulati, D. K., Hope, E., Teague, J., et al. 1991. Reproductive toxicity assessment by continuous breeding in Sprague-Dawley rats: a comparison of two study designs. *Fundam Appl Toxicol*, 17, 270-9.
- Haines, G. A., Hendry, J. H., Daniel, C. P., et al. 2001. Increased levels of comet-detected spermatozoa DNA damage following in vivo isotopic- or X-irradiation of spermatogonia. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 495, 21-32.
- Haines, G. A., Hendry, J. H., Daniel, C. P., et al. 2002. Germ cell and dose-dependent DNA damage measured by the comet assay in murine spermatozoa after testicular X-irradiation. *Biol Reprod*, 67, 854-61.
- Hardwick, R. J., Tretyakov, M. V. & Dubrova, Y. E. 2009. Age-related accumulation of mutations supports a replication-dependent mechanism of spontaneous mutation at tandem repeat DNA Loci in mice. *Mol Biol Evol*, 26, 2647-54.
- Kirkland, D., Levy, D. D., Lebaron, M. J., et al. 2019a. A comparison of transgenic rodent mutation and in vivo comet assay responses for 91 chemicals. *Mutat Res Genet Toxicol Environ Mutagen*, 839, 21-35.
- Kirkland, D., Uno, Y., Luijten, M., et al. 2019b. In vivo genotoxicity testing strategies: Report from the 7th International workshop on genotoxicity testing (IWGT). *Mutat Res*, 847, 403035.
- Knight, A. W., Little, S., Houck, K., et al. 2009. Evaluation of high-throughput genotoxicity assays used in profiling the US EPA ToxCast™ chemicals. *Regulatory Toxicology and Pharmacology*, 55, 188-199.
- Kong, A., Frigge, M.E., Masson, G. Et Al. 2012. Rate of de novo mutations and the importance of father's age to disease risk. *Nature*, 488, 471 - 475.
- Lambert, I. B., Singer, T. M., Boucher, S. E., et al. 2005. Detailed review of transgenic rodent mutation assays. *Mutation Research/Reviews in Mutation Research*, 590, 1-280.
- Lewis, S. E. & Simon, L. 2010. Clinical implications of sperm DNA damage. *Hum Fertil (Camb)*, 13, 201-7.
- Lupski, J. R. 2007. Genomic rearrangements and sporadic disease. *Nat Genet*, 39, S43-7.
- Marchetti, F., Aardema, M., Beevers, C., et al. 2018a. Simulation of mouse and rat spermatogenesis to inform genotoxicity testing using OECD test guideline 488. *Mutat Res Genet Toxicol Environ Mutagen*, 832-833, 19-28.

- Marchetti, F., Aardema, M. J., Beevers, C., et al. 2018b. Identifying germ cell mutagens using OECD test guideline 488 (transgenic rodent somatic and germ cell gene mutation assays) and integration with somatic cell testing. *Mutat Res Genet Toxicol Environ Mutagen*, 832-833, 7-18.
- Marchetti, F., Bishop, J. B., Cosentino, L., et al. 2004. Paternally transmitted chromosomal aberrations in mouse zygotes determine their embryonic fate. *Biol Reprod*, 70, 616-24.
- Marchetti, F., Rowan-Carroll, A., Williams, A., et al. 2011. Sidestream tobacco smoke is a male germ cell mutagen. *Proc Natl Acad Sci U S A*, 108, 12811-4.
- Marchetti, F. & Wyrobek, A. J. 2003. PAINT/DAPI analysis of mouse zygotes to detect paternally transmitted chromosomal aberrations. *Adv Exp Med Biol*, 518, 131-45.
- Marchetti, F. & Wyrobek, A. J. 2005. Mechanisms and consequences of paternally-transmitted chromosomal abnormalities. *Birth Defects Res C Embryo Today*, 75, 112-29.
- Oecd 2011. *OECD, Test No. 443: Extended One-Generation Reproductive Toxicity Study*, Paris, France, OECD Publishing.
- Oecd 2020. *OECD, Test No. 488: Transgenic Rodent Somatic and Germ Cell Gene Mutation Assays*, Paris, France, OECD Publishing.
- Oecd 2014. *OECD, Test No. 487: In Vitro Mammalian Cell Micronucleus Test*, Paris, France,, OECD Publishing.
- Oecd 2016a. Test No. 478: Rodent Dominant Lethal Test.
- Oecd 2016b. Test No 478: Rodent Dominant Lethal Test.
- Oecd 2016c. Test No. 483: Mammalian Spermatogonial Chromosomal Aberration Test.
- Oecd 2016d. Test No. 421: Reproduction/Developmental Toxicity Screening Test. *OECD Guidelines for the Testing of Chemicals, Section 4*.
- Oecd 2016e. Test No. 422: Combined Repeated Dose Toxicity Study with the Reproduction/Developmental Toxicity Screening Test. OECD Publishing.
- Oecd 2016d. Test No. 487: In vitro mammalian cell micronucleus test. OECD Publishing.
- Oecd 2016e. Test No. 474: Mammalian Erythrocyte Micronucleus Test. OECD Publishing.
- Oecd 2016f. Test No. 489: In Vivo Mammalian Alkaline Comet Assay. OECD Publishing.
- Oecd 2018a. Test No. 414: Prenatal Development Toxicity Study. OECD Publishing.
- Oecd 2018b. Test No. 443: Extended One-Generation Reproductive Toxicity Study. OECD Publishing.
- Oecd 2018c. Test No. 408: Repeated Dose 90-Day Oral Toxicity Study in Rodents. OECD Publishing.
- Pham, J., Shaw, C., Pursley, A., et al. 2014. Somatic mosaicism detected by exon-targeted, high-resolution aCGH in 10,362 consecutive cases. *Eur J Hum Genet*, 22, 969-78.
- Roach, J. C., Glusman, G., Smit, A. F., et al. 2010. Analysis of genetic inheritance in a family quartet by whole-genome sequencing. *Science*, 328, 636-9.
- Robinson, L., Gallos, I. D., Conner, S. J., et al. 2012. The effect of sperm DNA fragmentation on miscarriage rates: a systematic review and meta-analysis. *Hum Reprod*, 27, 2908-17.
- Sebat, J., Lakshmi, B., Troge, J., et al. 2004. Large-scale copy number polymorphism in the human genome. *Science*, 305, 525-8.
- Shanks, M., Riou, L., Fouchet, P., et al. 2008. Stage-specificity of spontaneous mutation at a tandem repeat DNA locus in the mouse germline. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 641, 58-60.
- Sills, E. S., Fryman, J. T., Perloe, M., et al. 2004. Chromatin fluorescence characteristics and standard semen analysis parameters: correlations observed in andrology testing among 136 males referred for infertility evaluation. *J Obstet Gynaecol*, 24, 74-7.
- Singer, T. M., Lambert, I. B., Williams, A., et al. 2006. Detection of induced male germline mutation: Correlations and comparisons between traditional germline mutation assays, transgenic rodent assays and expanded simple tandem repeat instability assays. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 598, 164-193.

- Speit, G., Vasquez, M. & Hartmann, A. 2009. The comet assay as an indicator test for germ cell genotoxicity. *Mutation Research/Reviews in Mutation Research*, 681, 3-12.
- Stankiewicz, P. & Lupski, J. R. 2010. Structural variation in the human genome and its role in disease. *Annu Rev Med*, 61, 437-55.
- Sun, J. X., Helgason, A., Masson, G., et al. 2012. A direct characterization of human mutation based on microsatellites. *Nat Genet*, 44, 1161-5.
- Vilariño-Güell, C., Smith, A. G. & Dubrova, Y. E. 2003. Germline mutation induction at mouse repeat DNA loci by chemical mutagens. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 526, 63-73.
- Wiszniewska, J., Bi, W., Shaw, C., et al. 2014. Combined array CGH plus SNP genome analyses in a single assay for optimized clinical testing. *Eur J Hum Genet*, 22, 79-87.
- Yauk, C. L., Aardema, M. J., Benthem, J., et al. 2015. Approaches for identifying germ cell mutagens: Report of the 2013 IWGT workshop on germ cell assays(). *Mutat Res Genet Toxicol Environ Mutagen*, 783, 36-54.

Abbreviations used in the document

HTT	Heritable translocation test
SLT	Specific locus test
DLT	Dominant lethal test
TGR	Transgenic rodent mutation assay
CNV	Copy number variant
ESTR	Expanded simple tandem repeat
MN	Micronucleus
SCSA	Sperm chromatin structure assay
TUNEL	Terminal deoxynucleotidyl transferase-mediated (TdT) deoxyuridine triphosphate (dUTP) nick end labeling assay
HTS	High throughput screening
SNV	Single nucleotide variants
WHO/IPCS	World Health Organisation / International Programme on Chemical Safety
GHS	Globally Harmonized System of Classification and Labelling of Chemicals
ICH	International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use