SPI-M-O Medium-Term Projections

17th March 2021

SPI-M-O Medium-term Projections


- These projections are not forecasts or predictions. They represent a scenario in which the trajectory of the epidemic continues to follow the trends that were seen in the data up to 15th March.
- The delay between infection, developing symptoms, the need for hospital care, and death means they will not fully reflect the impact of behavioural changes in the two to three weeks prior to 15th March.
- These projections include the potential impact of vaccinations over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes; it assumes an average of 4.1 million doses are administered per week across the UK.
- The projections assume vaccinations are administered according to JCVI's priority order, with 95% coverage in the over 80s and care home residents; 95% coverage in 70 to 80 year olds; 85% coverage in 50 to 70 year olds and care home workers; and 75% coverage in under 50 year olds.
- Modelling groups have used their expert judgement based on evidence from the <u>JCVI</u>, <u>Public Health England</u>, <u>Scottish universities and Public Health Scotland</u>, as well as other published sources when making assumptions about vaccine effectiveness. A table summarising these assumptions is available in the annex.
- Modelling groups have used data from contact surveys, <u>previous findings</u>, and their own expert judgement to incorporate the impact of re-opening schools. The projections do not include the effects of any other future policy or behavioural changes.
- Some of the projections this week have lower intervals that reach 0 for deaths and hospital admissions. Projecting forwards is difficult when the numbers of cases, admissions and deaths fall to very low levels.
- Not all modelling groups produce projections for both hospitalisations and deaths so there will be some differences between the models included in the combined projections for each metric.

Metrics:

- New hospitalisations per day: Number of individuals admitted with COVID-19 and inpatients newly diagnosed with COVID-19. Data definitions differ slightly across all four nations.
- New deaths per day (by date of death): The number of COVID-19 deaths within 28 days of a positive test. Data definitions differ slightly across all four nations.

New hospital admissions per day

These projections are based on current trends and will not fully reflect the impact of behavioural changes over the past two to three weeks. They are not forecasts or predictions.



Data notes:

England: Number of patients admitted with confirmed COVID-19 and the number of inpatients diagnosed with COVID-19 in the past 24 hours. Taken from NHSE COVID-19 Situation reports.

Wales: Number of patients admitted with confirmed COVID-19 and inpatients diagnosed with COVID-19. Provided by Public Health Wales.

Scotland: Number of patients who tested positive for COVID-19 in the 14 days prior to admission, on the day of admission, or during their stay in hospital. Readmissions within 14 days of a positive test are excluded. Provided by Public Health Scotland.

Northern Ireland: Number of patients admitted with confirmed COVID-19 and inpatients diagnosed with COVID-19. Provided by Health and Social Care Northern Ireland.

Key Real data Expected to Increase Projection Midpoint High and low estimates 5th to 95th percentile High and low estimates 25th to 75th percentile Models


These fan charts show the **90% credible interval and interquartile range** of the combined projections based on current trends. They cannot account for behavioural changes in the two to three weeks prior to 15th March, as these will not yet have been reflected in epidemiological data.


These projections include the potential impact of vaccinations over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes; with 95% coverage in the over 80s and care home residents; 95% coverage in 70 to 80 year olds; 85% coverage in 50 to 70 year olds and care home workers; and 75% coverage in under 50 year olds. The vaccine effectiveness assumptions used by each group are summarised in the annex.

Other than the reopening of schools, these projections do not include any effects of future policy or behavioural changes.

New hospital admissions per day

These projections are based on current trends and will not fully reflect the impact of behavioural changes over the past two to three weeks. They are not forecasts or predictions.






These fan charts show the **90% credible interval and interquartile range** of the combined projections based on current trends. They cannot account for behavioural changes in the two to three weeks prior to 15th March, as these will not yet have been reflected in epidemiological data.

projections include These the potential impact of vaccinations over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes; with 95% coverage in the over 80s and care home residents: 95% coverage in 70 to 80 year olds; 85% coverage in 50 to 70 year olds and care home workers; and 75% coverage in under 50 year olds. The vaccine effectiveness assumptions used by each group are summarised in the annex.


Other than the reopening of schools, these projections do not include any effects of future policy or behavioural changes.

New deaths per day

These projections are based on current trends and will not fully reflect the impact of behavioural changes over the past two to three weeks. They are not forecasts or predictions.



Key



These fan charts show the **90% credible interval and interquartile range** of the combined projections based on current trends. They cannot account for behavioural changes in the two to three weeks prior to 15th March, as these will not yet have been reflected in epidemiological data.

These projections include the potential impact of vaccinations over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for 95% modelling purposes; with coverage in the over 80s and care home residents; 95% coverage in 70 to 80 year olds; 85% coverage in 50 to 70 year olds and care home workers; and 75% coverage in under 50 year olds. The vaccine effectiveness assumptions used by each group are summarised in the annex.

Other than the reopening of schools, these projections do not include any effects of future policy or behavioural changes.


Data Notes:


The number of COVID-19 deaths (by date of death) within 28 days of a positive test.

The past data for England is taken from the PHE line list of deaths. The past data for Scotland, Wales, and Northern Ireland is taken from the Coronavirus (COVID-19) in the UK dashboard on Gov.uk.

New deaths per day

These projections are based on current trends and will not fully reflect the impact of behavioural changes over the past two to three weeks. They are not forecasts or predictions.





These fan charts show the **90% credible interval and interquartile range** of the combined projections based on current trends. They cannot account for behavioural changes in the two to three weeks prior to 15th March, as these will not yet have been reflected in epidemiological data.

projections These include the potential impact of vaccinations over the next three weeks. This has been based on a rollout scenario provided by Cabinet Office for modelling purposes; with 95% coverage in the over 80s and care home residents: 95% coverage in 70 to 80 year olds; 85% coverage in 50 to 70 year olds and care home workers: and 75% coverage in under 50 year olds. The vaccine effectiveness assumptions used by each group are summarised in the annex.

Other than the reopening of schools, these projections do not include any effects of future policy or behavioural changes.

Annex: SPI-M-O Vaccine Effectiveness Assumptions

Table 1: Vaccine reduction in risk of hospitalisation or death											
		LSHTM (Transmission)	Imperial	Manchester [1]	Warwick [2]	PHE	Scottish Government				
Pfizer-BioNTech	1st Dose	88%	86%	75%	75%	80%	88%				
	2nd Dose	94%	98%	75%	90%	95%	95%				
Oxford-AstraZeneca	1st Dose	70%	80%	75%	75%	50%	75%				
	2nd Dose	82%	80%	75%	90%	70%	84%				

Table 2: Vaccine reduction in risk of infection											
		LSHTM (Transmission)	Imperial	Manchester [1]	Warwick [2]	PHE	Scottish Government				
Pfizer-BioNTech	1st Dose	48%	65%	75%	60%	48%	48%				
	2nd Dose	60%	94%	75%	70%	60%	60%				
Oxford-AstraZeneca	1st Dose	48%	63%	75%	60%	48%	48%				
	2nd Dose	60%	63%	75%	70%	60%	60%				

[1] Manchester's model doesn't split vaccine effectiveness by vaccine type or different doses.

[2] Warwick's vaccine effectiveness assumptions are based on a weighted average of the two vaccines.