



Climate Change and Agricultural Commodities

CABI Position Paper

August 2009

Greg Masters g.masters@cabi.org

Peter Baker p.baker@cabi.org

Julie Flood j.flood@cabi.org

KNOWLEDGE FOR LIFE



Table of Contents

Table of Contents.....	2
Table of Figures	Error! Bookmark not defined.
Table of Boxes	Error! Bookmark not defined.
Executive Summary	4
1. Background	8
2. Review of climate change impacts	10
3. Climate change impacts on agricultural production	15
3.1 Effects of increasing temperatures	20
3.2 Changing precipitation patterns	23
3.3 Effects of increasing atmospheric CO ₂	23
3.4 Elevated CO ₂ , temperature and precipitation interactions	24
3.5 More frequent and extreme weather events	24
3.6 Changes in sea level.....	25
4. CABI Member Country Focus: India, agriculture and climate change	26
4.1 Climate and future climate of India	26
4.2 Impact of climate change on Indian agriculture	26
4.3 Implications of climate change impacts on Indian agriculture	28
4.4 What is CABI doing?	28
5. CABI Member Country Focus: Ghana, agriculture and climate change	30
5.1 Climate and future climate of Ghana	30
5.2 Impact of climate change on Ghanaian agriculture	30
5.3 Implications of climate change impacts on Ghanaian agriculture.....	31
5.4 What is CABI doing?	32
6. Commodity Focus: Cotton	33
6.1 Impacts of climate change on cotton	33
6.2 Pakistan, climate change and cotton	33
6.3 What is CABI doing?	34
7. CABI Commodity Focus: Coffee	35
7.1 Observed impacts of climate change on coffee.....	35
7.2 Impacts of climate change on coffee	35
7.3 What is CABI doing?	36
8. Summary and knowledge gaps	38
8.1 Immediate research needs	38
8.2 Priorities	39
8.3 Adaptation	40
9. References and further reading	42
contact CABI	46

Figures

Figure 1 Probability that projected average summer temperatures for (A) 2050 and (B) 2100 will be greater than the highest temperature recorded (1900-2006).	12
Figure 2 The impact of climate on water availability and then combined on agriculture (crop yield) ..	17
Figure 3 Agriculture and rainfall tendencies for eastern and southern Africa: Food-security analysis - rainfall, population, cropped area, seed use, and fertilizer use	18
Figure 4 Projected changes in global agricultural productivity by 2080 due to climate change (and carbon fertilization).....	19
Figure 5 Potential impacts of climate change on agricultural crops.	19
Figure 6 Comparison of temperature rise on crop yields in temperate and tropical regions	22
Figure 7 Projected impact of climate change on wheat: production (total yield in tons) estimates for India	27
Figure 8 Coffee exports for Africa and Vietnam.....	37



Boxes

Box 1: Observed and projected climate changes	7
Box 2: Ecosystem resistance, resilience and services and the link with sustainable development.	7
Box 3: Observed and projected climate changes (IPCC, 2007)	9
Box 4: Climate change impacts based on projections to the mid-late 21 st Century	11
Box 5: key impacts as a function of increasing global average temperature change.....	12
Box 6: Temperature relations for wheat grain.....	19
Box 7: Global warming threatens coffee collapse in Uganda	35
Box 8: Projected decline in production of coffee in Sao Paulo, Brazil	35
Box 9: The lesson from Brazil: government intervention works.....	39



Crop yields and changes in productivity due to climate change will vary considerably across regions and among localities, thus changing the patterns of production – in general, productivity is projected to increase in middle to high latitudes, depending on crop type, growing season, changes in temperature regime, and seasonality of precipitation

Robert T Watson

Former Chairman Intergovernmental Panel on Climate Change, 2001

...the cost of inaction will be great...

Nicholas Stern

Former UK Government Advisor and Director of Policy and Research for the Commission for Africa

It is irresponsible, reckless and deeply immoral to question the seriousness of the situation. The time for diagnosis is over and the time for action is now

Mrs. Gro Harlem Brundtland

UN Special Envoy of the Secretary General on Climate Change

Executive Summary

Agricultural commodities are any agricultural products which are traded internationally in response to demand. Climate change will have a profound impact on agricultural commodities including on their production and their productivity with consequences to both food supply and food security and many of these effects will be seen most in Least Developed Countries (LDCs).

Without doubt, climate change is occurring and is already having a dramatic impact on climatic variability, global temperatures and sea level. Climate change will have significant impacts on agriculture, reflecting the close link between climate (temperature and precipitation in particular) and productivity and these effects are likely to have greatest effect in the LDCs of the tropical zones where productivity will decrease. A greater frequency of extreme events, heat stress, droughts and floods, will increasingly have negative impacts on crop yields. In particular, water scarcity and the timing of water availability will increasingly constrain production. Climate change will increase pest and disease outbreaks thus negatively affecting food production. Rising carbon dioxide (CO₂) levels will also have effects, both detrimental and beneficial, on crop yields. The ability of farmers and rural societies to adapt to these changes is crucial to maintaining an adequate global food supply. While demand is likely to increase, due to rising global population size primarily, climate change will challenge agricultural production and food security (locality of production, supply, volume, quality). By 2080, agriculture output in Least Developed Countries (LDCs) may decline by 20% due to climate change and yields could decrease by 15% on average.

Immediate research needs

There is an urgent need to do more research on the collection and synthesis of data for regions and for individual countries so that ever more accurate models are produced to inform decision makers about the strategies they need to adopt. There is a need for more action on adaptation measures in agriculture, especially in LDCs, to implement these proposed strategies so assisting farmers, extensionists, researchers and stakeholders along the supply chain to reduce the adverse effects of climate change. This is particularly important in regions where there is already large, resource-poor populations suffering from food insecurity. In the immediate future, agricultural production needs to be made more sustainable and thus, resilient to change. To achieve this, increased knowledge and application of new irrigation technologies, water conservation measures, drought resistant varieties, new cropping patterns, improved management of pests and diseases through integrated pest management, crop diversification particularly with alternative crops, all have to be researched and implemented urgently.



Due to the complex physical and biophysical impacts of the interactions between the different climate variables, combined with varying irrigation, fertilizer and pest and disease management strategies and techniques, regional variability (e.g. topography, soils, land-use and management), and differences in crop growth, development, phenology, yield and resistance to drought and pest and diseases; the detailed impacts of these factors need to be investigated further. Conserving and promoting ecosystem resistance and resilience has to be a central tenet of adaptation strategies and needs to be balanced with any actions to alter or intensify agricultural production.

Specific recommendations for further research include:

- Increased precision in climate change projection with higher resolution on spatial and temporal scales;
- Linking of projections with agricultural production systems to suggest suitable options for sustaining agricultural production;
- Preparation of a database on climate change impacts on agriculture;
- Evaluation of the impacts of climate change in selected locations;

In terms of experimentation, there is a lack of knowledge of CO₂ and climate responses for the majority of crops other than cereal staples such as wheat, maize and rice, including many of importance to the rural poor (e.g. root crops, millet, brassicas). In addition to the staples such as root crops and cereals, more work needs to be initiated on how climate change will impact on traded commodities such as coffee, cotton and cocoa and how changes in production area, changes in yield etc will impact on the foreign exchange earnings of LDCs in particular those whose economies are based on one or two of these traded commodities. Quality of these traded commodities is also projected to change as well as quantity and improvement in post harvest technologies as pioneered in coffee in Africa by CABI and partners could be one way to limit the impact of changes in quality. The response of commodity pest and diseases to climate change is a knowledge gap and CABI's expertise and approach can address this research and implementation need. Changes in climate could allow endemic pests and diseases to increase both in severity and could provide conducive conditions to allow pests and diseases to expand their geographical range.

Where datasets exist (public/ private) of climate and preferably yield, of 50 years or greater then there is an opportunity to conduct an analysis and project changes for the near future (next 20 years). These projections should include, if possible, changes in mean, median, standard deviations, variance of temperatures and rainfall, but also the distribution of wet spells, dry spells, absolute maxima and minima, frequency of storms, frosts, and droughts. Valuable information can be gained to aid farmers in what to expect in the immediate future, for example dates of first and last frost, variation in rainfall patterns. This approach will enable future climate information and forecasts and thus better management of crops and cropping systems. CABI with member countries can source such datasets and run analyses. Results and conclusions can be disseminated by various means including via the CABI Environmental Change Web Portal.

Prioritisation

- **Awareness raising and training on planning and zoning** needed for donors and decision makers. Governments need to take firm action to protect the fertile uplands of their countries. These resources will become ever more valuable as climate change forces upslope migration of flora, fauna and human communities.
- **Capacity building** to reinvigorate institutions, including extension and research divisions. In general, agriculture has stagnated in development over the last 30 years, although recently, perhaps in response to environmental change and the growing threat of food insecurity and starvation, there are signs of this changing.
- **Adaptation planning and implementation:** in the short to medium term there is much that can be done to help farmers adapt to climate change. In areas of increased drought, options and knowledge exists to start activities in water harvesting, improved irrigation (e.g. drip irrigation,



being used for coffee in Nicaragua), drought resistant varieties, use of shade trees where appropriate, lean water use in processing, minimum tillage and many other technologies.

- **Coordination of the efforts of Governments, private organisations and other stakeholders** in mitigating or adapting to the challenges of climate change.

Prevention of climate change is not possible, mitigation can “buy-time” but adaptation strategies are needed for long term sustainability.

Adaptation

The United Nations Framework Convention on Climate Change (UNFCCC) instigated National Adaptation Programmes of Action (NAPAs) to provide a process for LDCs to identify priority activities that respond to their urgent and immediate needs to adapt to climate change – those for which further delay would increase vulnerability and/or costs at a later stage. Key strategies will include:-

- Use of new/ improved varieties, crops, or rotations (longer maturing varieties, heat and drought-tolerant). Alternatively, switching production from an increasingly unsuitable area to a climatic zone which is projected to be favourable for the commodity in the future;
- Design and implement irrigation and drainage;
- Adaptation to changed growing season or to shift timing of heat stress;
- Design and implement flexible cropping systems with mixed varieties, as trialled and implemented for commodities such as coffee;
- Improved use of sustainable technologies, including integrated pest management, post harvest management and soil health, of which CABI has expertise and programmes.

However, in developing adaptation strategies, agriculturists and other stakeholders need to ensure they are not adding to climate change. Agriculture, particularly intensive production, produces approximately a quarter of global greenhouse emissions. Given predicted decreases in agricultural production and an increasing demand for food, there is an argument to intensify and expand to meet the demand. This is a real challenge as it needs to be balanced against the need to adapt to climate change and to attempt to stabilize greenhouse gas emissions. Conserving and promoting ecosystem resistance and resilience has to be a central tenet of adaptation strategies and needs to be balanced with any actions to alter or intensify agricultural production.

Capacity building underpins much of foregoing discussion. Capacity building for all stakeholder groups will enable them to better deal with climate change in their own locality, country or region. Much of the focus should enable agriculturists to respond to current climatic variability, thus providing the stepping stones to adapt to future, more extreme climatic variability.

CABI (www.cabi.org) is a not-for-profit intergovernmental organization owned by a consortium of over 40 member countries and is dedicated to improving people's lives worldwide by providing information and applying scientific expertise to solve problems in agriculture and the environment. In 2007- 08, CABI organised three regional consultations with its member countries in Latin America and the Caribbean, Africa, and Asia. As a result, CABI has a mandate to support its member countries in adapting to climate change and in developing management strategies and policies. CABI's country membership, combined with a depth of practical on-the-ground experience plus our information and knowledge bases, puts us in a unique position to catalyse action by informing, up-dating and generally building capacity for countries, regions, communities and individuals to take concerted action now to adapt to the new realities of the impacts of climate change.

CABI has considerable, internationally recognised expertise among the staff in the field of agricultural commodities including coffee, cocoa, cotton, oil seeds. The unique combination of in-house experts in publishing, abstracting, databases, plus in-country scientists and practitioners allows us to deliver knowledge direct to those who need it web based portals and e-learning packages. Some of our initiatives in LDCs involve improved post-harvest processing for coffee and cocoa, integrated pest management for many pests and diseases of our focus commodities while community based



programmes such as our farmer field schools although originally used for IPM have also been adapted by us to improve quality and water management for cotton production.



1. Background

Climate change will impact global agriculture in terms of productivity, commodities (Box 1) and food security. Agriculture affects all livelihoods, occupying approximately 40% of the land globally, consuming 70% of global water resources and affecting biodiversity at all scales; genetic to ecosystem. One of our biggest challenges is how to balance the growing demand for agricultural commodities with an agricultural resource that is increasingly under pressure. Agricultural resources and their associated supply chains need to be managed sustainably, and climate change is a major challenge to sustainable development (Box 2).

Food security is a global issue that links with many aspects of food production, distribution, the environment and socio-economics. An estimated 2.5 billion people in Developing Countries depend on agriculture for their livelihoods (DfID, 2005). More than 50 Developing Countries, including a majority of the Least Developed Countries (LDCs), depend on exports of three or fewer agricultural commodities, typically tropical products, for between 20 and 90% of their foreign exchange earnings. Many LDCs are also net food importers, spending more than half their export earnings on commodity markets purchasing food imports to make up for shortfalls in domestic production. For example, 63% of all agricultural earnings from export in Ghana is due to cocoa (Section 5) and this single commodity constitutes over 70% of a farm household income. Changes to international commodity markets can make the difference between famine, survival or surplus.

Climate change will undoubtedly lead to more pressure on already volatile commodity prices and trading. The general consensus is that in many regions especially those in the tropics, agricultural productivity will decrease and that food shortages will increase (Section 3). Additionally, not only will climate potentially decrease the amount of land available for agricultural production but, there will be increased competition for resources with other development needs, such as infrastructure. Food and commodity prices have risen, and are expected to increase further due to instability in production quality and quantity due to climate change impacts.

If LDCs are to adapt to climate change they need to develop resilience by reducing long term vulnerability. Intercropping and increasing biodiversity, using drought tolerant varieties, utilising surrounding ecosystems and their services, implementing sustainable production, including integrated crop and pest management and targeted (or alternative) irrigation and fertiliser application are just some immediate strategies that can be used. If such knowledge is applied through a regional or local land management scheme developed and supported by extensionists, then resilience will naturally be increased. With an increasing demand for food, a predicted decrease in food production and a subsequent increase in food shortages, famine and poverty, there appears to be no alternative to the short term policy of importing food. Food insecurity will only get worse under climate change, threatening many millions of people in Developing Countries. Policies promoting the diversification of production in terms of number of commodities grown are simple risk-spreading strategies that could enable short-term adaptation to climate change.

Over the past 50 years, Man's activity has changed all ecosystems more rapidly and extensively than in any comparable period of time. The demand for ecosystem services (Box 2) grew significantly as the world population doubled to 6 billion people and the global economy increased more than six fold. To meet this growing economic and population demand, food production has increased by roughly 2.5 times and water use more than doubled.

Climate change will have a rapid and significant impact on the agricultural systems and the natural environment. Established production systems are becoming increasingly destabilised and environmental stress, particularly water stress, is growing in parallel with rising demand for food. It has been estimated that China will lose self-sufficiency in food production by 2030 due to climate change (Erda *et al.* 2008). The drive towards a select few varieties and crop species, results in genetic erosion. Such changes are great challenges to biodiversity which is the raw material used by breeders to create new crop/livestock varieties that will be needed to safeguard food and agriculture



for future generations as well as maintaining a broad gene pool, ensuring ecosystem resilience and resistance (Box 2). Genetically modified crops whilst useful in management of some commodity crops, such as cotton, has been a driver for much biodiversity loss as farmers opt for the new varieties over the older varieties and traditional landraces disappear.

Climate is central to agricultural production, it follows that as climatic parameters change then so will agriculture. This paper reviews climate change before considering the impact of climate change on some aspects of agricultural production. Focus case studies provide additional analysis and synthesis for the main agricultural commodities within CABI Member Countries, India and Ghana (particularly cocoa); other specific commodities include cotton (with a focus on Pakistan) and coffee. The final analysis identifies knowledge gaps.

Box 1: Agricultural Commodities

Definition

- Any item that is produced for which there is a demand and therefore a market.
- Any item that can be openly traded within such market.
- Agricultural commodity means: grain, livestock, poultry, fruit, timber or other items produced from agricultural activities.
- One of the characteristics of a commodity is that the price is determined as a function of the market as a whole. Established commodities have actively traded *spot* and *derivative* markets.
- Important agricultural commodities, according to the OECD-FAO (2008) are: wheat, rice, coffee, tea, maize, yams, cassava, sorghum, pulses, sugar cane, sugar beet, potatoes, fruit, vegetables (beans).
- CABI has internationally recognised expertise in a range of commodities including coffee, cotton, cocoa, oil seeds, wheat, and bananas

Box 2: Ecosystem resistance, resilience and services and the link with sustainable development

Every ecosystem is subject to perturbations such as climate, nutrient fluctuation, loss of biodiversity, and introduction of exotic species, that can alter ecosystem structure and function. The degree to which ecosystems respond to perturbations depends on their ability to withstand perturbation and maintain normal function (resistance) and/or recover from disturbance (resilience). The dynamic nature of ecosystems endows them with a degree of resilience to change. These two components of stability, in essence, address how a system responds to disturbances and knowledge about them is key to developing ecosystem recovery and restoration efforts.

We must learn to use ecosystem resources (services), without compromising the ability of ecosystems to maintain their integrity (structure and function) especially through understanding how ecosystems respond to perturbation (resilience/resistance). This is the core concept behind Sustainable Development.

Sustainable Development is development which meets the needs of the present without compromising the ability of future generations to meet their own needs (World Commission on Environment and Development, 1987).



2. Review of climate change impacts

Climate has changed many times in response to a variety of natural causes but the term 'climate change' usually refers to those changes that have been observed since the early 1900s and includes anthropogenic and natural drivers of climate. The main human influence on global climate is through emissions of greenhouse gases such as carbon dioxide (CO₂) and methane (CH₄). At present, about 6.5 billion tonnes of CO₂ are emitted globally each year, mostly through burning fossil fuels. Changes in land-use mean a further net annual emission of 1-2 billion tonnes of CO₂. Such increasing concentrations of greenhouse gases in the atmosphere since the industrial revolution have trapped more energy in the lower atmosphere, altering global climate.

The observed changes in global climate (Box 3) are due to a combination of both natural and anthropogenic causes. The Intergovernmental Panel on Climate Change (IPCC) concluded that the balance of evidence, from changes in global mean surface air temperature and from changes in geographical, seasonal and vertical patterns of atmospheric temperature, suggests a discernible human influence on global climate. There are uncertainties in key factors, including the magnitude and patterns of long-term natural variability (IPCC, 2001). The latest IPCC projections suggest anthropogenic climate change continuing at an alarming rate, with an average global warming by 2100 of up to 6.4°C (Box 3; Fig. 1; IPCC, 2007). These projections indicate that the warming would vary by region, and be accompanied by increases and decreases in precipitation. In addition, there would be changes in the variability of climate, and changes in the frequency and intensity of some extreme climate phenomena.

Already there is evidence indicating climate change impacts on physical and biological systems across the globe. Examples include glacial retreat and shrinkage, permafrost thawing, lengthening of growing seasons particularly at mid to high latitudes, poleward and altitudinal shifts in organism distribution and phenological shifts (e.g. earlier budburst). Generally, changes in biological and physical systems were in line with the expected change for directional climate change (Box 4; 5). The probability that the observed changes in the expected direction (with no reference to magnitude) could occur by chance alone is negligible and there is high confidence levels that recent changes in temperature are having discernible impacts on physical and biological systems (Box 4; 5).

Climate change therefore threatens economic development in many countries but this effect may be of particular consequence in the tropical climatic zone where many Least Developing Countries (LDCs) are situated (Fig 1) and where climatic variability is already a significant challenge to poverty alleviation. It is essential that key indicators of the onset of climate change impacts are developed and monitored, with appropriate thresholds set to establish not only when action should take place but what type of strategy should be adopted. Considering the Millennium Development Goals as a framework on which to base indicators (<http://mdgs.un.org/unsd/mdg/Default.aspx>; United Nations, 2008), LDCs in the tropical zone require help in developing appropriate indicators, such as measures of poverty reduction, economic development, sustainability, ecosystem impact, status of threatened species, and reform of trade and investment policies. The impacts of climate change on agricultural production are considered further in Section 3.

However, agriculture and food distribution also affect climate change, principally through the production and release of greenhouse gases such as carbon dioxide, methane and nitrous oxide (approximately 18% of the UK greenhouse gas emissions come from industrial agriculture and the food distribution system) but also by altering land cover, which changes the absorption or reflectance of heat and light, thus contributing to radiative forcing (key process of the greenhouse effect). Deforestation and desertification, together with fossil fuel use are major anthropogenic sources of carbon dioxide; agriculture itself is a major contributor to increasing atmospheric methane and nitrous oxide concentrations. Agriculture energy use (producing CO₂) includes the storage, drying, cooling and processing of food crops, the production of fertilizers and other agricultural inputs as well as feed concentrates and the need for intensive grazing pasture. Livestock and manure produce methane, a potent greenhouse gas. Greenhouse production and indoor livestock are also energy intensive.



The consequences of climate change are far reaching and will affect all sectors of society. The more that is learnt about climate change impacts, the more complicated the issue becomes but we have attempted to summarise the likely impacts of climate change on agricultural production in the next section.

Box 3: Observed and projected climate changes (IPCC, 2007)

Observed

- Global average surface temperature increased over the 20th Century by $0.74 \pm 0.18^{\circ}\text{C}$
- 11 of the last 12 years (1995-2006) rank among the 12 warmest years since instrumental records began (1850)
- In the past 50 years, cold days, cold nights and frosts have become less frequent while hot days, hot nights and heat waves have become more frequent
- Snow and ice extent, including mountain glaciers and snow cover have decreased, consistent with the warming pattern
- The area affected by drought has increased since the 1970s
- Precipitation has increased in eastern parts of North and South America, northern Europe and northern and central Asia but has declined in the Sahel, the Mediterranean, southern Africa and parts of southern Asia

Predicted

- For the next two decades there is likely to be warming of 0.2°C per decade
- The projected rate of warming is greater than any observed during the 20th Century
- Global average surface warming will increase by 1.1 to 6.4°C by 2100 relative to 1980–1999
- Global average water vapour concentration and precipitation increase during the 21st Century
- Global mean sea level will rise by 18 to 59 cm between 1990 and 2100
- Increase in Asian summer monsoon precipitation variability




Figure 1 Probability that projected average summer temperatures for (A) 2040–60 and (B) 2080–2100 will be greater than the highest temperature recorded (1900–2006). The areas shown in red have a greater than 90% probability that the future average summer temperature will exceed the highest temperature on record (from Battisti & Naylor, 2009).



Box 4: Climate change impacts based on projections to the mid-late 21st Century. These do not take into account any changes or developments in adaptive capacity. (IPCC, 2007)

Phenomenon and direction of trend	Likelihood of future trends based on projections for 21 st Century using SRES scenarios	Examples of major projected impacts by Sector			
		Agriculture, forestry and ecosystems	Water resources	Human health	Industry, environment and society
Over most land areas; warmer and fewer cold days and nights; warmer and more frequent hot days and nights	Virtually certain	Increased yields in colder environments; decreased yields in warmer environments; increased insect outbreaks	Effects on water resources relying on snow melt; effects on some water supplies	Reduced human mortality from decreased cold exposure	Reduced energy demand for heating; increased demand for cooling; declining air quality in cities; reduced disruption to transport due to snow, ice; effects on winter tourism
Warm spells/heat waves. Frequency increases over most land areas	Very likely	Reduced yields in warmer regions due to heat stress; increased danger of wildfire	Increased water demand: water quality problems e.g. algal blooms	Increased risk of heat-related mortality, especially for the elderly, very young, chronically sick and socially-isolated	Reduction of quality of life for people in warm areas without appropriate housing; impacts on the elderly, very young and poor
Heavy precipitation events. Frequency increases over most land areas	Very likely	Damage to crops; soil erosion; inability to cultivate land due to water logging of soils	Adverse effects on surface and groundwater; contamination of water supply; water scarcity may be relieved	Increased risk of deaths, injuries and infectious, respiratory and skin diseases	Disruption of settlements, commerce, transport and societies due to flooding; pressure on rural and urban infrastructures; loss of property
Area affected by drought increases	Likely	Land degradation; lower yields/crop damage and failure; increased livestock death; increased risk of wildfire	More widespread water stress	Increased risk of food and water shortage; increased risk of malnutrition; increased risk of water- and food-borne diseases	Water shortages for settlements, industry and societies; reduced hydropower generation potentials; potential for population migration
Intense tropical cyclone activity increases	Likely	Damage to crops; windthrow (up-rooting) of trees; damage to coral reefs	Power outages causing disruption of public water supply	Increased risk of deaths, injuries, water- and food-borne diseases; post-traumatic stress disorders	Disruption by flood and high winds; withdrawal of risk coverage in vulnerable areas by private insurers; potential for population migrations; loss of property
Increased incidence of extreme high sea level (excludes tsunamis)	Likely	Salinisation of irrigation water, estuaries and freshwater systems	Decreased freshwater availability due to salt water intrusion	Increased risk of deaths and injuries by drowning in floods; migration related health effects	Costs of coastal protection versus costs of land-use relocation; potential for movement of populations and infrastructure; also see <i>tropical cyclones</i> above



Box 5: Key impacts as a function of increasing global average temperature change.




Illustration of global impacts projected by climate changes associated with different amounts of global average temperature increase in the 21st Century. The black lines link impacts, the dotted lines indicate impacts continuing with increasing temperature. Entries are placed so that the left hand side of the text indicates the approximate area of a given impact. All sectors, water, ecosystems (natural and managed), food, coasts and health will be compromised and changed due to climate change. The impacts illustrated are those with the greatest links, either as a driver or as a consequence, to agricultural production. For example, there is widespread malnutrition projected but agricultural production will be further compromised, the ecosystems around cereal fields will experience changes in structure and function thus compromising functional biodiversity and the ability of farmers to use natural control strategies to deal with pests and water shortages are likely. Hence, increased food production to counter malnutrition is unlikely unless novel adaptation strategies are developed and implemented urgently. Adaptation to climate change is not included in these estimations. All entries are from published studies recorded in the chapters of the IPCC Working Group II Assessment. (Source: Modified from the IPCC, 2007).



3. Climate change impacts on agricultural production

Climate change will impact on agricultural commodity production globally due to the effects on plant growth and yield from elevated CO₂, higher temperatures, altered precipitation and transpiration regimes, and increased frequency of extreme events, as well as modified weed, pest and pathogen pressure.

Although the current green revolution in agricultural production has led to many technological advances such as improved varieties, GMOs, physiological improvement, seed treatment, targeted irrigation and fertilizer applications and integrated pest management, the local climate is still the greatest factor in determining how much food is produced in a given region or locality (Fig. 2). Even with these man-made improvements, temperature and precipitation relations are major factors in determining where a crop can grow, whether there is any yield, how many crops can be grown per year and what type of crop can be grown in a certain location. Seed viability, germination and plant growth, development, stature, phenology, fruiting and seed mass are all governed by temperature, water (precipitation/ soil moisture) and the interaction between these two climatic factors. Indeed, a “healthy” plant, which has just the right ratio of temperature and water for its optimum development is generally more resistant to attack by pests and diseases (conversely, stressed plants are more susceptible to attack). Crop growth models, used by agriculturists and agronomists, are climate dependent and so increasingly are being linked or incorporated into climate change models and general circulation models (GCMs).

Such relations underlie the susceptibility of agricultural production to climate change (Fig. 2). Climate is the prime driver of agricultural production and so, climate change will have a significant impact on agricultural commodities. This impact is projected to be positive and negative, depending on the severity of the climatic changes, the agricultural system, locality and capacity to adapt. In recent decades, international trade in agricultural commodities has grown rapidly, and so in addition to climate change affecting agriculture locally or at the regional scale, the impact of climate change on a global scale has to be considered. Certain crops are favoured by trade, to the extent that over 70% of the world’s food comes from just nine crops (rice, wheat, maize, oats, potato, barley, cassava, soybean, sugar cane), each of which is now cultivated far beyond the natural range to maximise income.

Farmers everywhere naturally manage their systems for climatic variability. Climate change, though, will have a sustained and increasing impact on agricultural systems, through increasing temperature, increasing greenhouse gases, and probably most importantly, increasing variability in frequency and severity of climatic events, particularly extreme weather events. Farmers generally use historic records to aid in planning what, when and where to plant or graze, but without accurate, reliable regional forecasts of climate change they are likely to experience greater pressure on production, as timing, location and crop choice are likely to be wrong or at least sub-optimal. In many places, growing seasons are changing, ecological niches are shifting, and rainfall is becoming more unpredictable and unreliable both in its timing and its volume (rainfall projection; Fig. 3). This is leading to greater uncertainty and heightened risks for farmers, and potentially eroding the value of traditional agricultural knowledge such as when to plant particular crops. Accurate predictions are essential to developing adaptation strategies (e.g. introducing a new crop more suited to the future climate) and implementing mitigation strategies if there is time to do so.

Climate change will have a range of impacts on agricultural systems and production (Fig. 4, 5). It is likely that the life cycle of crops will progress (advanced phenology) more rapidly; but with rising temperatures and variable rainfall, crops are expected to fail, especially if there is a decrease or increased variability in precipitation. However, the situation is more complicated; gains and losses are predicted for crop yields, depending on locality, crop type and production system (Fig. 4, 5). Global warming in the short-term is likely to favour agricultural production in temperate regions (largely northern Europe, parts of North America) and negatively impact tropical crop production (south Asia, Africa), where the majority of LDCs are located. For example, southern Africa has been



projected to lose over 30% maize yield by 2030 (Lobell *et al.* 2008). Additionally, over 10% losses of important staple crops such as maize, rice and millet are projected for South Asia within the same time period. Even small changes in temperature and rainfall could have significant effect on the quality of cereals, fibre and beverage crops, fruits, aromatic, and medicinal plants with resultant implications on their prices and trade.

The population of undernourished people in eastern and southern Africa has doubled in the last 25 years, often this is coupled with the cessation of development in rural areas (including agricultural capacity) and rural poverty increasing in the last 10–15 years. During the same time period, this region and the LDCs/ food insecure countries of the western rim of the Indian Ocean have been subjected to a 15% reduction in rain during the growing season. Such drought and the rural social disruption in food insecure countries are projected to lead to a 50% increase in the number of undernourished people by 2030. These relationships between rainfall, population, cultivated area, seed and fertiliser use are shown in Fig. 3, which also shows how mitigating or adapting to one of these components may offset the effects of the others. For example, modest increases in per capita agricultural productivity could more than offset the observed precipitation declines. Consequently, investing in agricultural development can help mitigate climate change while decreasing rural poverty and vulnerability.

Although general projections of the impacts of climate change are useful in some respects, they are subject to some uncertainty as it is the climate change model adopted for the analysis that can have a significant effect on the outcome. Therefore, recently crop models have been integrated with General Circulation Models (GCMs) for a range of increased CO₂ climate scenarios (Easterling *et al.* 2007; Hansen, 2005; http://www.adaptation.nrcan.gc.ca/perspective/summary_5_e.php) in an attempt to remove these uncertainties and this approach is proving to give consistent projections. Such analyses indicate that understanding the water cycle and how it is affected by temperature in particular is key in terms of agricultural production.

Increased moisture stress and drought are major concerns for both irrigated and non-irrigated crops. If adequate water is not available, production declines and entire harvests can be lost. While climate change is expected to cause moisture patterns to shift, there is still considerable uncertainty concerning the magnitude and direction of such changes. Longer growing seasons and higher temperatures are expected to increase demand for water, as would changes in the frequency of drought. In Africa and Latin America many rain-fed crops are near their maximum temperature tolerance, so that yields are likely to fall sharply for even small climate changes; falls in agricultural productivity of up to 30% over the 21st century are projected (Easterling *et al.* 2007). Water stress, increased temperature, the impact of an increased incidence and abundance of new invasive pests, diseases and weeds are almost certainly major factors in the projected agricultural declines in the tropics (Fig. 5). Invasive insect, disease and weed pests are likely to benefit most from climate change, leading to increased pesticide and herbicide use or greater reductions in yield.




Figure 2 The impact of climate on water availability and then their combined impact on agriculture (crop yield).

GDD: growing degree days; GSL: Growth season length. Adapted from M.A. Goheer, Global Change Impact Studies Centre (GCISC), Pakistan.




Figure 3 Agriculture and rainfall tendencies for eastern and southern Africa: Food-security analysis - rainfall, population, cropped area, seed use and fertilizer use

Rainfall is shown as the percentage of the 30 year average for 1951–1980. Other variables are expressed as percentages of averages for 1979–1981. Global radiative forcing is shown by the broken line. (Adapted from Funk *et al.* 2008; data originally from the FAO)




Figure 4 Projected changes in global agricultural productivity by 2080 due to climate change (and carbon fertilization).

Red areas indicate a decrease in productivity and green areas indicate an increase. Tropical areas are generally expected to have biggest decreases in agricultural production, which correlates to many LDCs. Consequently, the poor will be affected and will become poorer. UNEP/GRID-Arendal, Projected agriculture in 2080 due to climate change, UNEP/GRID-Arendal Maps and Graphics Library, <http://maps.grida.no/go/graphic/projected-agriculture-in-2080-due-to-climate-change>




Figure 5 Potential impacts of climate change on agricultural crops.

Climate promoted invasive species impacts (pests, diseases and weeds) are expected to be increasingly detrimental. http://www.adaptation.nrcan.gc.ca/perspective/summary_5_e.php



3.1 Effects of increasing temperatures

As a generality (Easterling *et al.* 2007) global food production is projected to increase overall with increases in average (local) temperatures of 1 – 3°C, but if temperatures increase above 3°C then global food production will decrease.

Temperature is fundamental in determining crop quality, quantity and where it can be grown. Due to the nature of this fundamental relationship in biology and ecology (Fig. 2), any changes in temperature, due to climate change, will have large impacts on crop (commodity) production.

All stages of crop development are sensitive to temperature. Development generally accelerates in a linear fashion within certain temperature boundaries (e.g. between 10 and 30°C for wheat; see Box 6), but with extreme temperatures the relationship becomes non-linear and increasingly difficult to predict. Higher temperatures often lead to heat stress which can result in increasing sterility and lowering overall productivity in crops. As temperature rises there is increased evaporation from plants and soils resulting in increased water requirements while lowering water availability, which causes further stress to the crop.

Chronic effects of warmer temperatures on crop growth and development are probably more important in projecting climate change impacts, than understanding extreme effects. Crop yields reflect the importance of season-long effects, where crops generally have a greater yield when the temperature is cooler during growth of the harvested component. The length of crop growth cycles are temperature dependent. Increased temperature leads to the time between sowing and harvesting being shorter (e.g. for maize this could be between one and four weeks shorter). This could be beneficial in terms of a greater number of cropping cycles, thus increasing yield over the course of a year. However, such a reduction in the duration of the crop cycle could lead to earlier senescence thus having an adverse effect on productivity. Additionally, a greater number of cropping cycles will lead to greater inputs, resulting in soil nutrient and moisture depletion, greater exposure (apparency) to pests and greater pressure on what might already be marginal land.

Temperate zones

In Europe, temperature increases will lead to a poleward migration of suitable cropping areas and a reduction in the growing period of determinate crops (e.g. cereals), but an increase for indeterminate crops (e.g. root crops). In northern Europe climate change (temperature change predominantly) is expected to have positive implications for agriculture, with the introduction of new crop species and varieties, higher crop production and expansion of suitable areas for crop cultivation. Disadvantages may be a greater risk from pests and disease and so greater plant protection needs, an increased risk of nutrient leaching and turnover of soil organic matter. In southern areas of Europe, the disadvantages will predominate. The possible increase in water shortage and extreme weather events may cause lower harvestable yields, higher yield variability and a reduction in suitable areas for traditional crops. These effects may reinforce the current trends of intensification of agriculture in northern and western Europe and extensification in the Mediterranean and south-eastern areas of Europe.

Mild warming is projected to result in initial increases in crop yields in some other temperate regions (Fig. 4). For example, as highlighted earlier, when crop models have been integrated with GCMs (multi factor climate projections) and applied to Canada it has been demonstrated that maize, sorghum, soybean and wheat yields are expected to increase by 20 – 124% under climate change, due to the projected temperature increases and increases in CO₂. Nevertheless, once temperature increases exceed 3°C, yields decline. Likewise, yields of sunflowers, potatoes, tobacco and sugarbeet will initially increase while peas, onions, tomatoes and cabbage will suffer a decrease in yields.

Increase in temperature will lengthen the effective growing season in areas where agricultural potential is currently limited by cold temperature stress. Thus, increased temperature will cause a



poleward shift of the thermal limits to agriculture. This poleward shift will be especially important for crops such as rice that have tropical centres of origin and adaptation but are also grown in temperate latitudes during warm seasons. Global warming impact will be greater in the northern hemisphere than the southern because there is more high-latitude area cultivated in the northern hemisphere.

In general, in temperate areas it is projected that warming will be greatest during the winter months but warmer winters are also expected to create problems for agriculture, especially with respect to pests as extreme winter cold is often critical for controlling populations.

Box 6: Temperature relations for wheat grain

A number of laboratory studies conducted with a temperature range of 25 – 35°C have shown:

- Average mean grain weight declined 16% for each 5°C increase in temperature
- Average grain yield decreased by 17% for each 5°C rise in temperature
- Average grain yield decreased by 8 – 10% (through having 5 – 6% fewer grains combined with 3 - 4% smaller grain weight) for every 1°C rise in temperature

When the mechanism of reduced grain filling because of higher temperatures was investigated, it was found:

- Photosynthesis had a broad temperature optimum from 20 to 30°C but declined rapidly above 30°C
- Rate of assimilate movement out of the flag leaf (phloem loading) was optimum around 30°C
- Rate of assimilate movement through the stem was temperature independent from 1 to 50°C
- Respiration effects did not result in decreased grain size with increasing temperature

Reduction of grain weight by heat stress may be explained mostly by effects of temperature on rate and duration of grain growth:

- When temperature was increased from 15/10°C (day/night) to 21/16°C, grain filling time reduced from 60 to 36 days and grain growth rate increased from 0.73 to 1.49 mg/grain/day (with a result of minimal influence on grain weight at maturity)
- When temperature was increased from 21/16°C (day/night) to 30/25°C, grain filling time reduced from 36 to 22 days with a minimal increase in grain growth rate from 1.49 to 1.51 mg/grain/day (resulting in mature grain weight being significantly reduced at the highest temperature)

So, in wheat, high temperatures reduce grain yields because of a reduced number of grains formed and a shorter grain growth time.



Tropical zones

In the tropics, the situation is very different. Any increase in temperature is projected to cause crop yield decline, particularly for maize and rice (whose centres of origin are in or close to the tropics) as these crops are already near the upper limits for their optimum growth (Fig. 6). Indeed, areas where coffee is currently produced will generally get hotter leading to a contraction in the top quality areas and the lower limit of cultivation rising by 3 to 7m every year. Overall there is likely to be less coffee produced of poorer quality (Section 7).

In tropical regions, the impact of increased temperatures is generally projected to be less than that expected in temperate regions (IPCC, 2007) but are still sufficient to significantly reduce the length of the effective growing season, particularly where more than one crop per year is grown. In semi-arid regions and other ecozones where there is wide diurnal temperature variation, relatively small changes in mean annual temperatures could markedly increase the frequency of highest temperature injury. For example, canopy temperature is 10 to 15°C higher in dryland cotton (*Gossypium hirsutum* L.) than in irrigated cotton (Section 6). Thus, global warming would reduce dry matter accumulation in dryland cotton because of increased respiration, and reduced photosynthesis and cellular energy.

Many crops in the tropics already experience heat stress. As temperatures increase due to climate change, crop failure in some traditional areas would become more commonplace. For example, crop growth models show that rice yields decrease 9% for each 1°C increase in seasonal average temperature. This chronic effect of high temperature differs significantly from the acute effect of short-term temperature events, because seasonal temperature effects are mostly a result of effects on crop development. For most grain crops, there is much greater genotypic variation in thermal requirements for vegetative than for reproductive development. As long-term temperatures increase, grain-filling periods decrease, and there appears to be little scope to manipulate this effect through existing genetic variation within species.




Figure 6 Comparison of temperature rise on crop yields in temperate and tropical regions (adapted from IPCC, 2007 in Nicholls *et al.* 2008).



3.2 Changing precipitation patterns

Like temperature, precipitation and water availability is central to agriculture, with productivity being optimal at a particular water balance. If this is decreased (e.g. going towards drought) or increased (towards saturation), then productivity will decrease. Although an increase in global average precipitation is projected (Box 3), this will be highly variable with certain regions experiencing prolonged droughts and/ or extensive heavy rainfall. In the UK for example, prolonged heavy rainfall over winter will be followed by periods of dry weather (drought) interspersed with intensive thunderstorms. Such changed precipitation patterns are expected to greatly impact agriculture, in terms of yield, distribution of crops across the landscape and opportunities for new crops and the expansion of marginal agricultural practices.

Agricultural productivity in tropical Asia is sensitive to changes in the nature and characteristics of the monsoon. Simulations of the impacts of climate change using crop models show that yield decreases due to climate change could have serious impacts on food security in this region. Climate change is likely to cause environmental and social stress in many of Asia's rangelands and drylands. In the arid and semi-arid tropics of Africa, which are already having difficulty coping with environmental stress, climate change resulting in increased frequencies of drought poses the greatest risk to agriculture. Impacts are related to projected temperature increases, the possible consequences to water balance of the combination of enhanced temperatures and changes in precipitation and sensitivity of different crops/cropping systems to projected changes. In Latin America, agriculture and water resources are most affected through the impact of extreme temperatures (excessive heat, frost) and the changes in rainfall (droughts, flooding).

Generally, increased precipitation would probably result in a greater risk of erosion, but could provide the soil with better hydration, according to the intensity of the rain. Soil erosion increases in regions with increased frequency and severity of rainfall, particularly in winter. Nutrient leaching may increase, likewise increasing salt in soil due to drought. The interactions between climate change, water scarcity and declines in agricultural productivity could lead to regional tensions and even open conflict between states already struggling with inadequate water supplies due to rising populations and over-pumping of groundwater.

3.3 Effects of increasing atmospheric CO₂

Plant response to elevated CO₂ alone, without climate change, is positive and was reviewed extensively by the IPCC (Easterling *et al.* 2007) among others. The effects of elevated CO₂ on plant growth and yield will depend on photosynthetic pathway, species, growth stage and management regime, such as water and nitrogen applications. On average, across several species and under unstressed conditions, compared to current atmospheric CO₂ concentrations, crop yields increase at 550ppm CO₂ in the range of 10–20% for C3 crops (e.g. rice, wheat) and 0–10% for C4 crops (e.g. maize, sorghum). Observed increase of above-ground production in C3 pastures is about 10%.

Simulations of unstressed plant growth and yield response to elevated CO₂ in the main crop-simulation models (e.g. CROPGRO, SIRIUS) are in line with recent experimental data, projecting crop yield increases of about 5–20% at 550ppm CO₂. The main crop and pasture models, CENTURY and EPIC, project above-ground biomass production in C3 species of about 15–20% at 550ppm CO₂, i.e. at the high end of observed values for crops, and higher than recent observations for pasture.

Importantly, plant physiologists and modellers recognize that the effects of elevated CO₂ measured in experimental settings and implemented in models may overestimate actual field- and farm-level responses, due to many limiting factors such as pests, weeds, competition for resources, soil, water and air quality, which are neither well understood at large scales, nor well implemented in models. Assessment studies should therefore include these factors where possible, while analytical capabilities need to be enhanced. It is recommended that yield projections use a range of CO₂ effects to better convey the associated uncertainty range.



Carbon dioxide is essential to plant growth. Rising CO₂ concentration in the atmosphere can have both positive and negative consequences. Elevated CO₂ increases the size and dry weight of most C3 plants and plant components. Relatively more photoassimilate is partitioned into structural components (stems and petioles) during vegetative development in order to support the light-harvesting apparatus (leaves). The harvest index tends to decrease with increasing CO₂ concentration and temperature. Selection of plants that could partition more photoassimilates to reproductive growth should be a goal for future research. As more is learned about the effects of anticipated climate changes on crops, more effort should be directed to exploring biological adaptations and management systems for reducing these impacts on agriculture and humanity.

3.4 Elevated CO₂, temperature and precipitation interactions

Temperature and precipitation changes in future decades will at least modify, if not limit, direct CO₂ effects on plants. High temperature during flowering may lower CO₂ effects by reducing grain number, size and quality. Increased temperatures may also reduce CO₂ effects indirectly, by increasing water demand. Rainfed wheat grown at 450ppm CO₂ demonstrated yield increases with temperature increases of up to 0.8°C, but yield declines with temperature increases beyond 1.5°C; additional irrigation was needed to counterbalance these negative effects (Easterling *et al.* 2007). In pastures, elevated CO₂ together with increases in temperature, precipitation and N-deposition resulted in increased primary production, with changes in species distribution and litter composition. Future CO₂ levels may favour C3 plants over C4, yet the opposite is expected under associated temperature increases; the net effects remain uncertain.

Climate impacts on crops will be dependent on precipitation. Over 80% of total agricultural land, and almost all grazing land, is rainfed; changes in precipitation as projected by GCMs will affect both the direction and magnitude of the overall impacts. Changes in precipitation and thus evapotranspiration modify ecosystem structure and function, as shown for UK agricultural grasslands by field-based research led by CABI and partners (Grime *et al.* 2000; Masters *et al.* 1998; Masters & Brown, 2001). Higher water-use efficiency and greater root densities under elevated CO₂ in field systems may alleviate drought but their large-scale implications are not well understood.

3.5 More frequent and extreme weather events

Extreme climatic events such as drought, flooding, hail, hurricanes and tornados are all expected to increase in frequency and severity as a consequence of climate change, and all can reduce agricultural production. Projected increases in frequency and severity of extreme events (in addition to the projected impacts of average climate change) will have significant consequences for food production and food insecurity. At the worst, this can be the complete destruction of agricultural systems but is more likely to be a greater reduction in crop yields and livestock productivity beyond the impacts due to changes in mean climate variables alone. For example, there is an increased risk of crop losses in Bangladesh, from increased flood frequency. In the USA, projected heavy precipitation will lead to excessive soil moisture causing production losses (a doubling has been estimated by 2030 to US\$3 billion p.a.; Rosenzweig *et al.* 2002). Increased soil erosion and salinization have also been projected.

More frequent extreme events will reduce yields over the longer-term by directly damaging crops at specific developmental stages, such as temperature thresholds during flowering, or by making the timing of field applications more difficult, thus reducing the efficiency of farm inputs. Climate variability and change also modify the risks of fires, and pest and pathogen outbreaks, with negative consequences for food production.



3.6 Changes in sea level

Obviously, changes in sea level will affect aquaculture (e.g. shrimps in the Indus Delta in Pakistan); rising seas will contaminate coastal freshwater aquifers with salt water, as experienced already by a number of island states. Such contamination of water quality will have knock-on effects on agricultural productivity. Higher seas also make communities more vulnerable to storm surges which can be 5–6 metres high.

A rise in sea level will reduce the available land area for agriculture and this is particularly so for some CABI member countries in Asia such as low lying areas of Bangladesh, India and Vietnam which will experience major loss of rice crop as sea levels rise. Vietnam relies heavily on the Mekong Delta for rice planting and any rise in sea level of up to one metre will submerge several sq. km. of rice paddies, rendering Vietnam incapable of producing its main staple and export of rice.

In the following Sections, we illustrate some of these generic points regarding the impact of climate change on agricultural production through focusing on two CABI member countries and two commodities that CABI works on together with its member country partners.



4. CABI Member Country Focus: India, agriculture and climate change

The impact of climate change is expected to be very high in India due to greater dependence on agriculture, limited natural resources, rapid increases in human and livestock population, changing pattern in land use and socio-economic factors that pose a great threat in meeting food, fibre, fuel and fodder requirements. There is a likelihood of a considerable impact on agricultural land-use due to snow melt, availability of irrigation, frequency and intensity of inter- and intra-seasonal droughts and floods, soil organic transformation matters, soil erosion and availability of energy as a consequence of global warming, impacting on agricultural production and hence the nation's food security. Climate change is expected to impact on the hydrological cycle, namely precipitation, evapotranspiration, soil moisture etc., which would pose new challenges for agriculture.



4.1 Climate and future climate of India

Observations by the Indian Meteorology Department and the Indian Institute of Tropical Meteorology have confirmed the trends reported by the IPCC for India's climate (monsoon rain, temperature, heat waves, glacial melt, droughts and floods).

Monsoon: At the country level, there is no consistent trend in monsoon rainfall during the last century, although there are some regional patterns. Areas with increasing monsoon rainfall are the west coast, north Andhra Pradesh and north-west India. Areas with decreasing monsoon rainfall are over east Madhya Pradesh and adjoining areas, north-east India and parts of Gujarat and Kerala (-6 to -8% of normal over 100 years).

Temperature: A significant average warming of 0.4°C has been recorded over the last century (1901–2000). A significant warming has been observed along the west coast, in central India and the interior peninsula and over north-east India. A cooling has been observed in the north-west and parts of southern India.

Drought, flood, storm, glacier: No significant long-term trend in the frequency of droughts or floods has been recorded over the past 130 years. Additionally, the frequency of cyclonic storms forming over the Bay of Bengal over the period 1887–1997 has remained constant. However, there is evidence that Himalayan glaciers are melting and receding rapidly.

Projected climate change for all of India indicates that by 2100 there will be an increase of 15–30% in rainfall across the country and mean annual temperature will increase by 3–6°C, with the maximum increase over northern India. The warming is also projected to be relatively greater in winter and post-monsoon seasons (Christensen *et al.* 2007).

By 2050, temperature is projected to increase by 2–4°C, there are expected to be marginal changes in monsoon rain during the monsoon months but large changes during the non-monsoon months. The number of rain days is projected to decrease by over 15 days and rain intensity to increase by 1–4mm/ day, also with an increase in the frequency and intensity of cyclonic storms.

4.2 Impact of climate change on Indian agriculture

Agriculture represents a core part of the Indian economy, representing 35% of the GNP and thus is central in the country's development (in the last half century, grain production has quadrupled).



Increased temperatures as projected by the IPCC by 2100 are expected to lead to a 10-40% crop productivity loss overall. This is supported by work from the Indian Agricultural Research Institute which projects that for every 1°C rise in temperature there is an associated loss of up to 5M tons in wheat production, assuming irrigation continues at today's level. The loss in total farm net revenue expected is between 9 and 25% for a temperature rise of 2-3.5%. It is expected that a major impact of climate change will be on any rainfed crop (other than rice/ wheat) which contributes approximately 60% of the cropland area. Rainfed agriculture is practiced by the poorest farmers.

Using historical records and crop-growth modelling, increasing temperature was projected to have varying negative impacts on the yields of barley, chickpea, mustard and wheat in north-west India.

Wheat: Climate change will lead to changes where in the country, wheat can be grown profitably. A 2–3°C increase in temperature will reduce yields in the majority of the wheat growing areas (Aggarwal & Sinha, 1993), and this yield reduction will be greater in non-irrigated (and thus water stressed) crops due to rainfall variability. So, the warmer regions will suffer crop losses. A minor increase of only 0.5°C during the winter is expected to decrease wheat yield by 0.45 t/ ha. However, as with most crops, the response of wheat yields is dependent on CO₂ fertilization, decreasing by up to 64% with no fertilization but ranging between +4 and -34% with CO₂ fertilization (Rao & Sinha, 1994). Climate projections indicate continual decrease in wheat yield (Fig. 7).




Figure 7 Projected impact of climate change on wheat: production (total yield in million tons) estimates for India.

(Source: Aggarwal *et al.* 2002 quoted in Chattopadhyay, 2008)



The growing season for wheat is limited by high temperatures at sowing and maturation. As wheat is grown over a wide range of latitudes in India, it is frequently exposed to temperatures above the threshold for heat stress. For example, high maximum and minimum temperatures in September (about 34/20°C) adversely affect seedling establishment, accelerate early vegetative development, and reduce canopy cover, tillering, spike size and yield. High temperatures at the end of February (25/10°C), during March (30/13°C) and April (30/20°C) reduce the number of viable florets and the grain-filling duration. The situation is similar for sorghum and pearl millet which are exposed to extreme high temperatures in Rajasthan (Abrol & Ingram, 1996; Abrol *et al.* 1991).

Rice: An increase in temperature of 2–4°C is expected to reduce yields. An increase of 2°C may decrease the rice yield by 0.75 t/ ha in high yield areas (Sinha & Swaminathan, 1991). Further work projects that a 1°C rise decreases rice yields by about 6%. There are though, regional impacts, with eastern regions having the greatest impact of increased temperature and decreased sunshine leading to fewer grains and shorter grain filling time. However, northern regions will not suffer a net reduction in yield because decreases to increased temperature are offset by increases to higher incidence of sunshine. The beneficial effect of CO₂ will be cancelled out by the negative effects of greater temperatures.

4.3 Implications of climate change impacts on Indian agriculture

Without doubt, increasing climatic variability (droughts, floods, tropical cyclones, heavy precipitation, extreme heat and heat waves) in the future will lead to large fluctuations (annual/ seasonal) in agricultural production, resulting in increased risk of food insecurity issues and thus will affect rural livelihoods throughout India.

Food security: changing crop yields (increasing in some areas and decreasing in others) combined with a redistribution of which areas/ regions can support growth, development and yield of which crops, will directly impact consistency, quality and quantity of food supply, and thus become a bigger food security issue.

Trade: it is highly likely that imports and exports for commodity crops will be affected and is particularly important for cash crops (e.g. chillies, coffee, tea, cotton, mango, grapes).

Livelihoods: any changes in crop yield and distribution will affect livelihoods, particularly if yields decrease and main crops in a region become marginal (or impossible) to grow due to climate change. Capacity building and education/training will be needed to cope with the new agricultural landscape.

Adaptation: Measures will need to be introduced to meet the changes imposed by climate change on agriculture on India, for example, alternative crops, new cropping patterns, water conservation, new irrigation technology and capacity building will all have to be researched and implemented as part of an adaptation strategy. Farmers and other stakeholders can begin to adapt, to a limited extent, to reduce agricultural losses. Simple adaptations such as changing sowing/ planting dates and using new crop varieties could help in reducing impacts of climate change to some extent in the immediate future. The Indian Agricultural Research Institute estimate that losses in wheat production in the near future could be reduced from 4 – 5 M tonnes to 1 – 2 M tonnes if the majority of farmers could change to more timely targeted planting and change to use better adapted varieties. This change of planting would, however, need to be examined from a cropping systems perspective.

4.4 What is CABI doing?

With partners, CABI is implementing a project investigating the resilience of coffee production to Coffee Leaf Rust and other diseases. Climate change is an important component as it will lead to warmer and more variable conditions in many areas, meaning that coffee varieties will be under increased stress in future years. This will raise many urgent issues, including the conservation of coffee genetic resources and continued productivity of the small holder coffee sector. Climate change affect many things, including the coffee genetic resources of a country – it will be necessary to review policies and practices to ensure that these resources are not eroded and that any new genetic material obtained is well conserved. The outputs of the project will lead directly to the uptake by



farmers of appropriate varieties, growing techniques and methods for farm adaptation and control of diseases. Valuable data will be shared on the combinations of tree stock and growing style that work best under different conditions, which can then be more widely replicated. The intention is that the concept of long term participatory trials promoted by the project will continue long after the project is finished.

Ghana is the CABI member country chosen as the focus for the West African sub region.



5. CABI Member Country Focus: Ghana, agriculture and climate change

The Human Development Index of 177 countries ranks Ghana at 138, where overall, approximately 40% of the population is considered to be poverty stricken. Poverty levels are highest in the north where drought/desertification are dominant and widespread. In the north about 88% of the population lives in poverty (WHO, 2006). Ghana's GDP consists of agriculture (37%), services (38%) and industry (25%). Major agricultural products include cocoa, rice, coffee and timber, of which cocoa and timber are exported (Dazé, 2007).

Land degradation is an increasing national problem. Desertification affects over 35% of Ghana's total land area, with the north-eastern region being at greatest risk. Unsustainable land uses, combined with natural vulnerability from the climate are the prime drivers of desertification. Greater incidence of droughts in the future will add to this vulnerability. Soil degradation is caused by erosion, nutrient leaching and increasing salinity. The northern savannah is highly vulnerable to soil erosion, and the situation is projected to get worse due to heavy rainfall/storms as a consequence of climate change. Unsustainable cultivation, deforestation, overgrazing, exposure, fire and population pressures all contribute to the depletion of nutrients. Combined with the impact of climate change, reduced soil productivity will make subsistence/ sustainable agriculture-based livelihoods almost impossible.



5.1 Climate and future climate of Ghana

Over the last 30 years, climatic data shows that there has been a 1°C increase in average temperature, a 20% decrease in precipitation and a 30% decrease in runoff. Future projections of climate change indicate that Ghana will experience a 2.5 – 3.2°C increase in average temperature and annual rainfall will decrease by 9-27% (the relatively large range is principally due to spatial variation) by 2100. This historical data from the last 30 years, also shows an annual sea level rise of 2.1mm. By 2100, sea level could have risen by 1m resulting in coastal inundation, coastal erosion, saltwater intrusion into freshwater sources and a possible increased earthquake risk due to liquefaction (Amponsah, 2004; Boko *et al.* 2007; Dazé, 2007).

5.2 Impact of climate change on Ghanaian agriculture

The agriculture sector employs over 60% of the workforce and is largely based on smallholdings (over 85% are 2 ha or less in size) and the majority of agriculture is rainfed. Cocoa is the principal cash crop, being grown on 40% of cultivated land. Cassava, yam and cocoyam are the most important root and tuber crops, which in total account for 40% agricultural GDP (Dazé, 2007; CIA, 2007; GINC, 2000). Millet is the staple crop in the savannah and maize and rice are widely consumed throughout the country. Cereals are relatively vulnerable to climate change and are important in terms of food security. Ghana's high temperatures result in low cereal yields due to a reduction in growing period and increased evapotranspiration (higher water use).

In the north of Ghana there is a high level of dependence on agriculture for livelihoods and the area is climatically sensitive with low and decreasing rainfall and frequent, recurring droughts, together making this the most vulnerable region to climate change. The dependence on rainfed agriculture across the country makes farmers in Ghana particularly vulnerable to climate change. Irrigation has recently been introduced but is not widespread. In 2000, only 0.5% of the cultivated area in Ghana



was irrigated (AQUASTAT, cited in Dazé, 2007) but climate change will increase demand for irrigation water. Coupled with the reductions in rainfall and runoff as projected by climate models (Boko *et al.* 2007), impacts on water availability for agriculture will be significant for this country.

Food crops: Cereals will be negatively affected by climate change. Increased variable rainfall will result in droughts, soil degradation and unpredictable growing seasons, all of which will reduce yields. Additionally, roots and tubers will be negatively impacted; cassava production is projected to decrease by 53% and cocoyam by 68% by 2080 (Anon, 2007; Dazé, 2007).

Cocoa: Cocoa accounts for 60–70% of foreign export earnings from agriculture or 20–25% total foreign export earnings. Over 800,000 smallholder families (350,000 farm owners), mainly in the western region, depend on cocoa production for their livelihoods, as cocoa represents 70–100% of their household annual incomes. Cocoa farms are small, generally 0.4–4ha each with a total cultivation area of 1.45 M ha, and do not use technology or any significant inputs. Cocoa can only be profitably grown within a temperature range of 18–32°C and is susceptible to drought and the pattern of cropping is related to rainfall distribution (CRIG; Dazé, 2007). Over four months, dry weather results in soil water deficit leading to seedling mortality, reduced bean size and increased pest attack. The plant is also highly sensitive to light, as prolonged exposure at high intensity disrupts photosynthesis but low light suppresses flowering. Given these climatic constraints, cocoa will be highly susceptible to climate change and this will have a large impact on the Ghanaian economy given its pre-eminence as a foreign exchange earner.

Production of cocoa is projected to become increasingly difficult as climate change will increase temperatures (thus increasing the probability of disrupting photosynthesis and decreasing growth) and lead to more erratic precipitation, which again will limit growth, development and yield. As a naturally occurring understory tree, cocoa is sensitive to high light intensities which can induce die-back of fine branches leading to attack by various pathogenic fungi and as more of the canopy becomes exposed to sunlight, the risk of further attack by mirids (already a very significant problem in Ghana) will increase. The geographic distribution of cocoa (and existing and new pests) will change, crop yields will decrease and there will be a greater incidence of crop loss: all of which will affect farm income, livelihoods and farm level decision making. A significant constraint to cocoa production currently in Ghana is that of aggressive black pod (*Phytophthora megakarya*) which can readily destroy a crop. As this pathogen thrives in humid conditions, any increases in precipitation or changes in patterns of rainfall could have significant negative effects on the spread and impact of this disease. Management of this disease is currently via fungicide sprays (recommended 8 times per year) so increased precipitation could mean more frequent spraying with associated environmental impacts.

5.3 Implications of climate change impacts on Ghanaian agriculture

Farmers need to develop and implement adaptation and mitigation strategies urgently. Simple strategies include the rehabilitation/ restoration to sustainable production on degraded farms to reduce the effect of land degradation that encourages the adverse effects of climate change. As with many countries, capacity building on the effects of climate change and how to deal with these changes needs to be urgently established.

Food security: Ghana will be subject to the same issues as many other developing countries. Decreasing crop yields (in general) combined with a redistribution of which areas/ regions can support growth, development and yield of which crops, will directly impact consistency, quality and quantity of food supply, and thus become a bigger food security issue.

Trade: it is highly likely that imports and exports for commodity crops will be affected and is particularly important for cocoa, the principle cash crop.

Livelihoods: any change in crop yield and distribution will affect livelihoods, particularly if yields decrease and main crops in a region become marginal (or impossible) to grow due to climate change. Ghana has an international reputation for producing cocoa of a high quality and as such, any climatic



changes affecting quality would have large implications on trade and foreign exchange earnings. Heavy daily rainfall prevents cocoa growers drying beans sufficiently to allow sale at a fair price. So, farmers withhold their crops from the market until the usual warm sunny weather materializes and beans can be dried, quality improves and farmers receive higher prices (value added) instead of selling wet beans. However, if warm sunny weather does not return and the cocoa cannot be sun dried, then farmer livelihoods may be affected as they will have to sell wet beans or invest in artificial driers which are currently beyond the means of most cocoa farmers. As Ghana is projected to become drier in the future under climate change then this may not be an issue. A current and future threat to livelihoods is the incidence and impact of pest and diseases reducing yields.

Currently, there are two production cycles of cocoa in Ghana and climate change could affect this; producers need to be made aware of likely impacts on their cash flow and prepare accordingly.

Adaptation: for cereals, adaptation strategies that could be implemented include the development or acquisition and use of drought tolerant varieties, the use of early maturing genotypes, use of conservation agriculture as much as possible including low tillage, altered planting dates to suit changing cropping cycles and an increased use of irrigation. Cropping can be maximized, such that in the savannah, early millet, groundnuts and cowpeas are planted; water conservation methods are introduced for rice production. In the savannah-forest transition, maize could be replaced with sorghum or cowpeas and vegetable production diversified. While in the forests, maize and cowpeas could be rotated. A similar approach to that for cereals can be adopted for root and tuber production, with improved or new capacity for improved farm technology, new drought tolerant varieties, shifted sowing time, the implementation of irrigation, improved post harvest technology and diversification of livelihoods.

A prime adaptation strategy for cocoa farmers is the implementation of drought management practices, including the exploration of irrigation, the continuing use of shade trees and improved drought tolerant varieties. Socio-economic strategies such as secure land tenure, effective pricing and access to credit schemes would increase the capacity for farmers and other stakeholders in the supply chain to understand and adapt to climate change. Fortunately, as most of Ghana's cocoa is produced under shade, some of the worst effects of changes in light intensity could be reduced but any cocoa cultivation occurring with little or no shade needs to have the impact of higher light intensities and temperatures on the crop taken into consideration.

5.4 What is CABI doing?

CABI has worked with the Cocoa Research Institute in Ghana (CRIG) for a number of years on various projects including examining alternatives to spraying pesticides to manage diseases such as aggressive black pod and most recently to help CRIG to survey pesticide use along the supply chain. Also with CRIG and Cadburys, CABI has pioneered the Ghana Cocoa Farmers Newspaper which could be a vehicle for introducing the concept of climatic change to cocoa farmers using cartoons and simplistic diagrams (<http://www.cabi.org/files/newspaperposter.pdf>).

CABI's work on irrigation and cotton (Section 6) can be applied to many areas where cotton is grown and water is limiting. This is the case in Ghana, and the irrigation of cotton in the north of the country could be a focus area for future work.

We now focus on two commodities, to give examples of CABI's work with cotton and coffee.



6. Commodity Focus: Cotton

Cotton (*Gossypium* species) is a leafy, green shrub, a member of the Hibiscus family, producing natural fibre. Within the native ecosystem, cotton is a perennial growing to over 3m in stature but commercially, cotton is now grown as an annual. Cotton is the most widely produced natural fibre in the world and comprises over 40% of the global textile market. Cotton is highly dependent on water for growth; it takes 20,000 l of water to make 1kg of cotton (which is enough for a single t-shirt and a single pair of jeans) and it is this water demand that makes cotton potentially susceptible to changes in climate. Places such as China, Pakistan and Central Asia, including India, are becoming increasingly water limited, but are also important cotton producing areas, so there is a potential conflict as water becomes increasingly limited.

6.1 Impacts of climate change on cotton

Although it was initially thought that cotton yields would be increased, due to the CO₂ fertilization effect, it now appears that water relations may be more important (Reddy *et al.* 2002). Such projected increases in cotton yield were negated when changes in temperature and precipitation were included in the analysis. Future climate change scenarios for the Mississippi Delta, USA, for example, estimate a 9% mean loss in fibre yield.

As discussed in general in Section 3, elevated CO₂ concentration in the atmosphere is projected to increase photosynthesis and water use efficiency in agricultural plants and thus increase growth and yield (a 40% increase in yield for cotton has been suggested). Yet, the situation is more complex, as the beneficial effects of increased CO₂ are expected to be offset by reduced rainfall, higher temperatures (heat stress) and their combined effect on evapotranspiration. Water limitation and temperature extremes will have detrimental effects on growth and development and this is expected to be greater than the growth promoting effects of increased CO₂.

Moderate increased temperatures could extend the cotton growing season and so have a positive effect on yield through an extended period for growth. Further positive impacts of increased temperatures include improved crop establishment, reduced cold shock, bigger plants with bigger leaves and longer fibres. However, this may be offset through an increased frequency of days/ nights with high temperatures (heat stress) which has been found to have detrimental effects on growth and development, including reduced photosynthesis, high respiration, reduced flowering, reduced cotton boll development and short fibres.

The overall effect of climate change on cotton growth and development is, on balance, going to be detrimental. The negative effect of water limitation will be greater than the beneficial effects of moderate temperature and elevated CO₂ and so cotton yield is expected to decrease under climate change.

6.2 Pakistan, climate change and cotton

Pakistan already has most of its arable land under cultivation, and agriculture accounts for a quarter of the GDP and employs 50% of the workforce. The economy is highly dependent on trade in textiles (and the vagaries of that market), particularly cotton. Pakistan is an agro-based economy and the sustainability of agriculture depends heavily on the availability of water.

The implications of climate change for Pakistan are likely to be very high because of direct and indirect effects on water and temperature relations. Climate change, through increasing temperatures, will negatively affect the volume of fresh water storage in Himalayan glaciers and its slow release for crops in addition to changing the timing, duration and intensity of monsoon rainfall. As the glaciers retreat at a rapid rate, the amount of fresh water released into rivers and streams, and hence available for agricultural and domestic use, is reduced as less water is stored. Additionally, if water is released too quickly then it is not available for agricultural use, rivers and the land quickly become saturated and flooding may occur. These climate driven changes will directly add to water,



food and energy insecurity. Rainfed agricultural areas will be particularly hard hit. Although irrigation in agriculture in general is scarce in Pakistan, scheduled irrigation is practiced in cotton production to the extent that overuse of irrigation water is very common in cotton.

For the last half century, the climate of Pakistan has been consistently changing, and this is expected to accelerate over the next century due to an increasing influence of climate change. Precipitation patterns are changing and the frequency and intensity of extreme precipitation events has increased along the foothills of the Himalayas. Temperature extremes, heat waves and prolonged cold spells, are becoming common.

The most significant climate change impact projected for Pakistan is increased variability in the monsoon rains. Summer monsoon rains could increase by up to 60%, resulting in flooding and reducing the ability to produce fibre and food, destroying irrigation systems and flattening crops such as cotton.

Pakistan used to have a water surplus overall, which enabled investment in the production of commodities like cotton. In many respects the Pakistan textile industry is now dependent on cotton to generate significant export revenues. However, the country is now in water deficit, and the situation is going to get worse as climate change continues. Extended droughts have reduced fresh water supplies, so highlighting the importance of adopting water conservation measures for the judicious use of this resource, especially in cotton production.

6.3 What is CABI doing?

CABI has investigated the impacts of cotton production on the freshwater ecosystem and identified good management practices to ensure sustainable cotton production with a minimum threat to the freshwater ecosystem. CABI empowered farming communities to identify and adopt sustainable cotton production technologies with long term benefits for water resources. This included developing water requirements and water use patterns of the cotton plant. CABI integrated water saving techniques with other crop management techniques and developed plant water requirement monitoring criteria to assess when a cotton field should be irrigated. As climate change accelerates and water becomes scarcer in the cotton growing areas of Pakistan, then targeted irrigation with appropriate knowledge transfer will become increasingly important. Such knowledge transfer is a key strength of CABI, where there is considerable expertise in training different stakeholders along the cotton supply chain, so integrating climate change into this capacity building will increase the ability of cotton producers and distributors to adapt to climate change (CABI Annual Review, 2008/2009; www.cabi.org).



7. CABI Commodity Focus: Coffee

Coffee is a globally significant crop, grown in over 50 countries. Coffee exports are important foreign exchange earnings for producer countries and most coffee farmers are smallholders, who depend on coffee for a substantial proportion of their livelihood. Coffee is suited to production on a small scale, as a family unit, as well as being appropriate for hillside farms where more demanding cash crops cannot be easily grown. The diversity of ways and environments in which this crop is grown means that the effects of climate change will be very diverse. Nevertheless, evidence of observed impacts of climate change on coffee production and the regions where coffee is grown, is increasing. After considering this aspect, the future impacts of climate change are examined. CABI's work on coffee is geographically widespread, covering Latin America, parts of India, Africa and PNG. Hence there is no specific country focus regarding coffee, but the work, in relation to climate change is highlighted.

7.1 Observed impacts of climate change on coffee

The climate has already changed in the coffee lands of Colombia; night-time minima have risen by 1°C over the last 50 years due principally to increased cloudiness. This increases pest pressure; for example the coffee berry borer, which is resurging after control successes in the 1990s. Coffee is no longer grown below 1200 m in Colombia, principally due to the favourable conditions for this pest. Similarly, in 2007, Costa Rica lost 1.6% of its harvest to western leaf spot (*Mycena citricolor*) which surges in the increasingly intense rainy season that precedes the harvest period; similar losses were recorded in 2005 after Hurricane Stan – the losses sound small, but margins for most coffee producers are very slender and the pathogen attacks above 1400 m which is where the best coffee grows. Production has also been affected in Cuba where in November 2007, the newspaper *Juventud Rebelde* reported a loss of 12% of the coffee harvest in the provinces of Granma and Santiago de Cuba due to extreme weather conditions; while in Nicaragua in 2006, the poor rains, caused by an El Niño event, caused production levels to fall by up to 50%. These events are becoming increasingly frequent but the surprise was that this time it affected not only the traditionally drier area of Segovias, but also the normally wetter areas of Jinotega and Matagalpa. Cooperative technical managers of PRODECOOP and CECOCAFEN definitively attribute this decline in production to climate change and are anxious to develop strategies to confront it.

In Africa too, many are starting to link coffee production problems to climate change (Box 7). In the mountains of Yemen, where some of the world's highest quality coffee grows, water extraction for irrigation is five times the quantity of precipitation. It is estimated that at current levels the region will dry up in a period of 20 years.

In a world of increased threat from flood and drought, it would make sense to distribute production over many countries to reduce the risk of shortages, but the opposite is happening with only seven countries responsible for three quarters of world production; Vietnam now exports more than the whole of Africa (Fig. 8). Hence we are seeing increased land brought into production to grow coffee in a few countries and abandonment of coffee in others, even though coffee in the latter may be more environmentally and socially sustainable. Short-term market considerations are sometimes moving the coffee further away from environmental sustainability, despite the actions of some commercial sustainable coffee schemes.

No single example above can be proved to be directly caused by climate change, but the pattern of events is exceptional and wholly consistent with projections from models of climate change.

7.2 Impacts of climate change on coffee

For coffee production in Mexico, regression models have shown that rainfall in Veracruz between 1969–98 decreased by 40 mm p.a. and temperatures increased by 0.02°C p.a. Extrapolating these changes to 2020, coffee production would decline by 34%, reducing profits of US\$200/acre, to less than \$20/acre. Yield declines of Arabica are virtually certain in a warming world because it is a montane species, not adapted to high temperatures. Robusta as a high temperature alternative is



not an option on the mountainsides because it is uncompetitive with mechanized production in Brazil. However, in Brazil, dramatic changes in production and revenue have been predicted from a physiological model by Pinto *et al.* (2005), who predicts yield declines as mean annual temperatures exceed 23°C, due to increased flower abortion (Box 8).

Quality would certainly be affected as mean temperatures rise, as the maturation process of the coffee cherry speeds up faster than the development of the bean, leading to lighter and lower quality berries. Also within coffee countries, zones where coffee is grown will change. A 3°C rise this century translates to the lower limit of coffee rising by 10 to 20 ft per year. Production will shift in many cases, either putting pressure on other agricultural land or on biodiversity of virgin ecosystems.

For countries with fewer comparative advantages, sustainable diversification options are few, especially in those countries which have a significant industrial economy that leads to an uncompetitive labour price. Thus in Colombia the coffee lands are being converted to lower cost cattle pasture and even annual crops. Although this may make economic sense, externalities such as soil compaction, erosion and run-off mean that the ecological services provided by a perennial such as coffee are diminished.

7.3 What is CABI doing?

CABI's work on coffee with partners in Latin America, Africa, India, Indonesia and in PNG, gives us many insights into how the industry will have to change to face the reality of climate change.

CABI's work on pests and diseases has found increasing signs that pests such as the coffee berry borer and diseases such as coffee leaf rust and American leaf spot are resurging. This is most likely due to more extreme weather events, for instance El Niño and hurricanes that provide conditions especially favourable to them – such occurrences used to be rare events but seem to be now more frequent. CABI's work with farmers will lead to more rapid updates on changing weather patterns and assistance to carry out prompt prevention and control measures. Working with support institutes, CABI is providing the knowledge and capacity to be able to identify and react quickly to new pests and diseases that are likely to appear. Much of the information about climate change and coffee is highlighted in Baker & Haggard (2007).

Another focus area is biodiversity. For example, CABI and partners examined the biodiversity associated with coffee, which even under intensive production such as in Colombia, harbours surprisingly high levels of biodiversity. Under climate change many species currently present in coffee are likely to disappear, but the patchwork fields of coffee, forest patches, streams, gardens and other crops will offer a refuge for species migrating upslope from warmer lower altitudes. Coffee too will decline at lower altitudes but it is clear that the landscape matrix that currently exists is capable of harbouring a wide range of biodiversity that will be greatly needed in the future for efficient ecosystem services, such as pollination and seed dispersal. Through participatory studies that CABI led, we have found that coffee farmers are very aware of environmental problems and are willing to adapt, as long as this does not affect their income. New trees and plants that can generate income and survive in changing conditions will have to be tested.

CABI has also pioneered farmer training schemes in Africa which have focussed on more environmentally friendly post-harvest processing for coffee, so reducing water usage whilst allowing producers to obtain a premium for their coffee.



Box 7: Global Warming Threatens Coffee Collapse in Uganda

"Climate change has affected coffee production already," said Philip Gitao, executive director of the East African Fine Coffees Association. The crop has had less time to mature because rain is falling at the wrong times, affecting coffee quality, Gitao said. And there have been more droughts in the past two to three years than ever before. "If the coffee beans face a lot of sunshine and less rain, the beans will be smaller and in lower yields," Ronald Buule, a central Ugandan coffee farmer, said as he stood at a coffee plot bordered by lush plants, muddy hills, and an orange dirt road. "We are worried about the temperature, but we have limited resources," he added, as he examined his crops under a dense thicket of banana leaves.

Alexis Okeowo in Nsangi, Uganda [from National Geographic News July 24, 2007]

Box 8: Projected decline in production of coffee in Sao Paulo, Brazil

Increase in temperature	Area suitable for coffee km ²	Production tonnes	Change in production tonnes	Change in revenue US\$
+1°C	145,202	269,082	-80,829	-113,160,600
+3°C	75,455	139,614	-210,297	-294,415,800

Source: Pinto *et al.* (2005)




Figure 8 Coffee exports for Africa and Vietnam

The amount of coffee exported from Africa (blue line) is variable between years, reflecting climate, markets and production, but shows a decline overall from the early 1970s. Coffee exports from Vietnam (magenta line) did not start until the early 1980s but has expanded exponentially since then, reflecting the land/ forest clearance for coffee and investment in production. Vietnam is now a globally significant coffee producing country, exporting more than Africa. Source: FAOSTAT



8. Summary and knowledge gaps

It is obvious that, in general, agricultural productivity will decrease, to a greater or lesser extent, under climate change, particularly with rising temperature and fluctuating extreme precipitation, i.e. increasing climatic variability. Consequently, food shortages and food insecurity will increase. LDCs with limited resources and little institutional capacity, will be the least able to adapt to current and future changing climate, variable agricultural conditions, highly fluctuating agricultural commodity markets and increasingly frequent and severe food shortages. Also many LDCs are particularly at risk due to their situation in the tropical zones that are projected to be most affected by climate change.

In areas of increased rainfall, more attention will be needed to reduce flooding, landslides and improve emergency responses. Varieties of all major crops that are more resistant to fungal diseases and pests will be needed as conditions will become more conducive for attack. There will be increased threats from new and different pests or diseases entering and becoming established in new regions or locales. In general, agricultural production technology and knowledge transfer needs revising and redesigning to make it efficient and sustainable to address the present and future threat of climate change.

8.1 Immediate research needs

There is an urgent need for more research and particularly action on adaptation measures in agriculture, especially in LDCs, to assist farmers, extensionists and supply chain stakeholders to reduce the adverse effects of climate change on agriculture. This is particularly so in the tropical zones, where LDCs are projected to be at great risk and often have large, resource-poor populations suffering from food insecurity.

Due to the complex physical and biophysical impacts of the interactions between the different climate variables, combined with varying irrigation, fertilizer and pest and disease management strategies and techniques, regional variability (e.g. topography, soils, land-use and management), and differences in crop growth, development, phenology, yield and resistance to drought and pest and diseases; the detailed impacts of these factors need to be investigated further.

Specific recommendations for further research include:

- Increased precision in climate change projection with higher resolution on spatial and temporal scales;
- Linking of projections with agricultural production systems to suggest suitable options for sustaining agricultural production;
- Preparation of a database on climate change impacts on agriculture;
- Evaluation of the impacts of climate change in selected locations;

In terms of experimentation, there is a lack of knowledge of CO₂ and climate responses for the majority of crops other than cereal staples such as wheat, maize and rice, including many of importance to the rural poor (e.g. root crops, millet, brassicas). Some CABI partners are beginning to conduct important studies looking at climate change drivers and crops of relevance to individual countries. The Indian Agricultural Research Institute, for example, has elevated CO₂ FACE and temperature gradient polytunnel experiments investigating cereal responses to projected climate change. CABI can partner these organisations and projects through our work on pest and disease management. The response of commodity pest and diseases to climate change is a knowledge gap and CABI's expertise and approach can address this research and implementation need.

In addition to the staples such as root crops and cereals, more work needs to be initiated on how climate change will impact on traded commodities such as coffee, cotton and cocoa and how changes in production area, changes in yield etc will impact on the foreign exchange earnings of



LDCs in particular– not least since until these countries own agricultural production is improved, these foreign exchange earners are important in the importation of food to maintain the very survival of the urban poor.

Much uncertainty remains regarding how changes in frequency and severity of extreme climate events, and climate change will affect all sectors. Improved projection of future impacts of climate change requires better representation of climate variability at scales from the short-term (including extreme events) to interannual and decadal. Where datasets exist (public/ private) of climate and preferably yield, of 50 years or greater then there is an opportunity to conduct an analysis and project changes for the near future (next 20 years). These projections should include, if possible, changes in mean, median, standard deviations, variance of temperatures and rainfall, but also the distribution of wet spells, dry spells, absolute maxima and minima, frequency of storms, frosts, and droughts. Valuable information can be gained to aid farmers in what to expect in the immediate future, for example dates of first and last frost, variation in rainfall patterns. This approach will enable future climate information and forecasts and thus better management of crops and cropping systems. The projections can have climate change parameters built into them. CABI with member countries and partners will source datasets and run the analyses and projections. Results and conclusions could then be transferred to the agricultural stakeholders through appropriate means, for example the CABI Environmental Change Web Portal.

For all of these research priorities, arrangements should be made for the sharing of experiences and the transfer of good practices in agriculture that can constitute mitigation and adaptation, again a role for the CABI Environmental Change Web Portal.

8.2 Priorities

One immediate response, as acknowledged by the IPCC (2007; Easterling *et al.* 2007), is to instigate capacity building to enable agricultural stakeholders to deal with climate change, but this development needs to focus on enabling agriculturists to respond to current climatic variability, thus providing the stepping stones to adapt to future, more extreme climatic variability.

CABI's country membership, combined with a depth of practical on-the-ground experience and extensive information and knowledge bases, puts us in a unique position to catalyse action by informing, up-dating and generally building capacity of countries, regions, communities and individuals (e.g. farmers, practitioners, extensionists, farmer groups, land managers) to take concerted action now to adapt to the new realities of the impacts of climate change. Action that CABI can lead and implement to tackle the issues, increasing problems and complexities agriculturists will face now and in the future include:

- **Awareness raising, training on planning and zoning** for donors and decision makers about the need for governments to take firm action to protect the fertile uplands of their countries. These resources will become ever more valuable as climate change forces upslope migration of flora, fauna and human communities. These territories are a country's primary asset and cultural heritage that should not be sacrificed to short term market goals;
- **Capacity building** to reinvigorate institutions, including extension and research divisions. Presently, research is concentrated in a limited number of countries and institutions (e.g. prototype GM coffee varieties with drought resistance, delayed maturation and herbicide tolerance being developed and trialled in Brazil). In general, agriculture has stagnated in development over the last 30 years, although recently, perhaps in response to environmental change and the growing threat of food insecurity and starvation, there are signs of a new green revolution utilizing biotechnology (including varieties, agronomy and processing) and a greater understanding of ecosystem services, resistance and resilience, all central to sustainable development (Box 2). Without a more equal approach to research, planning and active intervention, some countries will become ever less competitive. There is much to learn from Brazil's policies (Box 9).
- **Adaptation planning and implementation:** in the short to medium term there is much that can be done to help farmers adapt to climate change. In areas of increased drought, options and



knowledge exists to start activities in water harvesting, improved irrigation technologies (e.g. drip irrigation, being used for coffee in Nicaragua), drought resistant varieties, use of shade trees where appropriate, lean water use in processing, minimum tillage and many other technologies. There are many possible solutions that are not being developed, trialled nor implemented through lack of resources, knowledge and technology transfer, especially in LDCs.

- **Coordinates efforts of Governments, private organisations and other stakeholders** in mitigating or adapting to the challenges of climate change.

8.3 Adaptation

The United Nations Framework Convention on Climate Change (UNFCCC) instigated National Adaptation Programmes of Action (NAPAs) to provide a process for LDCs to identify priority activities that respond to their urgent and immediate needs to adapt to climate change – those for which further delay would increase vulnerability and/or costs at a later stage. About 40 such Action Programmes now exist and need to be implemented. CABI has relevant expertise to offer in several areas, including those relevant to commodities. Many NAPAs stress the need for introduction of new crops resistant to drought that will require considerable upgrading of extension services and farmer training. Other urgent topics include reforestation, selection and growing of tree species in farmland, soil and water conservation techniques, promotion of agroforestry systems and home-garden agriculture.

Agricultural ministries and supply chain stakeholders need to design and implement adaptation strategies. Adaptation of land use and management, if adopted now, can mitigate (particularly in the short term) climate change impacts. Key strategies, which have largely been discussed previously, include:

- Use of new/ improved varieties, crops, or rotations (longer maturing varieties, heat- and drought-tolerant, requiring less vernalization). Alternatively, switching production from an increasingly unsuitable area to a climatic zone which is projected to be favourable for the commodity in the future;
- Design and implement irrigation and drainage;
- Adaptation to changed growing season or to shift timing of heat stress;
- Design and implement flexible cropping systems with mixed varieties, as trialled and implemented for commodities such as coffee;
- Improved use of sustainable technologies, including integrated pest management, post harvest management and soil health, of which CABI has expertise and programmes.

In developing adaptation strategies, agriculturists and their stakeholders need to ensure they are not adding to climate change. Agriculture, particularly intensive production, produces approximately a quarter of global greenhouse emissions. Given predicted decreases in agricultural production and an increasing demand for food, there is an argument to intensify and expand to meet the demand. This is a real challenge as it needs to be balanced against the need to adapt to climate change and to attempt to stabilize greenhouse gas emissions.

Conserving and promoting ecosystem resistance and resilience has to be a central tenet of adaptation strategies and needs to be balanced with any actions to alter or intensify agricultural production.



Box 9: The lesson from Brazil: government intervention works

Agricultural losses in the 1990s were limiting the development of Brazilian agriculture. These losses were caused by two main factors:

- excessive rain during the harvest period (30% of all cases)
- dry spells during the flowering and grain-filling stage (60% of all cases)

These losses were related to a poor knowledge of rainfall distribution that led the farmers to plant at inappropriate times. To decrease risks, EMBRAPA (the National Institute for Agricultural Research) and the Brazilian Department of Agriculture started an official program of agricultural zoning in 1996 to define planting calendars for rice, beans, maize, soybean, wheat, sorghum, cotton, coffee, and fruits, based on simulation of cumulative water balance.

Agricultural zoning is based on the integration of crop growth models, climate and soil databases, decision analysis techniques, and geo-referencing. The planting calendars correspond to areas which produce more than 95% of Brazilian agribusiness GDP that is about US\$165 billion/year (total GDP US\$450 billion/year). The federal government provided subsidiary support for agriculture since 1965. Funding levels were US\$8 billion for the 2004/05 cropping season, with US\$2.5 billion available for small farmers.

Agricultural zoning has been used as a federal farm credit policy since the rural lenders, mainly the Bank of Brazil, have had to use the planting calendars when supplying federal credit to farmers. This program has helped farmers to use proper technologies, protect the soil and the environment, plan their activities, decrease the production costs and risks, and increase the national production and productivity.

Brazil has now a global top-three position in production of ten commodities, including coffee, soya, sugar and maize.

Adapted mostly from: Zullo JJ, Pinto HS, Assad HD (2006). Impact assessment study of climate change on agricultural zoning. *Meteorol. Appl. (Supplement)*, 69–80 (2006) doi:10.1017/S135048270600257X



9. References and further reading

- Abrol, Y.P. & Ingram, K.T. (1996) Effects of higher day and night temperatures on growth and yields of some crop plants. In: *Global climate change and agricultural production. Direct and indirect effects of changing hydrological, pedological and plant physiological processes*. Bazzaz, F. & Sombroek, W. (eds). FAO, Rome and John Wiley & Sons, Chichester.
- Abrol, Y.P., Bagga, A.K., Chakravorty, N.V.K. and Wattal, P.N. 1991. Impact of rise in temperature on productivity of wheat in India. In: *Impact of Global Climatic Change on Photosynthesis and Plant Productivity*. Y.P. Abrol *et al.* (eds). Oxford & IBH Publishers, New Delhi.
- Aggarwal, P.K. & Sinha, S.K. (1993) Effect of probable increase in carbon dioxide and temperature on productivity of wheat in India. *Journal of Agricultural Meteorology*, 48: 811–814.
- Amponsah, P.E. 2004. Seismic activity in Ghana: past, present and future. *Annals of Geophysics*, 47: 539–543.
- Anon (2007) Report on Sectoral Climate Change Impacts, Vulnerability and Adaptation Assessments in Ghana: Climate Change Impacts on Root and Tuber Crops Production, Technical Summary, September 2007.
- AQUASTAT, Irrigation in Africa in Figures, 2005. Accessed August 2007 by Dazé.
<http://www.fao.org/ag/agl/aglw/aquastat/countries/ghana/index.stm>
- Baker, P. & Haggard, J (2007) Global warming; the impact on coffee. Presentation at the Speciality Coffee Association of America conference (May 2007). Long Beach. USA
- Baker, P (2009) “Climate change brews up trouble for coffee growers” -
<http://www.guardian.co.uk/environment/2009/aug/06/climate-change-coffee> and
<http://www.scidev.net/en/>
- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, M., Brown, V.K., Buse, A., Butterfield, J.E.L., Coulson, J.C., Farrar, J., Good, J.E., Harrington, R., Hartley, S.H., Lindroth, R.L., Lincoln, D.E., Press, M. & Watt, A.D. (2002) Herbivory in global climate change research. Direct effects of rising temperature on insect herbivores. *Global Change Biology* 8, 1–16.
- Battisti, D.S. & Naylor, R.L. (2009) Historical warnings of future food Insecurity with unprecedented seasonal heat. *Science*, 323: 240–244.
- Berry, P.M., Harrison, P.A., Hossell, J.E. & Masters, G.J. (2005) Introduction, project aims and methods. In: *Modelling Natural Resource Responses to Climate Change (MONARCH). A Local Approach. Development of a conceptual and methodological framework for universal application* Berry, P.M., Harrison, P.A., Dawson T.P. & Walmsley, C.A. (eds). Chapter 1, pp. 1–12.
- Berry, P.M., Masters, G.J., Harrison, P.A., Hossell, J.E., Crick, H.Q.P., Freeman, S.N., Ellis, N.E., Orr, H.G., Walmsley, C.A. & Watts, O. (2005) Conclusions and future research. In: *Modelling Natural Resource Responses to Climate Change (MONARCH). A Local Approach. Development of a conceptual and methodological framework for universal application*. Berry, P.M., Harrison, P.A., Dawson T.P. & Walmsley, C.A. (eds). Chapter 10, pp. 251–265.
- Boko, M., I. Niang, A. Nyong, C. Vogel, A. Githeko, M. Medany, B. Osman-Elasha, R. Tabo and P. Yanda (2007) *Africa. Climate Change 2007: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J., van der Linden and C.E. Hanson (eds). Cambridge University Press, Cambridge UK, 433–467.



Brown, V.K., Masters, G.J. & Cheeseman, O.D. (1999) Predicting the floral future of a warmer Britain. *Plantlife Summer 1999*, p. 15.

CABI Final Technical Report (2003) Water conservation in cotton ecosystems through farmer field schools. CABI Pakistan 1068. 22p

CABI Bioscience (2000) Learning to cut the chemicals in cotton; case studies and exercises in farmer focused cotton IPM from around the world. CABI Bioscience Technical Support Group to the Global IPM Facility, Ascot, UK/PAN-UK, London, 92 pp. ISBN 0 9521656 3.5

Chattopadhyay, N. (2008) Climate Change and Food Security in India. International Symposium on Climate Change and Food Security in South Asia, August 25-30, 2008 Dhaka, Bangladesh. http://www.wamis.org/agm/meetings/rsama08/S402-Chattopadhyay-Climate-change_Food-Security.pdf

Christensen, J.H., B. Hewitson, A. Busuioc, A. Chen, X. Gao, I. Held, R. Jones, R.K. Kolli, W.-T. Kwon, R. Laprise, V. Magaña Rueda, L. Mearns, C.G. Menéndez, J. Räisänen, A. Rinke, A. Sarr and P. Whetton, 2007: Regional Climate Projections. In: *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.) Cambridge University Press, Cambridge, United Kingdom.

CIA World Factbook, Ghana, accessed June 2007 by Dazé. <https://www.cia.gov/library/publications/the-world-factbook/geos/gh.html>

Cock, M. J. W. (2003) Biosecurity and forests: an introduction – with particular emphasis on forest pests. FAO Forest Health and Biosecurity Working Paper FBS/2E, 2003, 61 pp.

CRIG (undated) Vulnerability of agriculture to climate change – impact of climate change on cocoa production, Cocoa Research Institute of Ghana.

Dazé, A. (2007) Climate change and poverty in Ghana. CARE International, Southern and West Africa, 23pp. <https://www.care.dk/files/ClimateChangeandPovertyinGhana-countryprofile.pdf>

DfID (2005) Growth and poverty reduction: the role of agriculture. DfID policy paper, 54pp, DfID, London.

Easterling, W.E., P.K. Aggarwal, P. Batima, K.M. Brander, L. Erda, S.M. Howden, A. Kirilenko, J. Morton, J.-F. Soussana, J. Schmidhuber and F.N. Tubiello (2007) *Food, fibre and forest products. Climate Change 2007: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK, 273–313.

Erda, L., Hui, J., Wei, X., Shiming, M., Xiu, Y., Chunsen, M., Zhanyun, M., Heran, W., Yingchun, L., Xue, H., Yingyu, H., Liyong, X. & Lijun, L. (2008) Climate change and food security in China: Executive summary. Greenpeace China, Beijing, pp. 8.

Funk, C., Dettinger, M.D., Michaelsen, J.C., Verdin, J.P., Brown, M.P., Barlow, M. and Hoell, A. (2008) Warming of the Indian Ocean threatens eastern and southern African food security but could be mitigated by agricultural development. *PNAS*, 105: 11081–11086.

Ghana's Initial National Communication (GINC) to the United Nations Framework Convention on Climate Change (UNFCCC), 2000. In Dazé, A. (2007) Climate change and poverty in Ghana. CARE International, Southern and West Africa, 23pp. <https://www.care.dk/files/ClimateChangeandPovertyinGhana-countryprofile.pdf>



Grime, J.P., Brown, V.K., Thompson, K., Masters, G.J., Hillier, S.H., Clarke, I.P., Askew, A.P., Corker, D. & Kieley, J.P. (2000) The response of two contrasting limestone grasslands to simulated climate change. *Science* 289, 762-765.

Hansen, J.W. (2005) Integrating seasonal climate prediction and agricultural models for insights into agricultural practice *Philos Trans R Soc Lond B Biol Sci.* 360: 2037–2047.

IPCC (2007) Climate Change 2007: Synthesis Report. A contribution of Working Groups I, II, and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri, R.K. and Reisinger, A. (eds)]. IPCC, Geneva, Switzerland. 104pp.

IPCC (2001) Climate Change 2001: Synthesis report. A contribution of Working Groups I, II and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Watson, R.T. and the Core Writing Team (eds). Cambridge University Press, Cambridge, UK, 398pp.

Lobell, D.B., Burke, M.B., Tebaldi, C., Mastrandrea, M.D., Falcon, W.P. & Naylor, R.L. (2008) Prioritizing Climate Change Adaptation Needs for Food Security in 2030. *Science*, 319: 607–610.

Masters, G.J. & Brown, V.K. (2001) Effects of experimental manipulation of climate on calcareous grassland plants and invertebrates. In: *Impacts of Climate Change on Wildlife*. RSPB publication on behalf of EN, WWF-UK, UNEP WCMC & RSPB. Eds. Green, R.E., Harley, M., Spalding, M. & Zöckler, C. pp. 57–59.

Masters, G.J. & Ward, N.L. (2005) Implications for the composition of species communities. In: *Modelling Natural Resource Responses to Climate Change (MONARCH). A Local Approach. Development of a conceptual and methodological framework for universal application*. Berry, P.M., Harrison, P.A., Dawson T.P. & Walmsley, C.A. (eds). Chapter 4, pp. 68–73.

Masters, G.J., Brown, V.K., Clarke, I.P., Whittaker, J.B. & Hollier, J.A. (1998) Direct and indirect effects of climate change on insect herbivores: Auchenorrhyncha (Homoptera). *Ecological Entomology*, 23, 45–52.

Nicholls, T., Norgrove, L., & Masters, G. J. (2008) Innovative solutions to new invaders: managing agricultural pests, diseases and weeds under climate change. Crawford Lecture Series.

OECD-FAO (2008) *Agricultural Outlook 2008–2017*. OECD Publications, Paris, 73pp.

Pinto H.S., Zullo Jr J., Assa E.D., de Avila A.M.H. (2005). Global warming and future Brazilian agriculture scenarios. 17th Int. Congr, Biometeorology. *Annalen de Meterologie* 41 223–226.

Rao, D.G. and Sinha, S.K. (1994) Impact of climate change on simulated wheat production in India, EPA Conference, New Delhi, India, pp. 1–8.

Reddy, K.R., Doma, P.R., Mearns, L.O., Boone, M.Y.L., Hodges, H.F., Richardson, A.G., Kakani, V.G. (2002) Simulating the impacts of climate change on cotton production in the Mississippi Delta. *Climatic Research* 22: 271–281.

Rosenzweig, C., Tubiello, F.N., Goldberg, R., Mills, E. & Bloomfield, J. (2002) Increased crop damage in the US from excess precipitation under climate change. *Global Environmental Change* 12: 197–202.

Sinha, S.K., and Swaminathan, M.S. (1991) Deforestation, climate change and sustainable nutrition security. *Climatic Change*, 19, 201–209.

Stern, N. (2006) *The Economics of Climate Change: The Stern Review*. HM Government and Cambridge University Press, Cambridge.



United Nations (2008) *Millennium Development Goals Report (2008)* United Nations, New York, pp. 56.

World Commission on Environment and Development (1987) *Our Common Future*. Oxford: Oxford University Press.

World Health Organization (WHO) (2006) Country cooperation strategy: Ghana.
http://www.who.int/countryfocus/cooperation_strategy/ccsbrief_gha_en.pdf.



contact CABI

europa

CABI Head Office

Nosworthy Way, Wallingford, Oxfordshire, OX10 8DE, UK

T: +44 (0) 1491 832111

E: corporate@cabi.org

CABI Europe - UK

Bakeham Lane, Egham, Surrey, TW20 9TY, UK

T: +44 (0) 1491 829080

CABI Europe - Switzerland

Rue des Grillons 1, CH-2800 **Delémont**, Switzerland

T: +41 (0) 32 4214870

asia

CABI South Asia

Opposite 1-A, Data Gunj Baksh Road, Satellite Town, PO Box 8, Rawalpindi-Pakistan

T: +92 (0) 51 9290132

CABI Southeast and East Asia

PO Box 210, 43400 UPM Serdang, Selangor, Malaysia

T: +60 (0) 3 89432921

CABI South Asia - India

2nd Floor, CG Block, NASC Complex, DP Shastri Marg, Opp. Todapur Village, PUSA, New Delhi - 110012, India

T: +91 (0) 11 25841906

CABI Southeast and East Asia - China

C/o CAAS-CABI Project Office

C/o Internal Post Box 56, Chinese Academy of Agricultural Sciences, 12 Zhongguancun Nandajie, Beijing 100081, China

T: +86 (0) 10 62112118

africa

CABI Africa

ICRAF Complex, United Nations Avenue, Gigiri, PO Box 633-00621, Nairobi, Kenya

T: +254 (0) 20 7224450/62

americas

CABI Caribbean & Latin America

Gordon Street, Curepe, Trinidad and Tobago

T: +1 868 6457628

CABI North America

875 Massachusetts Avenue, 7th Floor, Cambridge, MA 02139, USA



T: +1 617 3954051