

**EUROPEAN ASSOCIATION OF ANIMAL PRODUCTION
EAAP
53rd ANNUAL MEETING, CAIRO, EGYPT 1-4 September 2002
SESSION: THE USE OF DRAUGHT ANIMALS IN SEMI-ARID
FARMING SYSTEMS**

Aspects of Work Animal Use in Semi-arid Farming Systems

B G Sims and D H O'Neill

International Development Group

Silsoe Research Institute, Silsoe, BEDFORD MK45 4HS, UK

Summary

Small-holder farmers in the semi-arid regions of many developing countries are faced with a multitude of problems including soil erosion and fertility decline, erratic rainfall and increasing risk. Semi-arid region farming systems incorporate animal and crop production. Oxen are the preferred work animals, but necessity may require the use of cows and equids. Harnesses and equipment for equids is often under-developed and programmes of participatory technology development have produced viable options. The combination of high-lift harnesses and light-weight equipment has been a particularly successful example. Smaller animals consume less fodder and so may allow more widespread use of conservation farming practices, including direct seeding. Hillside soil and water conservation can be achieved whilst producing dry season fodder via the use of vegetative contour liver-barriers.

Keywords: soil moisture, fertility and conservation; fodder, high-lift harness; light-weight tools; equids; live-barriers

Introduction

Farm families struggling to secure a livelihood in the semi-arid regions of the World are often confronted by multiple difficulties which can make their situation particularly precarious when compared with that of small farm families working in more favourable climatic regimes.

By definition, moisture availability is a critical constraint and is fundamentally affected by soil manipulation and cover, and weed populations. Having the right amount of moisture available to crops at the critical times in their cycles can make the difference between harvest and no harvest. Coupled with the scarcity of total moisture supply is the factor of the unpredictability of the quantities received. Without irrigation semi-arid region farmers are at the mercy of the vagaries of the weather, which is frequently notoriously difficult to predict and allow confident farm planning.

Soil fertility decline is a frequently expressed problem facing semi-arid region small farmers. Crop yields decline year after year, more "thorough" cultivation increases organic matter oxidation resulting in soil fertility and water-holding capacity continuing on their downward slide. Regions with broken terrain, making it necessary

to cultivate on sloping land, have additional problems. Soil erosion resulting from heavy rainfall on cultivated soil bereft of vegetation will result in the preferential loss of the lighter soil particles which are also those richest in plant nutrients – or soil fertility.

As if these woes were not enough, semi-arid region small farmers are often forced to farm their marginal areas because of pressure on more fertile land by other sectors of society, sectors with more political muscle. The outcome is that small farmers, attempting to produce from marginal land in the first place, frequently find themselves marginalized politically, socially and economically as well. This can have a negative effect on access to markets, access to productivity enhancing inputs (seeds, fertilizer, crop-care products, etc), and still further marginalization.

Where animals are kept, either for production or for work, the pressure on the land is increased still further. Forage production will tend to be low *per se* and so will lead to the removal of all crop residues. Bare soil surfaces at the beginning of the rainy season exacerbate the erosion problem resulting in an accelerated loss of soil fertility.

These conditions constitute a challenge for the small farm family and the R&D community working to produce solutions to aid the families to farm sustainably whilst improving their prospects for better livelihoods. This paper examines some aspects of the work animal component of semi-arid farming systems, it looks at some of the problems associated with their use and suggests some solutions. The paper reflects and draws on the results of many years of work with small farmers in the semi-arid regions of Latin America and sub-Saharan Africa.

Semi-arid Farming Systems

Semi-arid agricultural production systems can be conveniently characterised as having 400 – 1200 mm annual rainfall, a mean air temperature higher than 18 C and having at least one season when evapotranspiration exceeds precipitation (LPP, 2001). Such circumstances lead to a shortage of water, or moisture, which is a major constraint on production, particularly crop production. No particular type of soil is especially prominent in the semi-arid areas that are farmed but, with conditions not generally promoting the growth of large volumes of vegetation, the soils are not rich in organic matter. They, therefore, tend to be infertile, degraded and have low water retention which, in many cases, exacerbates the shortage of rainfall.

Many, if not most, semi-arid farming systems depend on an integrated approach to crop and livestock production. With relatively poor soil, smallholder farmers being unable to achieve high levels of productivity need to cultivate fairly extensively to harvest sufficient for their family needs and market, if appropriate. It is attractive, therefore, for them to cultivate their plots using draft animal power (DAP) rather than depending only on human labour to increase the area cultivated and reduce drudgery. (Tractors are very rarely an option for smallholders). However, more extensive cropping may introduce problems with the timeliness of planting and the demands of crop care. Yield is forfeited if late planting results in the scarce rainfall not being fully exploited or if the weed infestation is too severe. A surprisingly small proportion of smallholders, even those with access to DAP, enhance their weeding with draft animal technologies. Two of the more likely explanations are that seeds

may have been broadcast (rather than planted in lines) and that weeding is regarded as a woman's job (men tend to control the draft animals and their use).

The basis of crop-livestock integration is that animals assist with crop production tasks (mainly land preparation) and transport, whilst the crop residues are used as fodder which, together with grazing, enable the animals to survive and provide organic fertilizer (manure). Animals, especially cattle, are important culturally in many semi-arid farming systems and their value provides a means of holding wealth with high liquidity.

Semi-arid farming, especially crop production, is risky primarily because of the combination of low rainfall and high temperatures creating crop moisture stress. Furthermore, semi-arid tropical climates tend to be characterised by erratic rainfall patterns leading to periods of water-logging and "mini-droughts", neither of which are conducive to prolific and healthy crop growth and reliable yields. As one means of managing, or ameliorating, these risks, smallholders may do their planting at different stages into the growing season and thereby hope to achieve a worthwhile total yield, although it may not be optimal. Other ways of reducing the risk of yield loss from mini-droughts include water conservation. These are generally fairly arduous soil and water management practices which can be facilitated by the use of DAP. Many problems could also be overcome by the use of an irrigation system but conventional systems demand a good, stable water source and a significant investment well beyond the reach of typical smallholder farming families. Such families depend on human labour but this is usually found wanting during periods of peak demand (planting, harvesting and especially weeding) and reliance on the extended family is a common solution. Households help each other out when tasks need to be done. Very little money changes hands and families are rewarded for their efforts by meals or reciprocal labour arrangements. Only the wealthier families can afford to hire labour, or draft power, for cash.

Solutions are urgently needed to enable smallholders to increase both their land and labour productivity whilst making more sustainable use of their natural resources for growing crops and raising livestock. A diversified use of draft animals, coupled with a move away from traditional land management to enhance water (rainfall) use and reverse declining soil fertility, would seem to offer a promising approach to reach a solution. For such an approach to succeed, there is a need for better information on the matching of draft animals, animal-drawn equipment, soil manipulation and moisture management within the context of the declining productivity of local farming systems and practices.

Work Animals

There are no major physical differences between draft animals and others of the same species, although certain breeds are more renowned for their draft capabilities. Almost any bovine (including buffaloes), equid or camelid can become a draft animal, provided that it is reasonably healthy and responds to training. In general, the most important criterion is body mass, or liveweight. This gives an indication of the amount of muscle on the animal and, thus, its potential to exert a force or, more specifically, a pull which the farmer can utilize through soil-working implements. A basic guide is that an animal can pull about 10 to 15 % of its weight for a working

period of around 4 hours, although there is some variation between species, breeds and working conditions. Summaries of the approximate work capabilities of different species are given in Table 1.

Table 1. Draft capability and power outputs of various animals.

Animal	Average weight (kg)	Approximate draft capability (N)	Average speed (m/s)	Power developed (W)
Bullock	500 - 900	600 - 800	0.56 - 0.83	560
Cow	400 - 600	500 - 600	0.70	340
Water buffalo	400 - 900	500 - 800	0.80 - 0.90	560
Light horse	400 - 700	600 - 800	1.0	750
Mule	350 - 500	500 - 600	0.9 - 1.0	520
Donkey	150 - 300	300 - 400	0.70	260
Camel	450 - 500	400 - 500	1.1	500
Man	60 - 90	300	0.28	75

Source: Campbell, 1990

The total amount of work that can be done depends on the animal's energy reserves and the onset of fatigue which, in turn, depend on the animal's condition and nutritional status. The energy cost of work can be found by measuring the animal's metabolic rate whilst working. The most direct determinant of metabolic rate is the animal's oxygen consumption, but this is not easily measured, especially in field conditions. A less direct but more accessible determinant is heart rate, which varies linearly with metabolic rate.

Any large domestic animal may be used for draft and farmers will choose what fits in with their system of farming from what is available and affordable. For the crop / livestock farmers of sub-Saharan Africa who, typically, own cattle, oxen would be the first choice. But, in the areas of southern Africa which have been most seriously affected by droughts in the last decade where many cattle have perished, donkeys are increasingly being used for crop production because, for smallholders, they are becoming more available and affordable than cattle. In the drier Sahelian region, camels would be the first choice but for how much longer smallholder farmers will be able to afford them is questionable.

The physical / physiological state of an animal, including its fitness for work, can be judged reasonably effectively by visual examination. Experienced farmers, who know their animals, have, of necessity, developed such a skill and now systems of body condition scoring have been developed. These are mainly to help extension workers and researchers judge the condition of animals, and have the added advantage of providing a consistency of appraisal. The body condition score reflects, primarily, how healthy an animal appears and integrates its level of nourishment and the presence of any obvious disease or injury. One example is that proposed by Nicholson and Butterworth (1986), for oxen, which runs from 1 to 9, representing emaciated to obese. For work, body condition scores of between 4 and 6 would seem to be optimum. Oxen with a body condition of more than 6 may be too overweight to

give optimum performance and may be more susceptible to heat stress than leaner animals.

A number of factors must be considered with regard to the feeding of working animals. Does the working animal have enough time to feed? Is enough feed available and is it of sufficient quality in terms of both energy content and digestibility? Does work affect appetite and the passage of food through the body? If so, in what way and to what extent? For more details, see Pearson (1996). For horses, oxen and buffalo, most of the questions relating to feed and feeding practices can be answered, but, for donkeys and camelids, less research has been done and recommendations are less well defined. For camels, in particular, the issues are further complicated by the camel's ability to dehydrate and thereby modify its metabolic processes.

There is an almost universal complaint from smallholder farmers who rely on DAP that there is insufficient to meet everyone's needs. Traditionally, farmers who have used animals for draft work have kept some males specifically for this purpose, but increasing pressures on land and feed, together with the underlying costs of maintaining and maybe purchasing them, have prompted a change. This is particularly the case for farmers who own cattle, who have tried to economize by using cows for work. It is still rare to find a farmer spanning or yoking cows exclusively but it is not uncommon to find cows being spanned with oxen to make a team of two or four, where the farmer cannot afford to buy or maintain a complete team of males. This raises the question of how much work a female animal can do before her milk production or, more seriously, her fertility is affected. If a female draft animal is worked to the point of infertility, the outputs of milk and calves (or foals) are lost and the farmer has lost an asset of greater potential value than a working animal. Recently, research has been undertaken to evaluate the effects of using cows for draft but the results have not been conclusive. It is clear, however, that when milking cows are used for work supplementary feeding is essential.

Implements and Harnesses

The potential for using lighter, smaller animals for draft work on the farm means that less fodder needs to be consumed, with important implications for cropping systems and protection against land degradation. But it also raises challenges for making the use of the available, reduced, power source more efficient. In many semi-arid regions, oxen remain the preferred farm-power source. However their use for a few days a year, principally on tillage and transport work, imposes a heavy forage burden throughout the year, and this can be especially difficult to meet in the dry season. Figure 1 shows the metabolic energy balance for draft oxen in the semi-arid inter-Andean valley region of Bolivia, and clearly indicates the period when energy demand exceeds the available supply.





Figure 1. Metabolic energy balance (MJ day^{-1}) for oxen during the year.

The consequences of seasonal forage shortage on farming systems and livelihoods can be quite dramatic. It is frequently necessary for farmers to sell their animals after the main soil cultivation season, and to buy new animals the following year. This involves journeys to the markets, uncertain prices, training new draft animals and risk.

Conversations with farm families has led to the suggestion that other, lighter, animals could possibly fulfil the functions of the heavier oxen. Equids (donkeys, mules and horses) are frequently already available as a transport option. However appropriate harnesses and light-weight equipment have not often been sufficiently developed. Whereas research can quite easily provide technical solutions, the existence of a manufacturing infrastructure is a vital ingredient in the process of successful development of adoptable technologies.

A process of participatory technology development in Bolivia (LPP, n.d.) worked with farm families on this theme and in conjunction with a local draft animal implement factory. Farmers recognised the untapped potential of their equines, but pointed out that existing draft animal tillage equipment was too heavy for use with lighter animals as it had been designed for use with pairs of oxen. Taking as a starting point Frank Inns' work on high-lift harnesses for reducing the draft of chain or rope-pulled implements (Inns, 1990 and 1991)¹ we were able to demonstrate that increasing the pull angle of the harness results in the reduction of the effective vertical force on the implement (by reducing parasitic soil/implement friction forces, and the effective implement weight). As a rule of thumb, by increasing the angle of pull from 15° to 30° it is possible to reduce the implement draft by up to 50% (Figure 2).

¹ And, in fact, incorporating Frank Inns into the R&D team



This harness gives an angle of pull of about 30° or slightly more

Figure 2. The force system acting on a draft animal. Increasing the angle of pull (α) will reduce the implement draft.

The materials used should be non-abrasive and padded where load is applied (i.e the hip and shoulder straps and, especially, the breast band). Local materials should be used and, once the principles have been well understood, local adaptation will be expected (Figure 3).



Figure 3. A horse using a high-lift harness made from fertilizer sacks. Capinota, Bolivia

The sustainable pull that an equid can produce will depend on its disposition, nutrition, health and physical condition, but also, crucially on its body weight. Equids can typically sustain pulls in the region of 10-15% of body weight which, for a 150 kg donkey translates to a pull force of 150–220 Newtons. This will usually still not be a high enough value for sustained pulling of equipment designed for pairs of draft oxen, and so lighter equipment needs to be designed and tested to ensure that it is within the capacity of the work animals. In the case of moldboard plows, for instance, we have found that a fit donkey can handle a plow with a share width of some 115 mm and this increases for single horse to 150 mm width.

However, although one important aspect of work animal use in semi-arid regions is the requirement to use less fodder, another is to reduce the amount of tillage and to increase rainwater infiltration.

Work in sub-Saharan Africa (Sims and Twomlow, 2000) has shown that the major problems associated with conservation tillage approaches currently promoted to the smallholder farmer are weed control, crop establishment and shortages of adequate draft power. To date little work has been carried out to assess the interaction of different conservation tillage options with different weeding regimes on maize yield and soil-water regimes. Therefore, it is essential that conservation tillage practices are developed that both conserve water, reduce draft animal power inputs, encourage timely crop establishment and weed control systems that take into account the resourcefulness of smallholder farmers.

The advantages have been shown of establishing a crop within either a rip line (with a narrow tine) or a planting furrow opened with a plow on winter or previously spring plowed land (Muza *et al.*, 1996). Although these techniques of crop establishment can reduce power inputs for seedbed preparation and allow timely planting, a heavier and earlier weed burden results compared with that for overall plowing and planting together, the prevailing practice in sub-Saharan Africa (Mabasa *et al.*, 1999). If combined with weeding with a plough (or a cultivator with ridging blades), timely planting, better soil water conservation, weed control and labour reduction could be achieved.

Recent work in Zimbabwe has focussed on developing weed control strategies that complement primary tillage techniques for the majority of communal area farmers who cannot afford purchased inputs, including herbicides. Because of poor returns to cropping and an acute shortage of labour in many households, tillage / weed control systems need to be based on low cost, labour saving technologies (Ellis-Jones and Mudhara, 1997). Riches *et al.*, (1997) reported that the use of the moldboard plow with body attached during weeding allows the creation of a ridge and furrow landform that can enhance soil water retention. The soil thrown towards the crop row smothers weeds and reduces the need for subsequent labour-intensive inter-plant weeding. Labor productivity, in terms of grain harvested, can be considerably higher with this technique than the use of existing hand-hoe or cultivator followed by hoe systems. While 76% of households own a plow in southern Zimbabwe, only 23% own a cultivator, so “plow-weeding” provides an opportunity for increasing the timeliness of weed control, for farmers who currently weed by hand, without the need for additional capital investment, if they have access to work animals.

Reduced vertical tillage has also been a feature of participatory technology development efforts in Latin America. Chisel plows to improve run-off infiltration on hillsides have been developed to be more efficient in terms of volume of soil moved per unit of energy imparted. One is a scaled-down version of a plow designed for oxen and drawing heavily on the design of the traditional ard-type plow. Reducing the width of the pointed share, and the width of the twin moldboards, will bring the implement to within the capacity of equines.

Narrow tines with a low rake angle have long been used for bursting and mixing soil in a limited width of work (Spoor, 1969), and in fact this is the basic principle on

which and plows function. To increase the efficiency of a chisel plow, fitting inclined wings to the rear of the chisel point has been found to be effective (Spoor and Godwin, 1978). Figure 4.



Figure 4. Wing-tined chisel plow for more efficient soil bursting during vertical tillage.

Surface cover is one of the best soil conservers and improvers that can easily be adopted by semi-arid region farmers. A problem arises, of course, over the competing demands for animal fodder, but the value and potential of mulch-based no-till agriculture has been well demonstrated (Wall *et al.*, 2001). Although no-till planting with animals is common practice in several Southern Cone countries, light weight, multi-row equipment for small grains was not available in the semi-arid valleys of the Andean region. Figure 5 shows a late prototype that has been developed for small cereals and is on trial in semi-arid regions of Bolivia, Mexico and India.



Figure 5. Direct seeder for small cereals designed to be pulled by a pair of oxen. Bolivia.

CHALLENGES FOR THE FUTURE

The principal aim of R&D efforts expended on semi-arid agriculture is to contribute to the resilience of the farming systems and to contribute to improved sustainability.

This in turn will add to the farm families' capital stock and contribute to an improvement in livelihoods.

Crop production is more uniform and less risky if soil is conserved and fertility maintained or improved. Capturing and directing run-off is a fundamental requirement for crop production in this hostile environment. We have seen some efforts that have been made to reduce tillage and promote vertical tillage, other approaches include using micro-relief for water capture. The direct seeder shown in Figure 5, for example, can be adapted to furrow-bottom planting of small grains with remarkable yield increases. Larger scale physical works and micro-catchments (earth dams and so forth) are also possible to construct with the aid of animal power and can provide sufficient supplementary irrigation in dry years to prevent total crop loss (Pacey and Cullis, 1986).

Domesticated animals are an integral part of most semi-arid farming systems, but they are also a source of land degradation and threaten the cropping area if the demand for dry-season fodder exceeds supply. Keeping fewer, lighter working animals may go some way towards ameliorating this situation and so an increase in the design, manufacture and supply of lightweight implements is a probable future development. The challenge is to develop this equipment in close collaboration with potential users and to produce prototype designs as a result of participatory (scientist and farmer) technology development.

Soil and water conservation, generally will be a priority for the future. With our present programme of World environmental destruction, forest devastation and population growth set to reach 9 billion (from the present 6 billion) in less than 50 years, and the seeming lack of worldwide concern for the damage that we are perpetrating, the situation is bound to get worse. Conservation tillage (including direct planting) will clearly have a role to play in conserving soil and water in semi-arid regions, but so, also, will conservation measures which respond to farm families' needs by, for instance, producing useful products whilst protecting the soil.

Recent work in the Bolivian semi-arid region (Rodríguez, and Sims, 2001) has produced, in collaboration with farm families, a variety of suitable species for contour-planted, soil and water conserving, vegetative live barriers which produce abundant dry season fodder whilst conserving fragile hillsides (Figure 6).



Figure 6. Contour barriers of *phalaris* grass (*Phalaris tuberoarundinacea*) and woolly-pod vetch (*Vicia sativa ssp. dasycarpa*) for soil conservation and forage production.

Acknowledgement

The technologies discussed in this paper are outputs of projects funded by the Rural Livelihoods Division of the UK Government's Department for International Development via their Crop Protection Programme (Project R7579) and their Livestock Production Programme (Projects R6970; R7401 and R7352). The views expressed are not necessarily those of DFID.

REFERENCES

- Campbell J K, 1990. Dibble sticks, donkeys and diesels. International Rice research Institute, Philippines. 329pp ISBN 971-104-185-5
- Ellis-Jones, J & Mudhara, M. 1997. Conservation Tillage for Resource Poor Farmers: The Critical Importance of Farm Power. *Bibliotheca Fragmenta Agronomica*, 2A/97 (presented to 14th ISTRO Conf. Agroecological and Ecological Aspects of Soil -Tillage). Pulawy, Poland.
- Inns, F. 1990. The mechanics of animal-draught cultivation implements. *The Agricultural Engineer* 45 (Spring 1990): 13-17.
- Inns, F. 1991. The design and operation of animal / implement systems: guidelines for soil cultivation implements. In: Fielding, D. and Pearson, R.A. (eds) *Donkeys, mules and horses in tropical agricultural development*. University of Edinburgh, Centre for Tropical Veterinary Medicine. pp 258-265. ISBN 0907146066.
- LPP, (undated). Management and use of draught equines by poor crop/livestock farmers. Project R6970. Chatham Maritime, Kent, UK. NR International. Livestock Production Programme. 2p.
- LPP. 2001. Livestock Production Programme. Project year book 2001-2002. Chatham Maritime, Kent, UK. Natural Resources International. Department for International Development. No page numbering.
- Mabasa, S. Riches, C.R, Nyahunzvi, S., Twomlow, S.J, Dhliwayo, H.H., Chatizwa, C., 1999. Tillage and Weed Control Interactions on A Semi-arid Granitic Catena. II. Weed Responses. In: CIMMYT and Ethiopian Agricultural Research Organization (Editors)). *Maize Production Technology for the Future: Challenges and Opportunities: Proceedings of the Sixth Eastern and Southern Africa Regional Maize Conference, 21-25 September, 1998, Addis Ababa, Ethiopia*.
- Muza, L. Dhliwayo, H. & Twomlow, S.J., 1996. Dryland maize response to different combinations of tillage and weeding methods. In: Ransom, J.K., Palmer, A.F.E., Zambezi, B.T., Mduruma, Z.O., Waddington, S.R., Pixley, K.V. & Jewell, D.C. (Eds) *Maize productivity gains through research and technology dissemination Proceedings of 5th Regional Maize Conference for*

- Eastern and Southern Africa, Arusha, Tanzania, pp 3-7. Addis Ababa, Ethiopia, CIMMYT. 110-114.
- Nicholson, M.J. and Butterworth, M. 1986. A guide to condition scoring of Zebu cattle. ILCA (International Livestock Center for Africa). Addis Ababa, Ethiopia. 36p.
- Pacey, A and Cullis, A. 1986. Rainwater harvesting. The collection of rainfall and run-off in rural areas. London. Intermediate Technology Publications. 216p.
- Pearson A, 1996. Feeding strategies for cattle and buffalo used for work. World Animal Review 87, 45-55. FAO, Rome.
- Riches, C.R., Twomlow, S.J. and Dhliwayo H.H. 1997 Low-input weed management and conservation tillage in semi-arid Zimbabwe. *Experimental Agriculture*, 33, 173-187.
- Rodríguez, F. and Sims, B.G. (2001). Barreras vivas asociadas para la conservación de suelo, agua y producción de forraje. Estrategias para la producción de forraje y control de erosión como complemento del manejo de malezas en laderas (PROFOCE). Department for International Development (DFID); Universidad Mayor de San Simón, Cochabamba, Bolivia. 4p.
- Sims, B.G., Twomlow, S.J. and Ellis-Jones, J. 2000. Evaluation of soil and water conservation practices for small farmers in marginal areas. Milwaukee, Wisconsin, USA. ASAE Annual International Meeting. Paper No. 002211. 15p.
- Spoor, G. 1969. Design of soil engaging implements. Parts I and II. London. *Farm Machine Design Engineering* 3:22-25, 28.
- Spoor, G and Godwin, R. J. 1978. An experimental investigation into the deep loosening of soil by rigid tines. *Journal of Agricultural Engineering Research* 23:243-258.
- Wall, P.C., Zambrana, L., Gámez, P, Sims, B. and Callisaya, A. 2001. Development of an animal drawn zero tillage seeder for small grains. Madrid, Spain. Xul. I World Congress on Conservation Agriculture. pp 631-635. ISBN 84-932237-2-7.