

Linking the hydrological cycle and rural livelihoods: A case study in the Luvuvhu catchment, South Africa

(ZF0150/R7937)

Hope, R.A.², Jewitt, G.P.W.¹ and Gowing, J.W.²

- (1) School of Bioresources Engineering and Environmental Hydrology, University of Natal, South Africa.
- (2) Centre for Land Use and Water Resources Research, University of Newcastle, Newcastle upon Tyne NE1 7RU, UK.

Abstract

The role of the hydrological cycle in contributing to the livelihoods of rural communities is often said to be important, but clear evidence of this is rarely offered. Furthermore, where such aspects are considered, they are largely focussed on the use of water from rivers, boreholes or some form of storage (blue water).

In this study, the hydrological cycle is considered in its entirety. Links between rural livelihoods, land use and the goods and services provided by the evaporation and transpiration components of the hydrological cycle (green water) are assessed through analyses of rural livelihoods in the Luvuvhu catchment, Limpopo Province, South Africa. Results highlight the importance of green water, and thus the importance of access to land and use of the natural resource base, in disaggregated rural community livelihood strategies

Finally, we describe a methodology for linking common outputs from hydrological models to rural livelihood impacts. In this way, the potential role of land use change in disaggregated rural livelihoods can be assessed for various development scenarios, such as increases in commercial afforestation and dryland agriculture.

Introduction

Society depends on the integrity of natural systems to provide goods and services for production and consumption. In particular, the goods and services generated by the movement of water through the landscape are assumed to make significant contributions to the livelihoods of rural communities. However, clear evidence is rarely offered to substantiate this position; the general tendency is to focus on the use of water from rivers, boreholes or some form of water storage for irrigated crop production. The omission of the contributions of other land uses under natural (rainfed) conditions has been emphasised as a major weakness in the analysis of land, water and development interactions.

Land use influences catchment hydrological responses by partitioning rainfall between return flow to the atmosphere as evaporation and transpiration ("green water") and flow to aquifers and rivers ("blue water"). Increasing attention is now being paid to understanding the dynamic inter-relationships between "green" and "blue" water, as these are considered to underpin essential terrestrial and aquatic ecosystem services (FAO, 2000; Jewitt, 2002; Falkenmark, 2003). The FAO suggests that almost all "green" water and a large proportion of "blue" water are needed to sustain ecosystem structures and functions, and maintain sustainable water supplies.

In this paper, links between rural livelihoods, land use change and the goods and services provided both green and blue water components of the hydrological cycle are assessed in the Luvuvhu catchment, Limpopo Province, South Africa. Specifically, a methodology to link the impacts of scenarios of land use change, such as, conversion of natural vegetation to commercial afforestation and irrigated agriculture on catchment hydrological functioning and rural livelihoods is illustrated in terms of green and blue water use. Furthermore, the analysis shows the impacts of land use change on poverty as a means of informing policy development (forestry, land affairs, water resources, institutional governance, compensation mechanisms, agriculture etc.) in an integrated and practical manner.

Land and water interactions and ecosystem goods and services

Movement of water through the landscape contributes to production (e.g., crops, timber, cattle), information (e.g., nature experiences), and regulation (e.g., formation of topsoil), functions of the

environment. Several authors have highlighted the importance of dryland crop production in rural areas and thus the importance of the water vapour phase (green water) of the hydrological cycle to rural livelihoods (e.g. Rockstrom et al, 1999). Typically, communities seek to optimize the conversion of available water, in the form of rainfall or irrigation water, through crop production resulting from the movement of green water through agricultural crops. However, natural resource harvesting of non-agricultural goods and services is also an important component of rural livelihoods.

Water both sustains and constrains land use, which in turn, influences catchment hydrological functioning by partitioning incoming precipitation into runoff, ground water recharge, interception and transpiration. Changes in upstream land use and resulting alterations in the movement of water through vegetation and air will impact on the downstream flow of water¹. An issue that has been highlighted, particularly in arid and semi-arid regions where green water flow dominates the hydrological cycle, is the potential for change in upstream land use and a resulting impact on the downstream flow of water. However, the impact of such land use change on the goods and services generated from green water flows is rarely considered explicitly.

Luvuvhu catchment

The Luvuvhu Catchment (5 941 km²) is located in Limpopo Province, South Africa, and together with the Letaba Catchment forms the Luvuvhu/Letaba Water Management Area (WMA), one of 18 WMAs identified by the South African Department of Water Affairs and Forestry (DWAf). It forms part of the larger Limpopo system, which drains into the Indian Ocean in southern Mozambique. For water resources management purposes, the catchment has been subdivided into 14 quaternary catchments² (Fig 1). Mean annual precipitation (MAP) over the Luvuvhu catchment is 608 mm, mean total evaporation (ET) is 1 678 mm and natural mean annual runoff (MAR) is estimated to be 520 x10⁶ m³. However, all of these show high spatial and temporal variation with highest rainfall and lowest ET over the Soutpansberg mountain range in the west and the lowest rainfall and highest potential ET in the arid areas in the west of the catchment adjacent to the Kruger National Park. A LANDSAT TM (1996) coverage provided by the Council for Scientific and Industrial Research (CSIR) has been used to estimate current land use in the catchment.

It is estimated that some 624 907³ people, including the populations of Thohoyandou (pop. 130 000), Louis Trichardt (pop. 88 000) (via inter-basin transfer) and Malamulele (pop. 2 000) depend on the basin for their water needs. Urban and industrial use was estimated at 6% of total water demand in 1997. However, it is predicted that this could reach 13% by 2010. Water use by commercial afforestation is 10%. A low confidence estimate of ecological water requirements of 105 x 10⁶ m³ per annum or 20% of MAR means that there is no surplus yield available in the water management area and that an over commitment of resources occurs. The bulk of irrigation water used in the catchment is supplied either from the Albasini Government Water Scheme or from private dams. In contrast to many other areas of the country, significant areas rely on groundwater for irrigation. DWAf (2002) report that ground water is extensively over-exploited, particularly in the vicinities of the Albasini Dam and Thohoyandou. Many small-scale irrigation schemes which utilize run-of-river flow and do not rely on an impounded water supply exist. It has been estimated that combined abstractions utilize all of the low flows in the river, particularly during the critically dry period of August to November.

Available soil surveys, indicate that 17% of the soils of the basins (960 280 ha) are potentially suitable for afforestation. However, only 14% of this area (14 750 ha) has been planted. Rainfall constraints, as well as difficulties faced in obtaining a Stream Flow Reduction Activity (SFRA) water use licence are the primary reasons for this. However, the potential for additional commercial afforestation does exist, and scenarios for such expansion form an important component of this study.

The upper catchment of the Mutale River, an important tributary of the Luvuvhu, and classified as quaternary catchment A92A by DWAf, forms the focus of this study. This is a relatively high MAP catchment (879mm) and includes areas of commercial afforestation, dryland and irrigated agriculture. A

¹ Change the partitioning of green and blue water in the catchment and may result in changes in livelihood responses at both community and catchment scale.

²The SA Department of Water Affairs and Forestry subdivides the countries catchments into quaternary catchments for water resources management purposes.

³ CSIR estimate from national census data.

summary of existing land uses for both the Luvuvhu and upper-Mutale catchments and potential scenarios of land use change are summarised in

Table 1.

Hydrological Modelling in Luvuvhu

The ACRU Agrohydrological modelling system has been configured for use in the Luvuvhu Catchment. The ACRU model is a multi-purpose and multi-level integrated physical conceptual model that can simulate streamflow, total evaporation, and land cover/management and abstraction impacts on water resources at a daily time step (Schulze, 1995). The model revolves around multi-layer soil water budgeting: streamflow is generated as stormflow and baseflow dependent upon the magnitude of daily rainfall in relation to dynamic soil water budgeting. Components of the soil water budget are integrated with modules in the ACRU system to simulate many other catchment components including irrigation requirements and sediment yield. Spatial variation of rainfall, soils and land cover is facilitated by operating the model in "distributed" mode, in which case the catchment to be modelled is sub-divided into sub-catchments. Within these sub-catchments, units of similar hydrological response, based largely on land use zones, are designated to facilitate simulation of land use changes, or isolation of, for example, riparian zones. Land cover and land use affect hydrological responses through canopy and litter interception, infiltration of rainfall into the soil and the rates of evaporation and transpiration of soil water from the soil and the model has been shown to be highly effective in this regard (Jewitt and Schulze, 1999).

The principal applications of the model are in the assessment of environmental and land use related impacts for water resources planning. Many verification studies have been performed, both on internal state variables and final model output (Schulze, 1995). In particular, forest water use and the impacts on streamflows have been verified at several locations in South Africa (Jewitt and Schulze, 1999).

Rural livelihoods and poverty

Approximately 75% of the 1.2 billion poor live in rural areas of the developing world (IFAD, 2001). In South Africa, of the 50% rural population 72% are considered poor (May, 1999). Rural livelihoods in the extensive semi-arid and arid areas of the developing world are influenced by their access to and use of local natural resources (e.g. water, rangeland⁴ etc.). A common perception that the natural resource base is unproductive is based more on reductionism⁵ than reality (Scoones et al., 1992; Guijt et al., 1995; FAO, 2001). Understanding how livelihood strategies are mediated by access to and use of water-dependent resources will allow a better understanding of how to reduce rural poverty. Analysis that includes common water-dependent livelihood activities may provide clearer direction on the trade-offs and likely outcomes of catchment water resources management, strategies for poverty reduction, hydrological integrity and economic development.

Livelihood analysis was conducted in Limpopo Province as it is one of the poorest of the nine South African provinces (Table 2). It has an estimated 4.9 million inhabitants across 123,910 km² representing 10% of the total land mass. It is the least urbanised province with a majority black population (96%). Across a range of development indicators the provincial status is poor in the national context, though its regional standing in the Southern African Development Community (SADC) is far better.

Livelihoods research in the Luvuvhu catchment is based on a household questionnaire (Jan, 2002; n=552), informant inquiry across 4 communities (Sept-Dec, 2002) and reference to secondary data sources⁶. Data are disaggregated by reported household income quintiles and are weighted by inverse probability of selection. Selected poverty indicators from the household survey are presented in Table 3.

⁴ Here, defined as communal land under Tribal Authority *de facto* management. Often, under mixed and complex open access, communal access and/or private tenure systems. The bulk of rangeland in the Luvuvhu falls within the former homeland/Bantustan of Venda.

⁵ The practice of analysing and describing a complex phenomenon in terms of its simple or fundamental constituents, especially when this is said to provide a sufficient explanation.

⁶ Shackleton's (1999a; 1999b; 2001) data from Limpopo province research into livestock and rangeland resources has been adapted with inputs from the Crookes et al. (2000) study.

Six water-dependent livelihood activities that conform to wider local, national and regional practices are identified for the analysis⁷: **1**) Water collection;; **2**) Livestock rearing (cattle and goats); **3**) Kitchen garden farming⁸. (0.04 ha; irrigated by IWS); **4**) Dryland farming (maize); **5**) Non-edible woodland products (NEWP: fuelwood, construction etc); and **6**) Edible woodland products (EWP: wild fruit and vegetables, herbs, worms etc).

The research quantifies the six activities by five criteria: **1**) Gross economic value (US\$/pa); **2**) Gross costs (US\$/pa); **3**) Net economic value (US\$/pa); **4**) Green water use (m³/pa); and **5**) Blue water use (m³/pa).

The opportunity cost of fuelwood gathering, water collection, subsistence farming and other natural resource harvesting activities is often described as a constraint to livelihoods in relation to children missing school, physical hardship and livelihood vulnerability due to illness or loss of an able-bodied family member (daughter/son leaves, father dies etc.). Whilst these very real costs are borne by the poor (including women and children in wealthier households) it is important to understand how these costs are distributed (Hope and Gowing, 2003; Budlender and Brathaug, 2002; StatsSA, 2001). This study has applied an effective local wage rate (US\$0.25 per hour) for these productive activities in order to compare it (gross costs) against the estimated values from water-dependent activities (gross value). With an unequal distribution of services (electricity, IWS, etc.) in rural areas it is highlighted which groups (here, income quintiles) have to bear the greater (opportunity) cost for not having access to these services.

Linking livelihood impacts and water use

The upper-Mutale case study represents an area of the Luvuvhu Catchment with a relatively high diversity of land uses and corresponding livelihood strategies. A weighted proportion of income quintiles in this quaternary catchment was generated from results of a survey from one of the communities sampled in the questionnaire.

Livelihood water use is estimated according to water use relative to a "natural" baseline condition represented by 100% cover with the dominant natural vegetation of the area (here, rangeland). In addition to the current land use in the catchment, the hydrological response of four additional land use scenarios was simulated with the ACRU model. In all scenarios, except the baseline, domestic blue water use is calculated at 25lcd for a population of 36 790. Only water collection remains unchanged in each scenario, though risk analysis of 'blue' water availability will assess the conditions under which dry season flow may be reduced to a level which fails to meet the specified basic human needs requirement (BHNR). These scenarios provide for a non-validated estimate of the direction and scale of water and economic impacts on the livelihood quintiles. It must be noted that complicated social, political and institutional factors that are likely to drive such land use change are not considered, as it is the management of such factors at policy level which this research is intended to inform.

Informant inquiry, secondary sources and careful deliberation are combined to assess how the land use change scenarios will affect livelihood water-dependent activities. Here, the following methodology has been followed with its inherent limitations⁹:

- Water collection remains static in each livelihood scenario analysis though a risk assessment is made within the hydrological analysis;
- Land, labour and other variables that contribute to kitchen garden and dryland agriculture are proportionally adjusted by active household and then weighted by income quintile class;
- Livestock and NEWP activities are proportionally adjusted by active household (own cattle, goat or collect fuelwood from rangeland) but labour is not adjusted as herding activities are argued to still involve similar labour demands and reductions in rangeland are likely to lead to more not less NEWP gathering effort. Active households are adjusted for each quintile;
- EWP are adjusted proportionally across all households.

⁷ Other activities (orchard, beer-brewing, brick-making etc) were considered but a limited and unrepresentative distribution across the study area limited their wider applicability.

⁸ Households with IWS (private taps in the home compound) are identified as the appropriate unit for analysis. The estimated area (0.04 ha) conforms to Perez de Mendiguren and Mabelane (2001) though is smaller than the 0.4 ha (4,000m²) used by High and Shackleton (2000). Kitchen garden water use is considered a proxy for irrigated agriculture and supplementary water application (assumed to be from IWS) is estimated from the difference between rainfall (859 mm) and irrigated agriculture evaporation (1149 mm).

⁹ The sustainability (ecological carrying capacity, resource availability, vegetation cover change, etc.) of both the current activities and scenarios is beyond the scope of this study.

It is assumed that a household with an IWS receives 25 lcd and that all other households collecting water from non-IWS sources (specifically, river water) consume 14 lcd¹⁰. Increases in commercial forestry and commercial agriculture land use are assumed to result in a loss of rangeland and that the “water use” of these land uses is calculated relative to the baseline condition (i.e. baseline water use = 0%). This is consistent with the definition of SFRAs in the South African NWA. Whilst there is uncertainty that water is always available in the reticulated system (IWS source), community research indicates that most households are likely to pursue a kitchen garden (food security) strategy given no supply constraints. The benefits of such a strategy provide insights into a potential poverty-focussed compensatory mechanism across the catchment with the relative benefits/costs calculated in water use (m³/pa), economic units (US\$) and food security (maize yield).

Results and discussion

A summary of household water-dependent activities is presented in Table 4. These data present the incremental difference in water use relative to the natural rangeland state. The four activities highlighted exhibit a hydrological impact that is moderated by asset ownership (IWS, cattle, dryland, etc). It is estimated that the bottom quintile depends on non-IWS sources for 68% of its domestic water requirements compared to 40% for the top quintile. Further the inequitable distribution of water provision disproportionately burdens the income poor both in the highest opportunity costs of sourcing non-IWS water and the associated health risks of consuming it. Analyses of daily flow volumes generated by the ACRU model shows that without significant storage associated with IWS systems, the runoff generated for Scenarios 2 and 3 will not be able to supply BHNR (25 lcd) in October at least once every ten years. Kitchen garden farming, which is reliant on IWS for winter irrigation, highlights the skewed distribution of livelihood opportunities (food security) associated with IWS delivery. Analysis of mean dryland maize production illustrates the vulnerability of the bottom quintile to the FAO grain food security threshold of 0.17 tonnes/capita/pa (

Table 5). ACRU simulations indicate that, due to climatic fluctuations, at least one in five years the maize production of the bottom quintile will fail to meet this threshold Whilst kitchen garden farming could potentially buffer this impact, the poorest are least well-served in this regard.

Related to dependency of the poorest on non-IWS sources is the livestock voluntary water intake (VWI) demand, which is mainly provided by river water. The quantity and quality of river water are connected as non-IWS households must often compete with livestock for low river flow and risk health problems associated with sharing an unhygienic water resource. It is the wealthier households with larger herds that compound the situation for the poor.

Dryland agriculture represents the dominant water impact of livelihood activities, which has the linked effect of reducing evaporation (shorter root crops, limited growing season) and concomitant increase in runoff relative to the natural vegetation. Access to land and a sufficient rainfall regime are critical to this activity. The level of green water use is linearly related to dryland plot size, which in turn determines crop (here, maize) production. Poorer households fail to capture the opportunity (income-generating or expenditure-saving) of dryland farming activities in comparison to wealthier households. The explanatory variables of community-level dryland distribution is beyond the scope of this paper but are broadly identified within the socio-cultural context of a patriarchal Tribal Authority system that is organised on kinship networks across generations and/or the ability to both overcome entry barriers (financial, in-kind, other) to land (size, location, fertility, etc.) and the resources to fund land development (traction, labour, market produce, etc.). Dryland farming generates the highest economic return for all income groups.

A summary of the scenarios analysed, and upper-Mutale catchment green and blue water flows for each scenario are provided in Table 7. The scenarios (Table 1) illustrate the baseline condition under natural vegetation (no human modification), the current land use, and four land use change permutations that highlight the hydrological impact of maximum commercial forest cover and/or increased commercial irrigation. The impact of livelihood activities within the rangeland land use on the flow regime (blue water) is presented as water use relative to the baseline condition for all scenarios.

¹⁰ The 14 lcd estimate was taken from a random sample of households with non-IWS and is supported by more extensive surveys of water use in Africa (Thompson et al. 2001; PDG, 1996).

The scenarios used have been selected in order to isolate one particular land use's hydrological influence within the case study area. The most significant hydrological impact occurs when commercial irrigation is increased to 15% land cover with commercial forestry at its maximum potential land use (Scenario 2). However, commercial forestry's contribution to a reduced catchment outflow is only limited in comparison to commercial irrigation (Scenario 3). The economic value generated from the resource base by each income quintile for each scenario is presented in Scenario analyses of land use change impacts on Thengwe catchment outflow (blue water) and contribution of livelihood activities on rangeland hydrology (million m³ per year)

		Rangeland (natural condition)	Commercial Dryland Agriculture	Forestry	Commercial Irrigation	Livelihood impact as % of rangeland	Catchment Outflow x 10 ⁶ m ³ per year
Baseline	Mm ³ /pa (area %)	200 (100%)	0	0	0	0	85
Current land use	Mm ³ /pa (area %)	118 (59%)	49 (27%)	30 (14%)	0	2%	89
Scenario 1	Mm ³ /pa (area %)	34 (17%)	49 (49%)	119 (56%)	0	1%	84
Scenario 2	Mm ³ /pa (area %)	34 (17%)	22 (12%)	119 (56%)	57 (15%)	1%	53
Scenario 3	Mm ³ /pa (area %)	118 (59%)	22 (12%)	30 (14%)	57 (15%)	2%	59
Scenario 4	Mm ³ /pa (area %)	70 (35%)	49 (27%)	81 (38%)	0	1%	86

Table 8.

Intuitively, the economic impacts on livelihoods reflect the proportional change in land availability. It is land availability *per se* rather than neighbouring land uses which determine the livelihood outcomes from this analysis. The impact that increased commercial irrigation has on hydrological functioning appears to have no direct ramification on livelihood outcomes¹¹. However, in Scenarios 2 and 3 where commercial forestry is at its maximum cover or present level with increased irrigation, non-IWS households will face the risk of river water flow being unable to supply the BHNR. The poorest quintile will face a more critical situation under Scenario 2 as it will record a net loss in the identified livelihood activities with the additional risk of failure of the river to supply the BHNR.

Evidence from this research suggests two policy implications that link land use change, water use and livelihoods. First, two of the scenarios will result in a significant reduction in the level of streamflow (blue water), leading to households without IWS facing a water crisis approximately once in every ten years. However, commercial forestry expansion (to 56%) has significantly less hydrological impact than commercial irrigation (to 15%). Scenario 4 makes this position clear with an increased forestry cover to 38% from 14% having only a limited impact on catchment outflow (reduced by 3 Mm³/year). However, the associated impact of an increase in forest cover (or commercial irrigation) would be to reduce the rangeland area and concomitant values harvested from the resource base by local communities. Under Scenario 4 (35% rangeland), there is a proportional reduction in gross values harvested by livelihood activities but the net effect (counting labour) is felt hardest by the poorest as their asset endowments (IWS, land, livestock) are lower than any other quintile. The poorest are more vulnerable to shocks from land use change due to their higher dependency on the resource base. Combining reported household income with water-dependent livelihood activities (non-market) (

Table 6) reveal a significant dependency on the natural resource base for the bottom quintile (62%) in comparison to the top quintile (17%).

Analysis of water-dependent livelihood activities has indicated both the hydrological, economic and food security significance of dryland agriculture within rangeland. The green water use (evaporation) of dryland crops is the most significant hydrological component of rural livelihoods subject to access to land and a sufficient rainfall regime. The high rainfall in the upper-Mutale catchment provides opportunities to develop income-generating and expenditure-saving food production strategies. Table 3 indicates how access to

¹¹ Forward and backward economic/employment linkages are likely to occur but are not captured here.

the dryland resource base is skewed in favour of wealthier quintiles, which are thus better able to exploit the resource. A more equitable distribution of dryland within communities (under *de facto* Tribal Authority management) could significantly reduce the vulnerability of the poor to food security and, potentially, create surpluses to supplement incomes that are less than US\$1 day. The connection between dryland food production and livelihood security mirrors the forestry situation in that rainfall is captured and converted into biomass, which then has value. Whilst land and water are prerequisites in this configuration, the negotiated process of land acquisition, land management and surplus marketing for dryland produce in communal areas are equally important.

Conclusions

The findings from this study suggest that small-scale dryland food production offers a tenable path to poverty reduction (here, increased food security) given relaxation of the identified constraints. However, upstream land use change that reduces the resource base must be weighed against the significant impacts it has on rural livelihood diversified value production systems. The impact of the two most likely land use changes reveal a significant difference in catchment hydrology that have both legal implications (Reserve) and very real human impacts. Land use change in the upper, water-endowed upper-Mutale catchment has the potential for economic growth (employment, production) but any land use change should have an equitable and compensatory mechanism for the negative impacts (externalities) that it is likely to trigger on downstream communities.

Acknowledgements

This publication is an output from a research project funded by the United Kingdom Department for International Development (DFID) for the benefit of developing countries. The views expressed are not necessarily those of DFID. (R7937-Forestry Research Programme). More information on the Catchment Management and Poverty (CAMP) project can be found at: <http://www.cluwrr.ncl.ac.uk/projects/camp/index.html>

GPW Jewitt is the SAPPI Senior Lecturer in Forest Hydrology at the School of Bioresources Engineering and Environmental Hydrology, University of Natal. The support of SAPPI in this regard is gratefully acknowledged.

References

Acocks, J.P.H. (1988) Veld types of South Africa (3rd edition) *Memoirs of the Botanical Society of South Africa*, **57**: 1-46.

Budlender, D. and Brathaug, A.L. (2002) Calculating the value of unpaid labour: a discussion document. Statistics South Africa: Pretoria.

Crookes, D., Ireland, C., de Wit, M., Geach, B., Kgame, W., and Netshiluvhi, T. (2000) Benefits and costs of natural resource use in Sekhukhuneland, Northern Province: case study evidence from Manganeng and Makua. CSIR Report: ENV-P-I-2000-013. CSIR: Pretoria.

FAO (1986) Water for Animals. Food and Agricultural Organization. Report: AGL/MISC/4/85. FAO: Rome.

FAO (2000) New Dimensions in Water Security – Water, Society and Ecosystem Services in the 21st Century. Land and Water Development Division, Food and Agriculture Organization of the United Nations, Rome.

FAO (2001) Non-Wood Forest Products in Africa: A Regional and National Overview. August, 2001. FAO: Rome.

Falkenmark, M (2003) Water cycle and people: water for feeding humanity. Land Use and Water Resources Research (3) 1. [online] URL: <http://www.luwr.com/issues/vol3.html>.

Guijt, I., Hinchcliffe, F. and Melynk, M. (1995) The Hidden Harvest – The Value of Wild Resources in Agricultural System. Institute for Environment and Development (IIED), London, 24pp.

High, C. and Shackleton, C.M. (2000) The comparative value of wild and domestic plants in home gardens of a South African rural village. *Agroforestry* **48**: 141-156.

- Hope, R.A. and Gowing, J.W. (2003) Managing water to reduce poverty: water and livelihood linkages in a rural South African context. Paper presented at the Alternative Water Forum, University of Bradford Centre for International Development, 1-2 May, 2003. Available at: <http://www.brad.ac.uk/acad/bcid/GTP/altwater.html>
- IFAD (2001) Rural Poverty Report 2001 – The challenge of ending rural poverty. International Fund for Agricultural Development. Oxford University Press: UK. Available at: <http://www.ifad.org/poverty/>
- Jewitt, G.P.W. and Schulze, R.E., (1999). Verification of the ACRU model for forest hydrology applications. *Water SA* 25: 483-489.
- Jewitt, GPW (2002) Can integrated water resources management sustain the provision of ecosystem goods and services? *Physics and Chemistry of the Earth* 27/11-22 pp. 887-895
- May, J. (1999) Poverty and Inequality in South Africa: Meeting the Challenges. Zed Books:UK.
- PDG (1996) Financial Modelling of Rural Water Supply: Synthesis Report. Palmer Development Group. Prepared for DWAF by Eberhard, R. Cape Town:RSA.
- Perez de Mendiguren, J.C. and Mabelane, M. (2001) Economics of productive uses of domestic water in rural areas: a case study from Bushbuckridge, South Africa. AWARD Research Report, Acornhoek, South Africa: 137.
- Rockström, J., L. Gordon, C. Folke, M. Falkenmark, and M. Engwall (1999) Linkages among water vapour flows, food production, and terrestrial ecosystem services. *Conservation Ecology* 3(2): 5. [online] URL: <http://www.consecol.org/vol3/iss2/art5>.
- Schulze, R.E. (1995). Hydrology and Agrohydrology : A Text to Accompany the ACRU 3.00 Agrohydrological Modelling System. Water Research Commission, Pretoria. Technology Transfer Report TT69/95. pp 552.
- Scoones, I., Melynk, M. and Pretty, J. (1992) The Hidden Harvest – Wild Foods and Agricultural Systems. A Literature Review and Annotated Bibliography. IIED, London.
- Shackleton, C.M., Shackleton, S.E. and Cousins, B. (2001) The role of land-based strategies in rural livelihoods: the contribution of arable production, animal husbandry and natural resource harvesting in communal areas of South Africa. *Development Southern Africa*, Vol. 18, No. 5, December 2001.
- Shackleton, C.M., Netshiluvhi, T.R., Shackleton, S.E., Geach, B.S., Ballance, A. and Fairbanks, D.F.K. (1999a) Direct use values of woodland resources from three rural villages. CSIR report: ENV-P-I 98210. CSIR: Pretoria.
- Shackleton, C.M., Shackleton, S.E., Netshiluvhi, T.R., Mathabela, F.R. and Phiri, C. (1999b) The direct use value of goods and services attributed to cattle and goats in the Sand River catchment, Bushbuckridge. CSIR report: ENV-P-C 99003. CSIR: Pretoria.
- StatsSA (1999) Rural survey, 1997. Statistical release: P0360. StatsSA: Pretoria.
- StatsSA (2001) A Survey of Time Use. How South African Women and Men Spend their Time. StatsSA: Pretoria.
- Thompson, J., Porra, I. T., Tumwine, J. K., Mujwahuzi, M. R., Katui-Katua, M., Johnstone, N. and Wood, L. (2001). Drawers of Water II: Domestic Water Use in East Africa. International Institute for Environment and Development (IIED). IIED: London.
- World Bank (2001) World Development Report 2000/2001: Attacking Poverty. World Bank: Washington, D.C.

Figure 1 Location of the Luvuvhu Basin and the upper Mutale quaternary catchment (A92A).




Table 1 Land Use in the Luvuvhu Catchment for current, “natural” and four potential development scenarios.

Land Use	Luvuvhu	A92A	Natural	Scen 1	Scen 2	Scen 3	Scen 4
Commercial forestry	4%	14%	0%	56%	56%	14%	38%
Commercial dryland agriculture	10%	22%	0%	17%	12%	12%	27%
Commercial irrigated agriculture	3%	5%	0%	0%	15%	15%	0%
Rangeland	50%	59%	100%	27%	17%	59%	35%
Conservation areas	30%	0%	0%	0%	0%	0%	0%
Urban areas	3%	0%	0%	0%	0%	0%	0%

Table 2 Comparative development profile of Limpopo Province, South Africa and selected SADC countries

	Access to improved water source	Access to sanitation	Adult literacy rate	GDP per capita (PPP\$) 1999
Limpopo Province	47	58	77	2 473
South Africa	70	46	85	8 318
Botswana	70	55	76	6 032
Kenya	53	77	80	975
Mozambique	32	21	42	797
Tanzania	49	86	73	478

Sources: World Bank (2001), SARPN and StatsSA¹²

Table 3 Household profiles by income quintiles (weighted catchment sample)

	Bottom 20%	40%	60%	80%	Top 20%
--	------------	-----	-----	-----	---------

¹² Southern African Regional Poverty Network (SARPN); RSA national statistics office (StatsSA).

Mean household size	5.3	5.2	5.9	6.2	6.2
Proportion with improved water supply	0.21	0.31	0.21	0.33	0.45
Proportion with pit latrine	0.73	0.69	0.73	0.71	0.81
Proportion cooking daily with fuelwood	0.96	0.93	0.95	0.94	0.70
Mean dryland plot (ha)	0.35	0.47	1.02	0.79	0.77
Mean cattle stock	0.12	0.28	1.75	1.83	2.96
Mean number state pension	0.00	0.05	0.04	0.34	0.34
Mean number of full-time employees	0.06	0.07	0.36	0.94	1.20
Mean HH income (US\$/pa, US\$1=R12)	167	526	778	1357	4158

Note: All economic data is reported in US\$, 2002 (US\$1=12 Rands)

Table 4 Household-level green water (evaporation) incremental impact of livelihood activities in relation to the natural hydrological condition (rangeland) by income quintiles in the Thengwe catchment (m³/pa)

	Bottom 20%	40%	60%	80%	Top 20%
Water collection	31	33	35	39	43
Kitchen gardens (irrigated maize)	46	58	46	60	74
Dryland (rainfed maize)	-214	-287	-622	-482	-470
Livestock (cattle/goats)	2	6	32	32	52

Note: NEWP and EWP are considered hydrologically neutral. Positive green water values are balanced by an equivalent reduction in blue water and vice versa, e.g. -100 m³/pa is a reduction in evaporation in comparison to rangeland with an equivalent +100 increase in runoff (blue water)

Table 5 Household maize production estimates (t/pa)

	Bottom 20%	40%	60%	80%	Top 20%
Mean kitchen garden yield	0.52	0.58	0.54	0.50	0.59
Mean dryland yield	1.38	1.86	4.03	3.12	3.04
Maize yield per capita	0.36	0.47	0.77	0.58	0.59

Table 6 Additional estimated household value from water-dependent livelihood activities (US\$/pa)

	Bottom 20%	40%	60%	80%	Top 20%
Gross value	780	935	1589	1315	1427
Gross costs	506	516	601	548	582
Net value	274	419	988	767	845
Proportion of net value to combined household income	0.62	0.44	0.56	0.36	0.17

Table 7 Scenario analyses of land use change impacts on Thengwe catchment outflow (blue water) and contribution of livelihood activities on rangeland hydrology (million m³ per year)

		Rangeland (natural condition)	Commercial Dryland Agriculture	Forestry	Commercial Irrigation	Livelihood impact as % of rangeland	Catchment Outflow x 10 ⁶ m ³ per year
Baseline	Mm ³ /pa (area %)	200 (100%)	0	0	0	0	85
Current land use	Mm ³ /pa (area %)	118 (59%)	49 (27%)	30 (14%)	0	2%	89

Scenario 1	Mm ³ /pa (area %)	34 (17%)	49 (49%)	119 (56%)	0	1%	84
Scenario 2	Mm ³ /pa (area %)	34 (17%)	22 (12%)	119 (56%)	57 (15%)	1%	53
Scenario 3	Mm ³ /pa (area %)	118 (59%)	22 (12%)	30 (14%)	57 (15%)	2%	59
Scenario 4	Mm ³ /pa (area %)	70 (35%)	49 (27%)	81 (38%)	0	1%	86

Table 8 Economic impact of land use change on livelihoods at household level (US\$; Gross values with net values in brackets)

	59% rangeland Current and scenario 3	35% rangeland Scenario 4	17% rangeland Scenario 1 and 2
Bottom 20%	780 (274)	427 (149)	293 (-19)
40%	935 (419)	503 (221)	361 (57)
60%	1589 (988)	779 (444)	550 (182)
80%	1315 (767)	655 (354)	448 (105)
Top 20%	1427 (845)	715 (400)	501 (150)