



Natural
Resources
Institute

Bulletin 77

CONDUCTING ETHNOBOTANICAL SURVEYS

An example from Ghana on
plants used for the protection
of stored cereals and pulses



CONDUCTING ETHNOBOTANICAL SURVEYS: AN EXAMPLE FROM GHANA ON PLANTS USED FOR THE PROTECTION OF STORED CEREALS AND PULSES

J.R. Cobbinah*, C. Moss, P. Golob and S.R. Belmain

*(Forestry Research Institute, Ghana)

Bulletin 77



Natural
Resources
Institute

The Natural Resources Institute (NRI) of the University of Greenwich is an internationally recognized centre of expertise in research and consultancy in the environment and natural resources sector. The Institute carries out research and development and training to promote efficient management and use of renewable natural resources in support of sustainable livelihoods.

Short extracts of material from this publication may be reproduced in any non-advertising, non-profit-making context provided that the source is acknowledged as follows:

COBBINAH, J.R., MOSS, C., GOLOB, P. and BELMAIN, S.R. (1999) *Conducting Ethnobotanical Surveys: an Example from Ghana on Plants used for the Protection of Stored Cereals and Pulses*. NRI Bulletin 77. Chatham, UK: Natural Resources Institute.

Permission for commercial reproduction should be sought from the Managing Editor, Natural Resources Institute, Central Avenue, Chatham Maritime, Kent ME4 4TB, United Kingdom.

Production of this publication was funded by the United Kingdom's Department for International Development. However the Department for International Development can accept no responsibility for any information provided or views expressed.

Price £5.00

Copies of this book can be obtained by writing to NRI Catalogue Services, CAB International, Wallingford, Oxon OX10 8DE, UK. When ordering, please quote **OB77**.

Natural Resources Institute

ISBN: 0 85954 502-4

ISSN: 0952 8245

Contents

Summaries	Page 1
SUMMARY	1
RESUME	1
RESUMEN	2
Introduction	3
Materials and methods	4
Results	4
Discussion and conclusions	10
Acknowledgements	11
References	12
Appendix	14

Summaries

SUMMARY

A survey was undertaken in the Ashanti Region of Ghana to assess the use of plant materials with insecticidal and repellent properties on local farms. Emphasis was placed on plant materials used to protect stores containing cereals and pulses. A total of 27 plant species were recorded as having protective qualities against storage pests of cereals and pulses. The species *Chromolaena odorata* (L.) [siam weed], *Azadirachta indica* A. Juss [neem] and *Capsicum annum* L. [chilli pepper] were the most commonly used to protect stored food. In addition to plants used to protect stored products, bio-activity against insect pests was reported in approximately 95 plant species. The most frequently mentioned plant species were *Bambusa vulgaris* Schrad. [bamboo], *Citrus* spp. [lime, lemon and orange], *Datura innoxia* Mill., *Manihot esculenta* Crantz [cassava] and *Piper umbellatum* L.

The survey found that 26% of the farmers interviewed used botanicals in some form for stored product protection; however, only a small percentage (7%) relied on them exclusively to protect their harvest from storage pests. Smoking of maize stores was the most common method of control in most districts (28%), with the exception of the major maize-growing districts (Ejura and Mampong), where the majority of farmers used conventional insecticides for the control of stored product pests.

RESUME

Une enquête a été effectuée dans la région d'Ashanti, au Ghana, afin d'évaluer l'utilisation de plantes ayant des propriétés insecticides et insectifuges dans les exploitations agricoles locales. L'accent a été mis sur les plantes utilisées pour protéger les magasins de céréales et de légumineuses. Au total, 27 espèces de plantes ont été enregistrées comme détenant des qualités protectrices contre les ravageurs des céréales et des légumineuses entreposées. Les espèces *Chromolaena odorata* (L.), *Azadirachta indica* A. Juss et *Capsicum annum* L. [piment] étaient les plus fréquemment utilisées pour protéger les aliments emmagasinés. Outre les plantes utilisées pour protéger les produits entreposés, une activité biologique contre les insectes ravageurs était signalée dans 95 espèces végétales environ. Les espèces végétales les plus fréquemment mentionnées étaient *Bambusa vulgaris* Schrad. [bambou], *Citrus* spp. [citron vert, citron et orange], *Datura innoxia* Mill., *Manihot esculenta* Crantz [manioc] et *Piper umbellatum* L.

L'enquête a indiqué que 26% des cultivateurs interrogés utilisaient des plantes d'une façon ou d'une autre pour protéger les produits emmagasinés; néanmoins, un petit pourcentage (7%) seulement d'entre eux dépendait exclusivement de celles-ci pour protéger leur récolte contre les insectes de stockage. La fumigation des magasins de maïs était la méthode la plus fréquente de lutte contre ces insectes dans la plupart des districts (28%), à l'exception des principaux districts de culture du maïs (Ejura et Mampong) dans lesquels la majorité des cultivateurs utilisait des insecticides conventionnels pour lutter contre les ravageurs des produits emmagasinés.

RESUMEN

En la región de Ashanti, Ghana, se llevó a cabo un estudio cuyo objetivo era evaluar el uso en las granjas de la zona de material vegetal con propiedades insecticidas y repelentes, habiéndose prestado particular atención al fitomaterial utilizado para proteger los almacenes de cereales y leguminosas. Se estableció que un total de 27 especies vegetales contaban con cualidades de protección contra las plagas de almacenamiento de cereales y leguminosas, habiendo sido las especies *Chromolaena odorata* (L.) [mala hierba de Siam], *Azadirachta indica* A. Juss [nóem] y *Capsicum annuum* L. [chile] las más comúnmente utilizadas para proteger los alimentos almacenados. Además de las plantas utilizadas para el fin antedicho, se obtuvo información sobre la bioactividad contra las plagas de insectos en un total aproximado de 95 especies vegetales. Las especies más frecuentemente mencionadas fueron: *Bambusa vulgaris* Schrad. [bambú], *Citrus* [lima, limón, naranja], *Datura innoxia* Mill., *Manihot esculenta* Crantz [mandioca] y *Piper umbellatum* L.

Este estudio estableció que el 26% de los agricultores entrevistados utilizaban productos botánicos de algún tipo para proteger los productos almacenados. Valga señalar, sin embargo, que solamente un reducido porcentaje (7%) dependía exclusivamente de dicho material para proteger sus cosechas contra las plagas del almacenamiento. Con la excepción de los principales distritos dedicados al cultivo de maíz (Ejura y Mampong), en donde la mayor parte de los agricultores utilizaban insecticidas tradicionales para el control de las plagas de los productos almacenados, en la mayor parte de los distritos (28%), el método de control más comúnmente utilizado era el ahumado de los almacenes de maíz.

Conducting ethnobotanical surveys: an example from Ghana on plants used for the protection of stored cereals and pulses

INTRODUCTION

Plants are well known for their medicinal and insecticidal properties, and considerable research around the world is conducted to screen plants for new drugs and agrochemicals. Because of the overwhelming diversity of plant species and costs involved in developing new drugs or chemicals, ethnobotanical surveys using indigenous knowledge systems have often been employed to develop new compound leads. Ethnobotanical surveys can, therefore, provide a range of information about how plants are used, where the breadth and scope of information is dependent upon the purpose of the survey and how the survey is conducted.

In 1992, the United Nations Convention on Biological Diversity (CBD) was agreed in Rio de Janeiro and has now been ratified by over 120 countries. The treaty reaffirms the sovereign rights of states over their own biological resources, and it is designed to protect indigenous knowledge systems, such as how plants are used, as a form of intellectual property. The CBD states that benefits from biological resources, including ethnobotanical information, should be shared equitably amongst all those involved. It is, therefore, in the interest of signatories to the CBD to invest in the conservation of their biological diversity and to add value to their biological resources by establishing current knowledge systems by indigenous populations.

There are many ways to conduct an ethnobotanical survey where the information required will ultimately be reflected in the methods of survey. However, conducting an ethnobotanical survey should be relatively straightforward. The objective of this Bulletin is to provide an example of how such a survey can be implemented and interpreted on a small scale with very little financial cost. Although the data in this Bulletin relate to the insecticidal properties of plants, and more specifically to stored-product insect pests of grain and legumes in one region of Ghana, the general survey techniques could apply to other ethnobotanical survey objectives, such as finding plants with specific medicinal properties. Although not written as a 'how to' self-help manual, it is to be hoped that the information presented herein will help others interested in the potential benefits of plants to develop their own ethnobotanical survey methodologies.

Historically, botanicals traditionally have been used for pest control in Ghana for generations. They are normally gathered locally and can provide an inexpensive method of control. There is little detailed information regard-

ing the use of local plants with insecticidal properties. However, laboratory studies have confirmed the insecticidal activity of a small number of Ghanaian plant materials traditionally used as protectants (Cobbinah and Osei-Wusu, 1988; Cobbinah and Appiah-Kwarteng, 1989, 1991; Cobbinah and Tuani, 1992; Niber, 1994; Tuani *et al.*, 1994).

The Ashanti Region of southern Ghana has a sub-tropical climate and is characterized by moist/dry semi-deciduous forest or savannah-fringe vegetation. The major crops cultivated in this region include maize (*Zea mays* L.), cassava (*Manihot esculenta* Crantz), groundnuts (*Arachis hypogaea* L.), cowpeas (*Vigna unguiculata* (L.) Walp.), rice (*Oryza sativa* L.), and vegetables (plantain, yam, pepper, garden eggs and tomatoes, etc.).

The main objective of this survey was to gather information on plant species used as protectants, with particular reference to those used against storage pests of cereals and pulses. Information regarding traditional storage methods and the use of conventional insecticides in this region was also assessed.

MATERIALS AND METHODS

A survey was undertaken of all the major farming districts in the Ashanti region of Ghana. Twelve out of 14 administrative districts were surveyed during the period 15 March–15 October, 1995. The two districts not visited were Kumasi, the regional capital, and Obuasi, a gold-mining centre, both of which are highly urbanized and have little agricultural land. Including other major farming areas, a total of 15 farming districts were visited by a two-man survey team, visiting an average of 12 towns and villages within each district (see Figure 1).

Contact was initially made with the chief, chief farmer, opinion leader or the assembly man for each town or village. Their assistance was requested as the objectives of the survey were explained. In the majority of towns and villages a discussion meeting was called of elders and farmers. Those individuals identified in the open forum as having relevant information were asked to complete the questionnaire with the assistance of the survey team (see Appendix). Information regarding cultivated crops and storage methods was obtained. Botanical protectants used by farmers were identified (by local and/or Latin names), as were the application methodology and perceived efficacy. Farmers were divided into two categories based on their level of literacy; literate individuals were defined as those who had been educated past primary school level.

RESULTS

Five hundred and twenty farmers were interviewed from 179 towns and villages in the 15 farming districts surveyed. The majority of farmers who took part were defined as illiterate (mean \pm sem, $64 \pm 3.3\%$). The age and sex of farmers interviewed were biased towards old males. Male respondents comprised 86% of the total number of farmers interviewed, and the mean (\pm sem) age of all respondents was 60 (± 3.1) years old. However, age range and sex ratio of respondents did not vary among districts (Kruskal-Wallis, $n = 15$, $P > 0.05$).




Figure 1 Map of the Ashanti region in Ghana showing the district capitals and other major farming centres that were covered by the survey. The numbers in brackets are the number of towns and villages visited in each farming area.

Plant species exhibiting insecticidal activity against storage pests

Twenty-seven plant species were recorded as being used by farmers in stores to protect cereals and pulses against storage pests (see Table 1). The most commonly mentioned of these species were *Chromolaena odorata* (L.) [siam weed], *Azadirachta indica* A. Juss [neem] and *Capsicum annum* L. [chilli pepper].

Chromolaena odorata was the most widely used of these plants (mentioned in 10 districts by 19% of respondents); however, there was a mixed reaction among farmers to its efficacy as a protectant. *C. odorata* was recognized mainly for its medicinal qualities, and it is used to dress wounds. Leaves of *C. odorata* have been reported to preserve the dead for up to two days. Its use as a protectant in maize (*Zea mays* L.) stores was considered a relatively new innovation, occurring during the last 10 years. It was more popular amongst the younger, more literate generation of farmers who were well informed about present-day storage methods. Harvested maize was generally stored in a barn made of bamboo or wood. A layer of freshly cut *C. odorata* was first spread on the floor, then alternate layers of maize cobs and *C. odorata* were added. Protection has been reported to last for six months using this treatment. *C. odorata* was also used to protect stored cowpea (*Vigna unguiculata* (L.) Walp.). The leaves were dried in the sun and then ground to form a powder. The powder was mixed with loose cowpeas before they were placed in storage sacks. Using this procedure, the cowpeas were reported to be free of insect infestation for four months.

Table 1 Plant species used as protectants of stored grains in the Ashanti region of Ghana

Plant species	Plant part or extract employed	Grain protected	Effects	Number of districts reported
<i>Chromolaena odorata</i>	leaves, leaf powder	maize, cowpeas	repellent	10
<i>Azadirachta indica</i>	leaves, leaf extract, seeds, seed extract	maize, cowpeas	repellent, toxic, oviposition inhibition	7
<i>Capsicum annum</i>	dried fruits, powdered fruits	maize, cowpeas	repellent, oviposition inhibition	4
<i>Citrus aurantiifolia</i>	ripe and unripe whole fruit, juice	maize, cowpeas	repellent	4
<i>Piper umbellatum</i>	leaves, leaf powder	maize	repellent, toxic	3
<i>Arachis hypogaea</i>	oil from seeds	cowpeas	oviposition repellent, ovicidal	2
<i>Cedrela odorata</i>	leaves	maize	repellent	2
<i>Datura innoxia</i>	leaves, leaf extract	maize, cowpeas	toxic	2
<i>Nicotiana tobacum</i>	leaf extract, leaf powder	maize	toxic	2
<i>Thevetia peruviana</i>	paste from fresh roots, leaves, leaf extract	maize	toxic, repellent	2
<i>Citrus sinensis</i>	dried peel	maize, cowpeas	repellent, adult mortality	1
<i>Cleistopholis patens</i>	bark powder	maize	toxic, repellent	1
<i>Cocos nucifera</i>	oil (seed kernel)	cowpeas	oviposition repellent, ovicidal	1
<i>Crossopterix febrifuga</i>	bark	maize	repellent	1
<i>Elaeis guineensis</i>	oil (seed)	cowpeas	oviposition repellent, ovicidal	1
<i>Ficus exasperata</i>	wood ash	maize	toxic, repellent	1
<i>Griffonia simplicifolia</i>	leaves	maize	repellent	1
<i>Khaya senegalensis</i>	leaf powder	cowpeas	oviposition inhibition repellent	1
<i>Mormodica charanta</i>	leaves, leaf extract (combined with lime juice)	maize		1
<i>Parkia clappertoniana</i>	pod extract	maize, millet, cowpeas	repellent	1
<i>Plumbago zeylanica</i>	leaves and leaf powder	rice	toxic	1
<i>Vitellaria paradoxa</i>	residue	cowpeas	repellent	1
Gyakporkporki (gruma)	leaves	maize	toxic	1
Kasali (daghani)	bark exudate	maize	toxic	1
Krubutu (banda)	bark powder	maize	toxic	1
Te (banda)	bark powder	maize	toxic	1
Vinlee (busanga)	leaves	maize, millet	repellent	1

The insecticidal properties of the two other commonly used species, *A. indica* and *Capsicum annum*, were consistently reported throughout the survey (mentioned in seven and four districts, respectively, by 11% and 8% of respondents, respectively). Therefore, both species were used less frequently than *Chromolaena odorata*. *A. indica* was more often recognized, like *C. odorata*, for its medicinal properties rather than its insecticidal properties. There were two methods of application for *A. indica* and *Capsicum annum*; the first was similar to the way *Chromolaena odorata* was used in that leaves or small branches were layered between maize cobs in stores. The sheaths were removed from the cobs to obtain the best results. The second method involved soaking ground, fresh neem leaves in a bucket of water for a day. The mixture was sieved and the extract sprayed on the maize before storage. Maize stored using this second method was reported to stay free of infestations for approximately one year. *C. annum* was used to protect stores of

both maize and cowpeas (*Vigna unguiculata* (L.) Walp.). Shelled maize was bagged and large amounts of dried chilli peppers were placed at the opening before sealing. Dried chillies were ground and moderate amounts of the powder were mixed with the cowpeas for protection. The cowpeas were then bagged and sealed; the chilli powder was reported to be effective at controlling infestation for well over a year.

Control methods and the effects of literacy

Amongst all farmers interviewed, the use of botanicals as the only control measure was relatively low (7.3%) (see Table 2). Literacy/illiteracy had no significant effect upon the percentage of farmers exclusively using botanicals (Binomial test, $P > 0.05$). However, the total use of botanicals (used alone and in combination with other control treatments) by all farmers (26.1%) showed that literacy increased the usage of plant materials against stored product pests (Binomial test, $P < 0.05$).

Table 2 Control methods used among literate and illiterate farmers in the Ashanti region of Ghana

Control method	Percentage of literate farmers	Percentage of illiterate farmers	Total percentage of farmers
Botanicals only	6.9	7.6	7.3
Chemical insecticides only	21.3	6.6	12.25
Smoke only	18.1	34.8	28.4
Botanicals and insecticides	8.0	4.0	5.5
Botanicals and smoke	11.2	9.3	10.0
Insecticides and smoke	12.2	12.3	12.25
Botanicals, insecticides and smoke	6.4	1.3	3.3
Other methods (e.g. ash)	16.0	24.2	21.0
Total botanical usage	32.5	22.2	26.1

Other methods of pest control used in the Ashanti region included conventional insecticides, smoke and ash. Smoke was the most popular of all control measures employed in the region (28.4%) (see Table 2). Generally, maize cobs were hung on cross-bars in the kitchen to allow smoke to pass over them. Alternatively, they were stored in raised barns under which a fire could be lit periodically. The practice of adding wood ash to maize stores was also reported. Ash was spread in alternate layers with maize cobs to provide protection against storage pests.

Between the two groups of farmers, literate and illiterate, the highest users of smoke were illiterate farmers (34.8%), and the use of insecticides was highest among literate farmers (21.3%) (see Table 2). In general, the literate farmers lived in Ghana's largest maize-growing districts (Ejura and Mampong) where farmers cultivate larger areas of land and store for longer periods (up to one year).

Plants exhibiting bioactivity against pests other than those of stored products

With regard to all plants mentioned in the survey, it was found that different parts of plants could be utilized: leaves, bark, fruit, seeds, roots and, in some cases, extracts or powdered material. For any given plant species, the plant parts utilized and the method of preparation of plant material varied amongst villages. There was a high usage of botanicals in districts where $90 \pm 4.2\%$ (mean \pm sem) of respondents mentioned more than three uses of botanicals

against pests. Approximately 95 plant species were cited during the survey as being used to control a variety of insect pests (see Tables 3 and 4). Eleven of these plants were previously mentioned as having activity against storage pests of cereals and pulses (see Table 1). For other pest problems, the most frequently mentioned species (30+ reports) were *Bambusa vulgaris* (bamboo), *Citrus* spp. (lime, lemon and orange), *Datura innoxia* Mill. (pepiewuo), *Manihot esculenta* Crantz (cassava) and *Piper umbellatum* (amumuaha). Plants listed in Table 4 are by local name only as reliable botanic names could not be found. Therefore, data in Table 4 may show different local names that refer to the same plant species, or different plant species referred to by the same local name.

Table 3 Plant species reported to have pesticidal properties in the Ashanti region of Ghana

Scientific name	Local name	Plant part or extract employed	Organism(s) affected
<i>Aframomum</i> spp.	Atiagya	leaves	insect repellent
<i>Albizia ferniginea</i>	Albizia	leaf extract	kills ticks
<i>Albuliton mauritianum</i>	Nwaha	leaves	repel lice on fowls
<i>Alcohoea cordifolia</i>	Gyamma	leaves	control lice on fowls
<i>Ananas comosus</i>	Pineapple	fresh peels and leaf extract	attract mosquitoes, control earth-borne insects
<i>Anchomanes deformis</i>	Awope/ope	tuber extract	kills lice on fowls
<i>Antrocaryon micraster</i>	Apokuma	leaves	control lice on fowls
<i>Aspilia africana</i>	Mfofoo	leaves	repel mosquitoes
<i>Bambusa vulgaris</i>	Bamboo	leaves	repel lice on fowls
<i>Bathia nitida</i>	Odwen	leaves	control insects
<i>Blighia sapida</i>	Akyee	leaves	control mosquitoes
<i>Borassus aethiopum</i>	Fan palm	fruit extract	controls lice on fowls
<i>Capsicum annum</i>	Chilli pepper	fruit powder (in soap solution)	controls field insects
<i>Carica papaya</i>	Pawpaw	leaves and root extract	kill ticks
<i>Cassia alata</i>	Nsempiidua	leaves	control insects
<i>Cassia occidentalis</i>	Nkodaabodee	fruit and leaves	mosquito repellent
<i>Celtis mildbraedii</i>	Esa	ash	insecticide
<i>Citrus aurantiifolia</i>	Lime	leaves (burnt)	mosquito repellent
<i>Citrus limon</i>	Lemon	fruit juice	insecticide
<i>Citrus sinensis</i>	Sweet orange	peel (burnt)	mosquito repellent
<i>Coix lacryma-jobi</i>	Job's tears	seeds soaked in oil (burnt)	mosquito repellent
<i>Cola gigantea</i> var. <i>glabrenscens</i>	Apupua	leaves	control ticks
<i>Costus</i> spp.	Kwekuotoma	leaves	repel honey bees
<i>Culcasia angolensis</i>	Konkrahae	leaves	kill bed bugs
<i>Dalbergia heudelotii</i>	Akasie	leaf extract	kills household insects
<i>Daniellia ogea</i>	Gum copal tree	gum (burnt)	mosquito repellent
<i>Daniellia oliveri</i>	Sanya	gum (burnt)	mosquito repellent
<i>Datura innoxia</i>	Pepiewuo	leaf extract (with clay)	controls earth-borne insects
<i>Datura metel</i>	Kwaseadua	leaf extract (with clay)	controls earth-borne insects
<i>Dombeya</i> spp.	Mfo	leaves	repel lice on fowls
<i>Drypetes floribunda</i>	Dwon	leaf extract	controls ticks
<i>Elaeis guineensis</i>	Oil palm	kernel oil and palm wine	control lice, earth-borne insects
<i>Entada africana</i>	Bonyo/kaboya	leaves	control lice on fowls
<i>Erythrophleum ivorense</i>	Odom	bark extract	insecticide
<i>Ficus exasperata</i>	Fiscus	leaves (burnt)	repel lice on fowls
<i>Griffonia simplicifolia</i>	Kagya	leaves (burnt)	repel lice on fowls
<i>Ipomoea batatas</i>	Sweet potato	leaf extract	controls lice
<i>Jatropha curcas</i>	Physic nut	leaves and ash from leaves	kill domestic insects and bed bugs

Scientific name	Local name	Plant part or extract employed	Organism(s) affected
<i>Justicia flava</i> <i>Lannea prob</i>	Ntumunumu	leaves fruits	mosquito repellent kill and repel lice on fowls
<i>Mangifera indica</i> <i>Manihot esculenta</i>	Mango tree Cassava	bark (burnt) leaves and root tuber (burnt)	mosquito repellent repel honey bees and mosquitoes
<i>Mareya micrantha</i> <i>Mormodia charantia</i>	Nkodaa bodee Nyanya	leaf powder plant and leaf extract	kills crawling insects control insects (especially ticks), and mosquito repellent
<i>Mucuna sloanei</i> <i>Musa paradisiaca</i>	Horse-eye bean Plantain	leaves ash from peel	mosquito repellent, mosquito and termite repellent
<i>Nicotina tobacum</i>	Tobacco	leaves	control domestic insects and termites
<i>Ocimum basilicum</i>	Eme	leaves (burnt)	repel mosquitoes and lice on fowls
<i>Ocimum gratissimum</i> <i>Okoubaka aubrevillei</i> <i>Olyra latifolia</i> <i>Parinari curatellifolia</i> <i>Parkia clappertoniana</i> <i>Piper umbellatum</i> <i>Psydrax parviflora</i> <i>Pycnocoma macrophylla</i>	Tea bush Odee Droben Atena Daudawa Amumuaha Kwawu nsusua Akorie-kofi	leaves seed extract leaves leaves pod extract leaves leaf extract leaf extract	mosquito repellent insecticide repel honey bees control insects controls insects repel lice on fowls repels honey bees toxic to insect pest of domestic animals repels honey bees control lice on fowls controls termites
<i>Solanum aethiopicum</i> <i>Solanum spp.</i> <i>Tectona grandis</i>	Egg plant Nkokodwe Teak	leaf extract leaves leaf extract (mixed with neem extract)	repels honey bees control lice on fowls controls termites
<i>Tetraptera tetraptera</i> <i>Theobroma cocoa</i>	Prekese Cocoa	fruit extract dried pods (burnt)	insecticide mosquito repellent
<i>Tragia spp.</i> <i>Trema orientalis</i> <i>Triplotaxis stellulifera</i>	Sansono Sesea Kooko	leaves leaves stalk and leaves, plants	insect larvae control ticks kill lice on fowls, controls termites
<i>Vernonia conferta</i> <i>Vernonia amygdalina</i> <i>Vitellaria paradoxa</i>	Flakwa Awoweno Shea butter tree	leaves leaf extract leaf powder	insecticide controls ticks kills ticks

Table 4 Local plants reported to have pesticidal properties (local names only) in the Ashanti region of Ghana

Local name	Language	Plant part or extract employed	Organism(s) affected
Afama	Twɩ	leaves	repel lice on fowls
Alata kwadu	Twɩ	fruit extract	kills insects
Aprowah	Twɩ	leaves	control ticks
Bondwaha	Twɩ	root extract	kills insects
Bruma	Kusasi	leaves	control insects
Dumkpaa	Dagbani	leaves (burnt)	repel mosquitoes
Emrodwa	Twɩ	leaves	repel lice on fowls
Epe	Dagomba	rhizome	kills lice on fowls
Godge-te	Ewe	leaves	kill mosquitoes
Kanden	Twɩ	leaves	repel lice on fowls
Kookopaa	Twɩ	seed extract	kills lice on fowls
Kuka/dadee	Twɩ	leaf extract	kills head lice
Mbrodwa	Twɩ	leaves	repel ticks
Mitsimitsi	Ewe	flowers and fruits	trap lice on fowls

Local name	Language	Plant part or extract employed	Organism(s) affected
Mmaa kube	Dagbani	fruit	repel lice on fowls
Nkokotaho	Twi	leaves and seeds (burnt)	control lice on fowls
Nulumnuuya	Dagbani	leaves	control mosquitoes
Nyanyanini	Twi	leaf extract	controls soil-borne insects
Nyutuyowa	Dagbani	leaves	repel mosquitoes
Ponopuw	Dagbani	leaves	repel lice on fowls
Posomono	Ewe	leaves (burnt)	repel mosquitoes
Prison	Kotokori	leaves	control ticks
Samanakatua	Twi	fruit and leaves burnt	repel mosquitoes and lice on fowls
Sampete	Twi	tuber extract	kills termites
Sumenku	Twi	leaves	control insects
Tuankrea	Twi	leaves	control ticks
Twomprowa	Twi	leaves	control ticks

DISCUSSION AND CONCLUSIONS

Knowledge about the insecticidal and repellent properties of plants was found widely among farmers in the Ashanti region of Ghana. However, this knowledge base varied greatly among villages and districts within the region. Although many plant species cited were often endemic across the region, most cases of their use were isolated to one or two districts (see Table 1). The restricted use of commonly occurring plants highlights the often poor communication between villages, perhaps reflecting cultural and ethnic biases that prevent the adoption of new knowledge. Although survey methods suggested that increased education may increase the usage of botanicals in combination with other control strategies, the knowledge base of botanical usage was biased towards old males. This could indicate that traditional information is failing to reach younger farming generations. Technology transfer among villages and villagers could result in greater efficacy of pest control using botanicals.

For most of the plants mentioned in this survey, reliable scientific research and mode-of-action studies on the insecticidal and/or repellent properties are lacking. For example, some laboratory studies showed extracts of *Chromolaena odorata* have no effect on growth and development of *Sitophilus zeamais*, *Callosobruchus maculatus* and *Pseudothermes militaris* (Cobbinah and Appiah-Kwarteng, 1991). However, similar research using extracts and slurries of *Chromolaena odorata* against *Prostephanus truncatus* and *Sitophilus oryzae* showed insecticidal activity at 1% (wt/wt) (Niber, 1994). Evidence is lacking on whether *C. odorata* interferes with host selection and chemo-orientation behaviour of stored-product pests, and it is unknown if volatiles of *C. odorata* are repellent to insect pests. The shoots and foliage of this plant are highly pubescent and physically abrasive. Therefore, if whole plant material were used, it could be possible that these physical characteristics would deter pests. However, *C. odorata* is the most widely distributed weed in Ghana, and its availability, rather than its efficacy, could account for its high usage. Research on the phytochemistry and bioactivity of *C. odorata* (Wollenweber and Roitman, 1996; Wollenweber *et al.*, 1995) and *Chromolaena* spp. (Biller, *et al.*, 1994), showing the presence of novel flavonoids and other secondary metabolites which may be responsible for the insecticidal properties demonstrated (Irobi, 1992; Pancho, 1983), is limited.

It is scientifically recognized that seeds, rather than the other parts, of the neem tree contain the highest proportion of insect antifeedant and growth-regulating factors (Sundaram, 1996; Gahukar, 1995). However, farmers in the survey region were using the leaves of neem trees to protect their produce. Several factors may account for this, such as lack of knowledge, seasonality of seed production, and time and effort required to process the seeds to extract the oil. Education from appropriate sources could increase the efficacy of neem usage. The use of neem against storage pests (Dales, 1996; Xie *et al.*, 1995; Chinwada and Giga, 1993), its chemical constituents (Johnson *et al.*, 1996; Salimuzzaman *et al.*, 1992), insecticidal and growth-regulating properties (Ascher, 1993) and vertebrate toxicity (Oguge *et al.*, 1997; Cohen *et al.*, 1996; Chinnasamy *et al.*, 1993) are well researched. The use of hot peppers against post-harvest pests has also been previously researched (Mbata *et al.*, 1995; Onu and Aliyu, 1995; Lale, 1992).

Traditional methods of stored-product protection were not considered viable alternatives in the large-scale farming of maize that occurred mainly in the districts of Ejura and Mampong. Literate farmers in these districts regarded conventional insecticides as the safest method to ensure the control of post-harvest pests. The most commonly used insecticides by farmers for this purpose were Actellic, Phostoxin, Thiodan, Endosulfan, Gammalin 20 and Cymbush. The usage of botanicals or smoking in large-scale storage is generally not common (Proctor, 1994; Poswal and Akpa, 1991). Botanical usage is dependent upon local availability and collection of plant material; large stores would require very large amounts of plant material. As information and scientific support on botanical protectants is lacking, large-scale farmers also feel more locally traditional methods cannot guarantee the safety of their produce. Although increasing, the amount of research on botanicals used for protection in stored-food systems is still inadequate (Dales, 1996). Therefore, difficulties arise in recommending botanicals as a replacement for chemical insecticides because efficacy levels of botanicals often vary among storage pests, application methods, and stored products. Reliable vertebrate toxicity data and active component/mode-of-action studies are not often available for traditional botanicals, making it impractical to recommend the widespread use of potentially harmful materials.

The survey has found that, while approximately one-quarter of the farmers interviewed used botanicals in some form, only a small percentage relied on them exclusively to protect their harvest from stored-product pests. Literacy did increase the chances that a farmer would consider botanicals as an aspect of a protection regime. Increasing education should, therefore, increase the farmer's ability to choose the types of protection most suitable to his needs. The slow pace of scientific recognition and development of botanicals combined with the increasing availability of conventional pesticides has prevented wide-scale adoption of botanically derived protectants. Increased financial and political support for research on the use of botanicals as alternatives to synthetic pesticides may eventually lead to better stored-product protection at farm level.

ACKNOWLEDGEMENTS

The authors would like to thank the UK Department for International Development (DFID, formerly ODA) for their financing of this project and the Ghanaian Ministry of Agriculture for their support and permission to conduct and publish this research.

REFERENCES

- ASCHER, K.R.S. (1993) Nonconventional insecticidal effects of pesticides available from the neem tree, *Azadirachta indica*. *Archives of Insect Biochemistry and Physiology*, **22**(3-4):433-449.
- BILLER, A., BOPPRE, M., WITTE, L. and HARTMAN, T. (1994) Pyrrolizidine alkaloids in *Chromolaena odorata*, Chemical and chemoeological aspects. *Phytochemistry*, **35**(3): 615-619.
- CHINNASAMY, N., HARISHANKAR, N., KUMAR, P.U. and RUKMINI, C. (1993) Toxicological studies on debitterized neem oil (*Azadirachta indica*). *Food and Chemical Toxicology*, **31**(4): 297-301.
- CHINWADA, P. and GIGA, D. (1993) Vegetable and neem oils as protectants of stored beans against bruchids. Second meeting of the Pan-Africa working group on bean entomology. *CIAT-African Workshop Series*, **25**: 40-49.
- COBBINAH, J. R. and APPIAH-KWARTENG, J. (1989) Effects of various components of neem on stored pests of maize. *Insect Science and its Application*, **10**: 89-92.
- COBBINAH, J. R. and APPIAH-KWARTENG, J. (1991) Pesticidal action of some local plants. *Forig Technical Bulletin*, **10**: 11-18.
- COBBINAH, J. R. and OSEI-WUSU, K. (1988) The effects of neem seed extracts on pests of egg plant, okra and cowpea. *Insect Science and its Application*, **9**: 601-607.
- COBBINAH, J. R. and TUANI, G. K. (1992) Antifeedant effect of *Jatropha curcas* L. seed oil and extracts on the variegated grasshopper, *Zonocerus variegatus* L. (Orthoptera: Acrididae). *Journal of the University of Science and Technology*, **12**: 136-139.
- COHEN, E., QUISTAD, G.B., JEFFERIES, P.R. and CASIDA, J.E. (1996) Nimbolide is the principal cytotoxic component of neem seed insecticide preparation. *Pesticide Science*, **48**(2): 135-140.
- DALES, M.J. (1996) *A Review of Plant Materials used for Controlling Insect Pests of Stored Products*. NRI Bulletin 65. Chatham, UK: Natural Resources Institute.
- GAHUKAR, R.T. (1995) *Neem in plant protection*. Nagpur, India: Agri-horticultural Publishing Ltd.
- IROBI, O.N. (1992) Activities of *Chromolaena odorata* (Compositae) leaf extract against *Pseudomonas aeruginosa* and *Streptococcus faecalis*. *Journal of Ethno-Pharmacology*, **37**(1): 81-82.
- JOHNSON, S., MORGAN, E.D. and PEIRIS, C.N. (1996) Development of the major triterpenoids and oil in the fruit and seeds of neem (*Azadirachta indica*). *Annals of Botany*, **78**(3): 383-388.
- LALE, N.E.S. (1992) Oviposition deterrent and repellent effects of products from dry chilli pepper fruits, *Capsicum* spp., on *Callosobruchus maculatus*. *Postharvest Biology and Technology*, **1**(4): 343-348.
- MBATA, G.N., OJI, O.A. and NWANA, I.E. (1995) Insecticidal action of preparations from the brown pepper, *Piper guineen* Schum, seeds to *Callosobruchus maculatus* (Fabricius). *Discovery and Innovation*, **7**(2): 139-142.

- NIBER, B.T. (1994) The ability of powders and slurries from ten plant species to protect stored grain from attack by *Prostephanus truncatus* Horn (Coleoptera: Bostrichidae) and *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *Journal of Stored Products Research*, **30**(4): 287-301.
- OGUGE, N., NDUNG'U, D. and OKEMO, P. (1997) Effects of neem plant (*Azadirachta indica* Juss, Meliaceae) products on maize grain consumption by three common rodent pests in Kenya. *Belgian Journal of Zoology*, **127**: 129-135.
- ONU, I. and ALIYU, M. (1995) Evaluation of powdered fruits of four peppers (*Capsicum* spp.) for the control of *Callosobruchus maculatus* (F) on stored cowpea seed. *International Journal of Pest Management*, **41**(3): 143-145.
- PANCHO, J.V. (1983) Plants poisonous to livestock in the Philippines (*Chromolaena odorata*, *Lantana camara*, *Pteridium aquilinum*). *Kalikasan, Philippines Journal of Biology*, **12**(3): 193-284.
- POSWAL, M.A.T. and AKPA, A.D. (1991) Current trends in the use of traditional and organic methods for the control of crop pests and diseases in Nigeria. *Tropical Pest Management*, **37**(4): 329-333.
- PROCTOR, D.L. (1994) Grain storage techniques. Evolution and trends in developing countries. *FAO Agricultural Services Bulletin. No. 109*, 277 pp.
- SALIMUZZAMAN, S.; SHAHEEN, F.; SIDDIQUI, B.S.; GHIASUDDIN, S.; SIDDIQUI, S. and FAIZI, S. (1992) Constituents of *Azadirachta indica*: isolation and structure elucidation of a new antibacterial tetranortriterpenoid, mahmoodin, and a new protolimonoid, naheed in. *Journal of Natural Products*, **55**(3): 303-310.
- SUNDARAM, K.M.S. (1996) Azadirachtin biopesticide – a review of studies conducted on its analytical chemistry, environmental behaviour and biological effects. *Journal of Environmental Science and Health Part B – Pesticides*, **31**(4): 913-948.
- TUANI, G.K., COBBINAH, J.R. and AGBODAZE, P.K. (1994) Bioactivity and phytochemical studies on extractives of some Ghanaian plants. *Ghanaian Journal of Forestry*, **1**: 43-47.
- WOLLENWEBER, E., DORR, M. and MUNIAPPAN, R. (1995) Exudate flavonoids in a tropical weed, *Chromolaena odorata* (L.). *Biochemical System Ecology*, **23**(7/8): 873-874.
- WOLLENWEBER, E. and ROITMAN, J.N. (1996) A novel methyl ether of quercetagenin from *Chromolaena odorata* leaf exudate. *Biochemical Systematic Ecology*, **24**(5): 479-480.
- XIE, Y.S., FIELD, P.G. and ISMAN, M.B. (1995) Repellency and toxicity of azadirachtin and neem concentrates to three stored-product beetles. *Journal of Economic Entomology*, **88**(4): 1024-1031.

APPENDIX

Questionnaire Part I

Name: _____ Town/village: _____

Age: _____ House no.: _____

Sex: _____ District: _____

Educational status: _____ Region: _____

Occupation: _____ Date: _____

1. What major crop(s) do you grow?
2. What varieties do you cultivate (if known)?
3. How much of your produce do you store (all, half, third, etc.)?
 - (a) Chemical insecticide (specify)
 - (b) Plant part or extract
 - (c) Any other method (specify)
4. If you ticked 3b (above) mention the name of the plant
5. Where did you obtain it?
6. How did you come by it?
7. How long have you been using it?
8. Did you buy it?
9. What storage methods do you use?
10. What is the length of storage period?
11. What method is used to protect against insect pests?
12. Which part of the plant is used (i.e. stem, bark, leaves, roots, etc.)?
13. How do you process it before use?
14. How do you administer it (i.e. spraying, sprinkling, etc.)?
15. What quantities are used?
16. How effective is it against target insect (i.e. kills, repels, etc.)?
17. How long does it remain effective?

Questionnaire Part II

1. Apart from the above, do you know about any plant(s) which is used in some way to control insects in any situation (i.e. control of mosquitoes, cockroaches, houseflies, termites, etc.)?
2. If yes, give details
 - (a) Name of plant
 - (b) Purpose
 - (c) Source
 - (d) Others
3. Other comments

The Bulletin series presents the results of research and practical scientific work carried out by the Natural Resources Institute. It covers a wide spectrum of topics relevant to development issues ranging from land use assessment, through agricultural production and protection, to storage and processing.

Each bulletin presents a detailed synthesis of the results and conclusions of work carried out within one specialized area, and will be of particular relevance to colleagues within that field and others working on sustainable resource management in developing countries.

Plants are well known for their medicinal and insecticidal properties, and considerable research around the world is conducted to screen plants for new compounds. Ethnobotanical surveys are often used to target

research activity because of the diversity of plant species and costs involved in developing new compounds.

The United Nations Convention on Biological Diversity reaffirms the sovereign rights of States over their own biological resources. Adding value to biological resources by establishing current indigenous knowledge systems could, therefore, lead to financial gains and effective biodiversity conservation. In **Conducting Ethnobotanical Surveys: an example from Ghana on plants used for the protection of stored cereals and pulses** information is presented

to show how ethnobotanical data can be used towards promoting research and conservation of indigenous plant species.

This bulletin will be of special interest to those working in the field of storage entomology, the control of insect pests, the medicinal and insecticidal properties of plant materials and indigenous knowledge system management.

