



## Local Tobacco Control Profiles

### Indicators at a glance (February 2015)

Tobacco summary of indicators - February 2015				
Indicator	Period	Value	Unit	Change from previous
Smoking Prevalence (IHS)	2013	18.4	%	↓
Smoking prevalence - routine & manual	2013	28.6	%	↓
Successful quitters at 4 weeks	2013/14	3524	Crude rate per 100,000	
Successful quitters (CO validated) at 4 weeks	2013/14	2472	Crude rate per 100,000	
Completeness of NS-SEC recording by Stop Smoking Services	2013/14	86.2	%	
Smoking status at time of delivery	2013/14	12.0*	%	↓
Low birth weight of term babies	2012	2.8	%	↔
Smoking prevalence age 15 years - regular smokers	2013	8	%	↔
Smoking prevalence age 15 years - occasional smokers	2013	10	%	↔
Lung cancer registrations	2009 - 11	75.5	DSR per 100,000	↔
Oral cancer registrations	2009 - 11	12.8	DSR per 100,000	↔
Deaths from lung cancer	2011 - 13	60.2	DSR per 100,000	↔
Deaths from chronic obstructive pulmonary disease	2011 - 13	51.5	DSR per 100,000	↑
Smoking attributable mortality	2011 - 13	288.7	DSR per 100,000	↓
Smoking attributable deaths from heart disease	2011 - 13	32.7	DSR per 100,000	↓
Smoking attributable deaths from stroke	2011 - 13	11	DSR per 100,000	↓
Smoking attributable hospital admissions	2012/13	1688	DSR per 100,000	↔
Cost per capita of smoking attributable hospital admissions	2010/11	36.9	Crude rate per 100,000	
Accidental fires ignited by smoking related materials	2012/13	3143	Count	↓
Fatalities from accidental fires ignited by smoking materials	2012/13	73	Count	↓
Smoking prevalence modelled estimates – occasional smokers aged 11-15 years	2009-12	1.4	%	
Smoking prevalence modelled estimates – occasional smokers aged 15 years	2009-12	3.9	%	
Smoking prevalence modelled estimates – occasional smokers aged 16-17 years	2009-12	5.8	%	
Smoking prevalence modelled estimates – regular smokers aged 11-15 years	2009-12	3.1	%	
Smoking prevalence modelled estimates – regular smokers aged 15 years	2009-12	8.7	%	
Smoking prevalence modelled estimates – regular smokers aged 16-17 years	2009-12	14.7	%	
Indicative tobacco sales figures (£ millions)	2013	15446.1	Count	↑

Notes:

- Changes are only shown if they are statistically significant. An upwards arrow represents a significant increase in the indicator value, a downwards arrow represents a significant decrease. A sideways arrow is displayed if there has been no significant change.
- The arrows are coloured green and red for those indicators where a change can be described as improving or worsening respectively