

Protecting and improving the nation's health


Local summary of Legionnaires' disease in residents of England and Wales, 2013

South Midlands and Hertfordshire

This report provides a short summary of Legionella activity in residents of South Midlands and Hertfordshire, with particular emphasis on cases with onset of symptoms during 2013. Data for years 2011 and 2012 and in some tables and graphs, for years 2004 to 2012, are presented for comparative purposes.

For a wider analysis of Legionella activity across England and Wales, please refer to the annual report, Legionnaires' disease in England and Wales; 2013, available on the Public Health England website at; https://www.gov.uk/government/statistics/legionnaires-disease-in-england-and-wales-2013

Figure 1: Cases of legionellosis (confirmed and presumptive) in residents of South Midlands and Hertfordshire by year of onset of symptoms; 2004 – 2013



PHE publications gateway number: 2014392

Published: October 2014

Table 1: Cases of legionellosis (confirmed and presumptive) in residents of South Midlands and Hertfordshire by disease type and year of onset; 2011 – 2013

	Number of confirmed (presumptive*) cases					
	2011	2012	2013	Total (2011 - 2013)		
Legionnaires' disease	6	13	18	37		
Non-pneumonic Legionellosis	-	-	1	1		
Pontiac fever	-	-	-	0		
Total	6	13	19	38		

^{*} Presumptive cases with only a single high titre or PCR result

Table 2: Number & population rate per million of confirmed cases of Legionnaires' disease in residents of South Midlands and Hertfordshire by year of onset; 2011 - 2013

Region	2011		2012		2013		Total (2011 - 2013)	
	n	Rate	n	Rate	n	Rate	n	Rate
South Midlands and Hertfordshire	6	2.96	13	6.42	17	8.39	36	5.93
National	235	4.18	306	5.45	284	5.06	825	4.90

Figure 2: Number of confirmed Legionnaires' disease cases in residents of South Midlands and Hertfordshire by year of onset and category; 2004 – 2013




Table 3: Number and proportion of confirmed Legionnaires' disease cases in residents of South Midlands and Hertfordshire by likely exposure category; 2011 – 2013

Category	2011	2012	2013	Total	National Total (2011 - 2013)
Community [∓]	3 (50.0)	6 (46.2)	12 (70.6)	21 (58.3)	466 (56.5)
Nosocomial	1 (16.7)	1 (7.7)	-	2 (5.6)	28 (3.4)
Travel abroad	2 (33.3)	6 (46.2)	5 (29.4)	13 (36.1)	331 (40.1)
Total	6	13	17	36	825

Ŧ includes travel UK cases

Table 4: Number and proportion of confirmed Legionnaires' disease cases in residents of South Midlands and Hertfordshire by age group and gender, with age specific casefatality rates; 2011 – 2013

Age Groups	Number (%) of cases			No.	Age Specific	
	Female	Male	Total (2011-2013)	Deaths	CFR (95%CI)	
Under 50 years	2 (33.3)	4 (66.7)	6 (16.7)	0	0.0 (0.0 - 45.9 [‡])	
50-59 years	4 (36.4)	7 (63.6)	11 (30.6)	0	0.0 (0.0 - 28.5 [‡])	
60-69 years	6 (42.9)	8 (57.1)	14 (38.9)	3	21.4 (4.7 - 50.38)	
70 years & over	-	5 (100.0)	5 (13.9)	1	20.0 (0.5 - 71.6)	
Total	12 (33.3)	24 (66.7)	36 (100.0)	4	11.1 (3.1 - 26.1)	

≠ one sided confidence interval

Figure 3: Number of confirmed Legionnaires' disease cases in residents of South Midlands and Hertfordshire by month and year of onset; 2011 – 2013




Table 5: Legionnaires' disease cases in residents of South Midlands and Hertfordshire by diagnostic test and year of onset; 2011 - 2013~

	2011	2012	2013	Total (2011 - 2013)	National Total (2011 - 2013)
Case Numbers [†]	6	13	17	36	825
Culture	3 (50.0)	1 (7.7)	6 (35.3)	10 (27.8)	190 (23.0)
Urine	6 (100.0)	13 (100.0)	16 (94.1)	35 (97.2)	810 (98.2)
Serology (SHT)	2 (33.3)	-	-	2 (5.6)	5 (0.6)
Serology (FFR)	-	-	-	0 (0.0)	1 (0.1)
PCR	4 (66.7)	-	8 (47.1)	12 (33.3)	198 (24.0)

[~] This table shows the investigations carried out on each confirmed case

[†] Please note that some cases have more than one diagnostic test carried out, therefore the sum of diagnostic tests does not equal the annual 'case number'.

Table 6: Number of confirmed Legionnaires' disease cases in residents of South Midlands and Hertfordshire by Local Authority (Upper-Tier) and year of onset; 2011 – 2013

Local Authority	2011	2012	2013	Total (2011-2013)
Bedford UA	-	2	-	2
Broxbourne	1	-	-	1
Central Bedfordshire UA	-	-	1	1
Corby	-	1	1	2
Dacorum	-	-	3	3
Daventry	-	-	1	1
East Hertfordshire	1	-	1	2
East Northamptonshire	1	1	2	4
Hertsmere	-	2	-	2
Luton	1	-	-	1
Milton Keynes	-	1	3	4
North Hertfordshire	1	-	-	1
Northampton	-	2	-	2
South Midlands and Hertfordshire	-	-	1	1
St Albans	-	2	1	3
Three Rivers	-	1	1	2
Wellingborough	1	-	1	2
Welwyn Hatfield	-	1	1	2
Total	6	13	17	36

Table 7: Number and type of outbreaks/clusters with associated number of South Midlands and Hertfordshire cases of Legionnaires' disease; 2011 – 2013

Outbreak/Clusters

No outbreaks or clusters were identified with association to South Midlands and Hertfordshire.

First published: October 2014

© Crown copyright 2014

Re-use of Crown copyright material (excluding logos) is allowed under the terms of the Open Government Licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/version/2/ for terms and conditions.