





NHS Cervical Screening Programme Screening Protocol Algorithm for HPV Triage and TOC



July 2014 1 of 5






(i) The management of women with abnormal cytology at this second 12 month follow up test will mirror that at the first 12 month repeat test.

2 of 5 July 2014






3 of 5 July 2014



Cancer Screening Programmes



July 2014 4 of 5



Cancer Screening Programmes

KEY TO COOLS & ABBREVIATIONS

Action codes

routine recall Α Rm early repeat in in months suspend from recal S

PROVISIONAL Result codes

?glandular neoplasia (non cervical) Ø* ?qlandular neoplasia (non cervical) (HP Qest) o G *

1 inadequate

2 negative (not HPV tested) Ν negative (HPV tested)

3 low grade dyskaryosis (not HPV tested) low grade dyskaryosis (HPV tested) M 4 high grade dyskaryosis (severe)

5 high grade dyskaryosis ?invasive squamous carcinoma

6 ?glandular neoplasia of endocervical type

7 high grade dyskaryosis (moderate)

8 borderline change in squamous cells (not HPV tested) В borderline change in squamous cells (HPV tested)

9 borderline change in endocervical cells

Ε borderline change in endocervical cells (HPV tested) Infection codes

Ø (zero) **HPV** negative 9 (nine) **HPV** positive

HPV result inadequate/unreliable U

Miscellaneous

NTDD Next Test Due Date

BLUE indicates codes used on NHAIS in format Cytology result – HPV infection code – Action code Vithdrawn on 06 April 2016 RED indicates manual action required to reset NTDD

July 2014 5 of 5

^{*} non-cervical neoplasia treated as negative for CSP management (2) Used to denote both categories of negative result (negative and ?glandular neoplasia (non cervical)) or both categories of borderline result (borderline change in squamous cells and borderline change in endocervical cells.)