



**Tuberculosis notifications by site of disease (respiratory/non-respiratory), England and Wales,
1913-2012**

Year	Respiratory	Non-respiratory	Total
1913	80,788	36,351	117,139
1914	76,109	23,388	99,497
1915	68,309	22,283	90,592
1916	68,109	22,799	90,908
1917	68,801	20,884	89,685
1918	71,631	18,942	90,573
1919	61,154	16,357	77,511
1920	57,844	15,488	73,332
1921	56,334	15,368	71,702
1922	53,422	15,837	69,259
1923	57,007	18,494	75,501
1924	57,737	18,711	76,448
1925	58,545	19,228	77,773
1926	56,212	18,608	74,820
1927	54,048	17,697	71,745
1928	53,047	17,977	71,024
1929	52,634	16,544	69,178
1930	50,583	16,818	67,401
1931	49,798	16,084	65,882
1932	46,579	15,499	62,078
1933	44,482	14,095	58,577
1934	43,034	13,694	56,728
1935	39,635	12,435	52,070
1936	39,336	12,268	51,604
1937	39,630	12,661	52,291
1938	37,879	12,810	50,689
1939	34,930	11,275	46,205
1940	36,151	10,421	46,572
1941	39,499	11,465	50,964
1942	40,629	11,990	52,619
1943	42,410	11,932	54,342
1944	43,794	10,519	54,313
1945	42,165	9,944	52,109
1946	42,173	9,116	51,289
1947	43,159	8,566	51,725
1948	43,971	8,605	52,576
1949	44,480	7,561	52,041
1950	42,435	6,923	49,358
1951	42,695	6,744	49,439
1952	41,904	6,189	48,093
1953	40,917	5,629	46,546
1954	36,973	5,375	42,348
1955	33,580	4,554	38,134
1956	31,342	4,162	35,504
1957	28,870	3,799	32,669
1958	26,391	3,447	29,838
1959	24,280	2,820	27,100
1960	20,799	2,806	23,605

1961	19,051	2,696	21,747
1962	17,845	2,674	20,519
1963	16,348	2,589	18,937
1964	15,019	2,580	17,599
1965	13,552	2,551	16,103
1966	12,355	2,246	14,601
1967	11,023	2,183	13,206
1968	10,681	2,273	12,954
1969	9,675	2,490	12,165
1970	9,475	2,426	11,901
1971	9,119	2,578	11,697
1972	8,689	2,384	11,073
1973	8,710	2,446	11,156
1974	8,102	2,575	10,677
1975	8,208	2,610	10,818
1976	7,712	2,386	10,098
1977	7,121	2,399	9,520
1978	7,708	2,604	10,312
1979	6,807	2,459	9,266
1980	6,670	2,472	9,142
1981	5,859	2,269	8,128
Year	Pulmonary	Non-pulmonary	Total
1982	5,466	1,944	7,410
1983	4,880	1,923	6,803
1984	4,535	1,606	6,141
1985	4,293	1,564	5,857
1986	4,345	1,648	5,993
1987	3,640	1,446	5,086
1988	3,675	1,489	5,164
1989	3,816	1,616	5,432
1990	3,618	1,586	5,204
1991	3,596	1,840	5,436
1992	3,816	1,983	5,799
1993	3,961	1,960	5,921
1994	3,694	1,897	5,591
1995	3,711	1,897	5,608
1996	3,690	1,964	5,654
1997	3,947	1,912	5,859
1998	3,926	2,161	6,087
1999	3,827	2,317	6,144
2000	4,232	2,340	6,572
2001	4,233	2,481	6,714
2002	4,160	2,593	6,753
2003	3,972	2,546	6,518
2004	3,923	2,800	6,723
2005	4,347	3,281	7,628
2006	4,434	3,187	7,621
2007	3,968	3,021	6,989
2008	4,260	3,059	7,319
2009	4,334	2,906	7,240
2010	4,076	3,724	7,800
2011	4,272	4,128	8,400
2012	4,563	4,096	8,659

Sources: 1913-1982 Statutory Notifications of Infectious Diseases (NOIDs); 2010-2012 Enhanced Tuberculosis Surveillance (ETS)
Prepared by: Centre for Infectious Disease Surveillance and Control, Public Health England

Figure 1: Tuberculosis notifications, England and Wales, 1913-2012




Figure 2: Tuberculosis notifications by site of disease (pulmonary/non-pulmonary), 1982-2012, England and Wales

