



Public Health
England

Spotlight on sexually transmitted infections in South Midlands and Hertfordshire Public Health England Centre

2013 data

About Public Health England

Public Health England exists to protect and improve the nation's health and wellbeing, and reduce health inequalities. It does this through advocacy, partnerships, world-class science, knowledge and intelligence, and the delivery of specialist public health services. PHE is an operationally autonomous executive agency of the Department of Health.

Public Health England

Wellington House

133-155 Waterloo Road

London SE1 8UG

Tel: 020 7654 8000

www.gov.uk/phe

Twitter: @PHE_uk

Facebook: www.facebook.com/PublicHealthEngland

Prepared by: Lynsey Emmett and Richard Puleston, Field Epidemiology Services

For queries relating to this document, please contact: lynsey.emmett@phe.gov.uk

© Crown copyright 2014

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v2.0. To view this licence, visit OGL or email psi@nationalarchives.gsi.gov.uk. Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned. Any enquiries regarding this publication should be sent to lynsey.emmett@phe.gov.uk

Published August 2014

PHE publications gateway number: 2014351



Contents

1. Summary	4
2. Charts, tables, maps	5
3. Information on data sources	11
4. Further information	12
5. Acknowledgements	12

1. Summary

Sexually transmitted infections (STIs) represent an important public health problem in South Midlands and Hertfordshire. Out of all the Public Health England centres it has the seventh highest rate of new STIs in England.

Over 19,800 new STIs were diagnosed in South Midlands and Hertfordshire residents in 2013, representing a rate of 735 diagnoses per 100,000 adults. Rates ranged by local authority from 485 new STI diagnoses per 100,000 population in Central Bedfordshire to 974 new STI diagnoses per 100,000 population in Bedford.

The number of new STIs diagnosed in South Midlands and Hertfordshire residents rose by 7% between 2012 and 2013.

PHE recommends that local areas should be working towards achieving a chlamydia diagnosis rate of at least 2,300 per 100,000 in the 15 to 24 year old age group and this is now an indicator in the Public Health Outcome Framework. In 2013 the chlamydia diagnosis rate for the 15 to 24 year old age group was 2,018 per 100,000 residents.

Men and women have similar rates of new STIs (767 and 681 per 100,000 residents respectively). Where gender and sexual orientation are known, men who have sex with men account for 6% of South Midlands and Hertfordshire residents diagnosed with a new STI in a GUM clinic (76% of those diagnosed with syphilis and 30% of those diagnosed with gonorrhoea).

STIs disproportionately affect young people. South Midlands and Hertfordshire residents aged between 15 and 24 years accounted for 57% of all new STI diagnoses in 2013.

Nationally, black ethnic groups have the highest rates of new STIs, although white ethnic groups contribute the highest numbers.

2. Charts, tables, maps

Figure 1: New STI diagnoses by public health centre (PHEC) of residence: England 2013.

Data source: GUMCAD and CTAD





Figure 2: Diagnoses of the five main STIs: South Midlands and Hertfordshire, 2009-2013

Data source: GUMCAD, CTAD, NCSP and laboratory chlamydia data



Any increase in gonorrhoea diagnoses may be due to the increased use of highly sensitive Nucleic Acid Amplification Tests (NAATs) and additional screening of extra-genital sites in MSM.

Any decrease in genital warts diagnoses may be due to a moderately protective effect of HPV-16/18 vaccination.


Any increase in genital herpes diagnoses may be due to the use of more sensitive NAATs.

Increases or decreases may also reflect testing practices.

Due to changes to the surveillance of chlamydia in 2012, comparisons to previous years are not robust.

Figure 3: Diagnosis rates of the five main STIs: South Midlands and Hertfordshire, 2009-2013.

Data sources: GUMCAD, CTAD, NCSP and laboratory chlamydia data



Any increase in gonorrhoea diagnoses may be due to the increased use of highly sensitive Nucleic Acid Amplification Tests (NAATs) and additional screening of extra-genital sites in MSM.

Any decrease in genital warts diagnoses may be due to a moderately protective effect of HPV-16/18 vaccination.

Any increase in genital herpes diagnoses may be due to the use of more sensitive NAATs.

Increases or decreases may also reflect changing testing practices.

Due to changes to the surveillance of chlamydia in 2012, comparisons to previous years are not robust.

Table 1: Percentage change in New STI diagnoses: South Midlands and Hertfordshire.

Data sources: GUMCAD, CTAD, NCSP and laboratory chlamydia data

Diagnoses	2013	% change 2009-2013	% change 2012-2013
New STIs	19,885	-	7%
Syphilis	73	3%	20%
Gonorrhoea	859	60%	17%
Chlamydia	9,047	-	8%
Genital Herpes	1,426	50%	9%
Genital Warts	3,242	-3%	1%

Figure 4: Rate of new STIs per 100,000 residents by age group in South Midlands and Hertfordshire, 2013.

Data sources: GUMCAD and CTAD




Figure 5a: Ethnicity of South Midlands and Hertfordshire residents diagnosed with a new STI: 2013.

Data sources: GUMCAD and CTAD




Figure 5b: Rates by ethnicity for South Midlands and Hertfordshire residents diagnosed with a new STI: 2013

Data sources: GUMCAD and CTAD





Figure 6: Diagnoses of the five main STIs among MSM in GUM clinics: South Midlands and Hertfordshire, 2009-2013.

Data source: GUMCAD



GUMCAD started in 2009. Reporting of sexual orientation is less likely to be complete for earlier years, so rises seen may be partly artefactual. Any increase in gonorrhoea diagnoses may be due to the increased use of highly sensitive Nucleic Acid Amplification Tests (NAATs) and additional screening of extra-genital sites in MSM. Any decrease in genital warts diagnoses may be due to a moderately protective effect of HPV-16/18 vaccination. Any increase in genital herpes diagnoses may be due to the use of more sensitive NAATs. Any increase or decrease may reflect changes in testing.

Figure 7: Rate of new STI diagnoses per 100,000 population among South Midlands and Hertfordshire residents by upper tier local authority of residence 2013.

Data sources: GUMCAD and CTAD




Figure 8: Rate of chlamydia diagnoses per 100,000 population aged 15-24 years in South Midlands and Hertfordshire by upper tier local authority of residence: 2013.

Data sources: GUMCAD and CTAD




Figure 9: Rate of gonorrhoea diagnoses per 100,000 population in South Midlands and Hertfordshire by upper tier local authority of residence: 2013.

Data source: GUMCAD





Figure 10: Map of new STI rates per 100,000 residents by upper tier local authority in South Midlands and Hertfordshire 2013.

Data source: GUMCAD & CTAD



Contains Ordnance Survey data © Crown copyright and database right 2014.
Contains National Statistics data © Crown copyright and database right 2014.
Map produced using PHEGIS. Contact GIS Team, ER D/MRA, Porton Down,
01980-616937 or gis@phe.gov.uk

3. Information on data sources

Genitourinary Medicine Clinic Activity Dataset (GUMCAD)

This disaggregate reporting system collects information about attendances and diagnoses at genitourinary (GUM) clinics. Patients can be de-duplicated within a clinic but not between clinics. Information about the patient's area of residence is collected along with demographic data and other variables. GUMCAD superseded the earlier KC60 system and can provide data from 2009 onwards. GUMCAD is the main source of data for this report. The data extract used was provided in May 2014.

Chlamydia Testing Activity Dataset (CTAD)

The Chlamydia Testing Activity Dataset (CTAD) is a universal disaggregate dataset for the collection of data on all NHS and LA/NHS-commissioned chlamydia testing carried out in England. The CTAD dataset is comprised of all chlamydia (NAATs) tests for all ages (with the exception of conjunctival samples), from all venues and for all reasons. CTAD enables unified, comprehensive reporting of all chlamydia data, to effectively monitor the impact of the NCSP through estimation of population screening coverage, proportion of all tests that are positive and diagnosis rates.

Date of data extract eg May 2014.

New STIs

New STI diagnoses comprise diagnoses of the following:

Chancroid, LGV, donovanosis, chlamydia, gonorrhoea, genital herpes (first episode), HIV (acute and AIDS defining), *Molluscum contagiosum*, non-specific genital infection (NSGI), non-specific pelvic inflammatory disease (PID) and epididymitis, chlamydial PID and epididymitis (presented in chlamydia total), gonococcal PID & epididymitis (presented in gonorrhoea total), scabies, Pediculosis pubis, syphilis (primary, secondary & early latent), trichomoniasis and genital warts (first episode).

Calculations

Confidence Intervals were calculated using Byar's method.

ONS mid-year estimates for 2012 were used as a denominator for rates for 2013.

4. Further information

Please access the online Sexual and Reproductive Health Profiles for further information: <http://fingertips.phe.org.uk/profile/sexualhealth>

For more information on local sexual health data sources please access the guide: http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1317141016664

Local authorities have access to LA STI epidemiology reports (LASERs) and the HIV and STI portal. They should contact lynsey.emmett@phe.gov.uk if they do not have access to this information.

5. Acknowledgements

We would like to thank the following:

- local sexual health clinics for supplying the GUM clinic data
- local laboratories for supplying the CTAD data
- PHE Centre for Infectious Disease Surveillance and Control (CIDSC) HIV and STI surveillance teams for collection, analysis and distribution of data