

Vitamin D status of older adults

Older adults are at increased risk of poor vitamin D status due to reduced exposure to sunlight, decreased synthesis in the skin, reduced dietary intake, impaired intestinal absorption, and impaired hydroxylation in the liver and kidneys.^{1,2} A person 70 years of age exposed to the same amount of sunlight as a 20 year old person makes approximately 25% of the vitamin D that the 20-year-old person can make,³ placing them at higher risk of vitamin D insufficiency.

A UK study has shown that only 16% of men and 13% of women ≥ 65 years have serum 25(OH)D levels ≥ 75 nmol/l.⁴ The 2014 National Diet and Nutrition Survey shows that year round in the UK, up to a quarter of people aged ≥ 65 years have a serum 25(OH)D concentration < 25 nmol/l. This figure increases to almost a third during the winter months.⁵

The Department of Health recommends that people aged ≥ 65 years should take a daily supplement containing 400 IU (10 μ g) of vitamin D.⁶ The Reference Nutrient Intake (RNI) for vitamin D for adults ≥ 65 years is also 400 IU.⁷ However, there is emerging evidence that a higher level of daily supplementation of 700-1000 IU would be beneficial for older adults.^{8,9,10}

The impact of poor vitamin D status on the health of older people is well established, and includes an increased risk of falls and fractures¹¹ and future nursing home admissions.¹² In hip fracture patients aged ≥ 65 years, 80% appear to have 25(OH)D levels below 50 nmol/l. Less than 5% have serum levels of 75 nmol/l, considered by some as the level required for optimal fracture reduction.¹ Vitamin D deficiency is also associated with poor muscle strength and weakness,¹¹ and poorer physical performance, as measured by the walking test, chair stands and tandem stand, and with a greater decline in physical performance over time in older men and women.¹³ Finally, lower serum 25(OH)D concentrations in older people may be associated with an increased risk of mortality.¹²

Vitamin D supplementation in vitamin D deficient older people has been shown to improve muscle strength, improve functional ability and lead to fewer falls and fractures.²

The beneficial effects of reducing falls and fractures in older adults by providing supplemental vitamin D appear to be dose-related. In the clinical publications discussed as part of this review, only subjects receiving 700-1000 IU vitamin D daily experienced a significantly reduced risk of falling,⁹ and only those receiving between 482–770 IU each day had a reduced risk of fractures.¹⁴

A study by Zhu *et al.*, 2010 investigated the effects of supplementation with vitamin D on muscle strength and mobility in 302 community-dwelling women aged 70-90 years with vitamin D insufficiency. After daily supplementation of 1000 IU vitamin D, the vitamin D group had significantly higher serum 25(OH)D levels than the control group. The same study found that for subjects in the experimental group with baseline values in the lowest tertile, vitamin D improved muscle strength, and the 'Timed Up and Go Test' (TUAG) by 17.5%.⁸

A review which examined the relationship between vitamin D deficiency, muscle function and falls concluded that vitamin D supplementation in vitamin D deficient older people can improve muscle strength, walking distance and functional ability, and leads to a reduction in falls and nonvertebral fractures.²

A meta-analysis of randomised controlled trials found that a daily dose of 700-1000 IU supplemental vitamin D reduced falls by 19% in those with a mean age of 65 years and over,⁹ and a randomised, multiple-dose study in nursing home residents found that those receiving 800 IU vitamin D for 5

months daily had a 72% lower falls rate than the control group (who received no vitamin D supplementation).¹⁰

In 2009, Bischoff-Ferrari *et al.* performed a meta-analysis on the efficacy of oral vitamin D supplementation in preventing nonvertebral and hip fractures in older adults (≥65 years) and found that 482–770 IU/day of supplemental vitamin D reduced nonvertebral fractures by 20% and hip fractures by 18%.¹⁴

In 2011, EFSA concluded that ‘a cause and effect relationship has been established between the intake of vitamin D and a reduction in the risk of falling.’ The EFSA state that in order to obtain the claimed effect, 800 IU of vitamin D should be consumed daily. The target population is people ≥60 years of age.¹⁵

This is supported by other sources. Venning, 2005 states that supplementation with 800 IU of vitamin D daily is needed to have an effect on falls,¹¹ and Brouwer-Brolsma *et al.*, 2013 concluded that to reach a serum 25(OH)D level of 50 nmol/l, older adults aged ≥65 years are recommended to meet a mean daily vitamin D intake of 20 µg (800 IU).¹

References

- 1 Brouwer-Brolsma EM *et al.* *Osteoporos Int* 2013;24:1567-1577.
- 2 Janssen HCJP *et al.* *Am J Clin Nutr* 2002;75:611-615.
- 3 Holick MF. *Am J Clin Nutr* 2004;80(1):1678S-1688S.
- 4 Hirani V *et al.* *Age Ageing* 2010;39:62-68.
- 5 Public Health England & the Foods Standards Agency, 2014: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/310995/NDNS_Y1_to_4_UK_report.pdf.
- 6 Department of Health, 2012: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213703/dh_132508.pdf.
- 7 Department of Health. Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. HMSO, London 1996 (8th ed), pp80
- 8 Zhu K *et al.* *J Am Geriatr Soc* 2010;58:2063-2068.
- 9 Bischoff-Ferrari HA *et al.* *BMJ* 2009;339:b3692.
- 10 Broe KE *et al.* *J Am Geriatr Soc* 2007;55:234-239.
- 11 Venning G. *BMJ* 2005;330:524-526.
- 12 Visser M *et al.* *Am J Clin Nutr* 2006;84:616-622.
- 13 Wicherts IS *et al.* *J Clin Endocrinol Metab* 2007;92(6):2058-2065.
- 14 Bischoff-Ferrari HA *et al.* *Arch Intern Med* 2009;169(6):551-561.
- 15 EFSA Journal 2011;9(9):2382: <http://www.efsa.europa.eu/en/efsajournal/doc/2382.pdf>.