

# Salmon Stocks and Fisheries in England and Wales in 2014





# **SALMON STOCKS AND FISHERIES IN ENGLAND AND WALES, 2014**

**Preliminary assessment prepared for ICES, March 2015**



**Acknowledgement:**

This report has been compiled jointly by staff from the Cefas Salmon and Freshwater Fisheries Team at Lowestoft and fisheries personnel from the Environment Agency and Natural Resources Wales. The monitoring and assessment of salmon stocks is funded by Defra and Welsh Government. All the participating organisations would like to extend their thanks and recognition to the various operational fisheries staff who have collected and compiled the data for this report. Thanks are also due to: the National River Flow Archive at the Centre for Ecology and Hydrology for providing river flow data; to the Game and Wildlife Conservation Trust for data relating to the River Frome; to the General Secretary of ICES for permission to cite the reports of the ICES Working Group on North Atlantic Salmon; and to Graphic Solutions (E.A) Limited for preparation of the final report.

# **CONTENTS**

<b>FOREWORD</b>	<b>4</b>
<b>HIGHLIGHTS FOR 2014</b>	<b>5</b>
<b>REPORT ON SALMON FISHERIES IN 2014</b>	<b>6</b>
<b>1. Description of Stocks and Fisheries</b>	<b>6</b>
<b>2. Fishery regulation measures</b>	<b>9</b>
<b>3. Fishing effort</b>	<b>13</b>
<b>4. Declared catches</b>	<b>20</b>
<b>5. Catch per unit effort (CPUE)</b>	<b>44</b>
<b>6. Exploitation rates</b>	<b>48</b>
<b>REPORT ON STATUS OF STOCKS IN 2014</b>	<b>52</b>
<b>7. Stock monitoring</b>	<b>52</b>
<b>8. Assessment of stock status</b>	<b>61</b>
<b>9. Factors affecting stocks, fisheries and catches</b>	<b>72</b>
9.1 Management measures	72
9.2 Other factors	72
<b>10. Existing and emerging threats to salmon populations</b>	<b>76</b>
<b>11. References</b>	<b>77</b>
<i>ANNEX 1. NASCO's request for scientific advice from ICES in 2015</i>	<i>78</i>
<i>ANNEX 2. Net Limitation Orders applying to salmon net fisheries in England &amp; Wales</i>	<i>81</i>
<i>ANNEX 3. Byelaws applying to salmon rod fisheries in England and Wales</i>	<i>82</i>

## ***FOREWORD***

Annual reports on the status of salmon stocks and fisheries in England and Wales have been produced since 1997. These reports present a preliminary assessment for the latest year to assist ICES in providing scientific advice to NASCO and to provide early feedback to fishery managers and anglers. The list of questions posed by NASCO to ICES for consideration in 2015 is provided at Annex 1 of this report.

For much of the period, the annual reports were prepared by the Centre for Environment, Fisheries and Aquaculture Science (Cefas) and the Environment Agency. However, from 1 April 2013, the functions of the Environment Agency in Wales transferred to a new body, Natural Resources Wales (Cyfoeth Naturiol Cymru) (NRW). This body is now responsible for salmon management and regulation in Wales. All three organisations have therefore contributed to production of the annual assessment report since 2013.

Until 2013, each annual assessment report was designed to stand alone, to avoid the need to refer to previous reports for background information. However, this resulted in much of the descriptive text being very similar year-on-year. From 2014, therefore, and in the interest of streamlining procedures, the report has been split into two separate documents. A background report provides the regulatory framework and describes the various methods and approaches used in the assessment process; this report is expected to change relatively little year on year. The most recent annual assessment (this report) then provides a relatively short description of developments in the latest year together with updated tables and figures. Both reports are available on line at the same location on the gov.uk website.

It should be noted that the data for the most recent year are provisional and will be updated and confirmed as complete catch data are obtained and records validated. The final confirmed data for the current year will be included in next year's report. The Environment Agency and Natural Resources Wales also publish separate Salmonid and Freshwater Fisheries Statistics reports. These are also available at gov.uk: <https://www.gov.uk/government/collections/salmonid-and-freshwater-fisheries-statistics>.

## **HIGHLIGHTS FOR 2014**

- The provisional declared salmon catch by nets and fixed engines in 2014 (11,391 fish; 43.0 t) was 37% below that taken in 2013 and 31% below the average of the previous five years. Most of this catch (90%) was taken in the north east coast fishery. There has been a marked decline in net catches over the past 15-20 years as a consequence of increased regulatory controls and the phasing out of mixed stock fisheries.
- The provisional declared rod catch in 2014 (9,552 fish) is the lowest in the available time series. Catches of both 1SW salmon (grilse) and MSW salmon were well below the recent 5-year means. However, while MSW salmon remained well within the range of previous catches, the catch of 1SW fish was the lowest in the time series.
- Flows were well below average on many rivers for much of the summer and early autumn and thus conditions for angling were often unfavourable. This is expected to have contributed to the poor rod catches, and particularly those of grilse which return later in the season.
- Since 1993, rod catches include an increasing proportion of fish that have been caught and released. In 2014, it is provisionally estimated that 7,368 salmon (77% of the catch) were released. This is the highest percentage ever recorded. It is estimated that released fish contributed an additional 15 million eggs to the breeding population.
- Since the introduction of the national measures to protect spring salmon in 1999, anglers have been releasing a greater proportion of all fish caught, and of large salmon in particular.
- The majority of adult counts and returning stock estimates in monitored rivers in 2014 were below the average of the previous five years. In some rivers counts were particularly poor and at or among the lowest in the available time series.
- Spawning escapement in 2014 was estimated to be above the conservation limit (CL) in only 12 of the 64 principal salmon rivers in England and Wales, the lowest number since assessments have been undertaken. Catch-based assessment procedures may have tended to underestimate actual spawner numbers in 2014 due to the poor angling conditions. However, the assessment outcome is consistent with the low numbers of returning fish seen on many of the rivers with fish counters.
- Formal compliance assessment in 2014 indicated that only 4 rivers (6%) were classified within the top two categories – i.e. had a greater than 50% probability of achieving the management objective (MO) of exceeding the CL in 4 years out of 5, on average. No rivers were classified as 'not at risk' (> 95% probability of meeting the MO) and only 4 were classified as 'probably not at risk' (50-95% probability of meeting the MO).
- Salmon with swollen and/or bleeding vents (Red Vent Syndrome) continued to be observed in 2014 (RVS was first reported in 2004). Affected fish show a degree of recovery in freshwater and appear to be able to spawn successfully.

# REPORT ON SALMON FISHERIES IN 2014

## 1. DESCRIPTION OF STOCKS AND FISHERIES

There are 49 rivers in England and 31 rivers in Wales that regularly support salmon (Figure 1), although some of the stocks are very small and support minimal catches; of these, 64 rivers have been designated 'principal salmon rivers'. Conservation limits (CLs) and Management Targets (MTs) have been set for the 42 principal salmon rivers in England and 22 in Wales and are used to give annual advice on stock status and to assess the need for management and conservation measures.

Rod fishing for salmon is permitted on all rivers supporting salmon stocks, and net or fixed engine fisheries operate on a proportion of these, usually in the river estuaries. Descriptions of the different salmon fishing methods employed in England and Wales can be found in the background report.

Many of the tables and figures presented in this report summarise statistics for England and Wales at a regional level. Following a reorganisation in 2014, the Environment Agency ceased to operate on a regional basis. However, in the interests of maintaining existing time series, data are still aggregated and reported on a regional basis. The full statistics, reported on a river by river basis, are provided in the catch statistics reports which are published annually by the Environment Agency and Natural Resources Wales. A list of the individual rivers falling within each region is provided in Table 1.

**Table 1. The main salmon rivers in England and Wales aggregated according to their former regional jurisdictions. The table also provides details of those rivers with Salmon Action Plans (SAP) and those designated as Special Areas of Conservation (SAC).**

Country	Region (pre-2014)	Region (pre-2011 where different)	River	SAP for river	SAC designation	Comments	
England	North East		Aln				
			Coquet	Yes			
			Tyne	Yes			
			Wear	Yes			
			Tees	Yes			
			Yorkshire Esk	Yes			
		Anglian				No salmon producing rivers, but has coastal fishery	
	South East	Thames Southern		Thames	Yes		
				Itchen	Yes	Yes	
				Test	Yes		
	South West			Hampshire Avon	Yes	Yes	
				Stour	Yes		
				Piddle	Yes		
				Frome	Yes		
			Axe	Yes			
			Exe	Yes			
			Teign	Yes	Yes		
			Dart	Yes	Yes		
			Avon (Devon)	Yes			
			Erme	Yes	Yes		
			Yealm	Yes	Yes		
			Plym	Yes			
		Tavy	Yes	Yes			
		Tamar	Yes				
		Lynher	Yes				
		Looe					



**Table 1. continued**

		Fowey	Yes	
		Camel	Yes	Yes
		Taw	Yes	Yes
		Torr ridge	Yes	
		Lyn	Yes	
Midlands		Ouse		
		Trent	Yes	
		Severn	Yes	
North West		Mersey		
		Ribble	Yes	
		Wyre	Yes	
		Lune	Yes	
		Kent	Yes	
		Leven	Yes	
		Crake	Yes	
		Duddon	Yes	
		Esk (Cumbria)	Yes	
		Irt	Yes	
		Ehen	Yes	Yes
		Calder	Yes	
		Derwent	Yes	Yes
		Ellen		
		Eden	Yes	Yes
		Esk (Border)	Yes	
Wales	Welsh	Wye	Yes	Yes
		Usk	Yes	Yes
		Taff	Yes	
		Ogmore	Yes	
		Afan	Yes	
		Neath		
		Tawe	Yes	
		Loughor	Yes	
		Gwendraeth Fawr & Fach		
		Tywi	Yes	
		Taf	Yes	
		E & W Cleddau	Yes	
		Nevern	Yes	
		Teifi	Yes	Yes
		Aeron		
		Ystwyth		
		Rheidol	Yes	
		Dyfi	Yes	
		Dysynni	Yes	
		Mawddach	Yes	Yes
		Whion		
		Artro		
		Dwryrd	Yes	
		Glaslyn	Yes	
		Dwyfach & Dwyfawr	Yes	
		Llyfni		
		Gwyrfa		Yes
		Seiont	Yes	
		Ogwen	Yes	
		Conwy	Yes	
		Clwyd	Yes	
		Dee	Yes	Yes

Note: Those rivers designated as SACs have salmon identified as a qualifying species in all or part of the catchment. This confers additional protection for salmon in these rivers and any associated on-line lakes.




Figure 1. Map of England and Wales showing the main salmon rivers and denoting those with Salmon Action Plans (\*) and those designated as Special Areas of Conservation (\$) in which salmon must be maintained or restored to favourable conservation status.

## 2. FISHERY REGULATION MEASURES

Salmon fisheries in England and Wales are primarily regulated by effort controls, which specify the nature of the gear that may be operated, along with where, when and how it may be used. A full description of the gears used and the controls is provided in the background report; summary details of the current Net Limitation Orders (NLOs) and byelaws related to rod fisheries are provided in Annex 2 and Annex 3, respectively. The following tables summarise some of the other current controls:

- Table 2 provides details of the rod bag limits and catch limits on net and fixed engine fisheries that are currently in force.
- Table 3 summarises the progress in phasing out those net fisheries that exploit predominantly mixed-stocks where our capacity to manage individual stocks is compromised. A policy to phase out such fisheries has been in place since 1996.
- Table 4 provides details of other arrangements to reduce netting effort in 2014, principally by agreement to release fish alive or by compensating netsmen not to fish for the periods shown.
- Table 5 provides a summary of the effort restrictions recorded in Table 3 over the available time series, 1993 to the present.

In response to the widespread decline in stocks of early-running multi-sea-winter (MSW) salmon, national measures were introduced in 1999 to reduce the levels of exploitation of this stock component. Most netsmen were banned from fishing for salmon before 1 June, with a small number allowed to continue where netting is predominantly for sea trout, on the basis that any salmon caught are returned alive. The national measures also introduced mandatory catch-and-release of salmon by anglers prior to 16 June and imposed other method restrictions. Following review and consultation, the total package of national spring salmon measures was renewed for a further 10 years in December 2008. A brief evaluation of the effect of these measures is included in Section 4.

**Table 2. Statutory rod bag limits and catch limits on net and fixed engine fisheries in force for salmon in 2014.**

EA Region / NRW	River	Rod fishery bag limits			Other constraints	Net/FE catch limits	
		Salmon Bag Limits – per				Fishery	Measure
		day	week	season			
North East		No bag limits apply				Solway haaf nets	Seasonal catch limits apply
Anglian		No bag limits apply					
South East	Thames	2					
South West	Taw	2	3	10	No fish > 70 cm to be retained after 1 August	Tavy seine nets	Seasonal catch limits apply
	Torridge	2	2	7		Tamar seine nets	
Midlands		No bag limits apply				Severn fixed engines	Seasonal catch limits apply
						Severn lave nets	
						Severn seine nets	
North West	Ribble				2	Voluntary carcass tagging scheme	
	Lune				4		
	Crake				11	Limit applies to catch on whole river by all anglers; mandatory carcass tagging scheme	
	Derwent	2				No female fish to be retained after 30th Sept. Voluntary 1 fish per day and 5 fish per season limit encouraged by Derwent Owners Association	
	Eden	2				No female fish to be retained after 10 Sept. Voluntary 1 fish per day and 6 fish per season limits encouraged by River Eden and District Fisheries Association.	
	Border Esk	2				No female fish to be retained after 10 Sept.	
Wales	Taf	2	5		No fish to be retained after 8 Oct.		
	Tywi	2	5		No fish to be retained after 8 Oct.		
	E&W Cleddau	2	5		No fish to be retained after 8 Oct.		
	Nevern	2	5				
	Teifi	2	5				
	Aeron	2	5				
	Ystwyth	2	5				
	Rheidol	2	5				

**Table 3. Number of licences issued each year in net fisheries subject to phase outs (zero NLOs) and closures, 1992–2014.**

Year	Phase Outs										Closures <sup>(a)</sup>												
	Fishery	Phase out commenced	1993	2012	1996	1997	1997	1997	1997	1997	1997	1998	2000	2002	2003	2004	2004	2004	2005	2005	2014		
1992			142		129	17	2	2	0	2	8	4	1	14 <sup>(b)</sup>	6	14	5	4	4	13	2	0	0
1993			124		93	11	1	3	0	2	8	4	1	14 <sup>(b)</sup>	6	14	5	4	4	21	1	0	0
1994			114		72	16	2	2	0	2	8	4	1	14 <sup>(b)</sup>	6	14	5	5	4	18	0	0	0
1995			99		65	9	2	2	0	2	8	4	1	14 <sup>(b)</sup>	6	14	5	5	4	14	0	0	0
1996			89		59	0	2	1	2	2	8	4	1	12	6	14	5	4	14	14	0	0	0
1997			81		56	1	2	2	0	2	8	4	1	14	6	14	5	5	4	15	0	0	0
1998			75		54	<b>0</b>	2	<b>0</b>	<b>0</b>	1	8	4	1	14	6	15	5	5	4	14	0	0	0
1999			72		54		2			1	8	1	1	14	6	14	5	4	4	12	0	0	0
2000			71		46		1			<b>0</b>	<b>0</b> *	1	1	14	6	14	5	4	4	10	0	0	0
2001			70		46		<b>0</b>				1	1	1	14	6	14	5	4	4	8	0	0	0
2002			69		46						1	1	1	3*	6	14	5	4	4	12	0	0	0
2003			16*		45						1	1	1	3	4	14	5	4	4	12	#	0	0
2004			16		40	#	#	#	#	#	#	<b>0</b>	1	3	4	3* <sup>(b)</sup>	1* <sup>(b)</sup>	2* <sup>(b)</sup>	4	11		0	#
2005			16		39					#			1	3	4	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>	4	13			
2006			16		36							1	1	3	3	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>	3*	9*			
2007			16		35							1	1	3	3	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>	2*	4*			
2008			16		33		#	#	#	#		1	1	3	3	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>	<b>0</b> *	3*			
2009			15		30		#	#	#	#		<b>0</b>	3	3	2	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>	<b>0</b> *	<b>0</b> *			
2010			14		30								3	3	2	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>					
2011			14		26								3	3	2	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>					
2012			14	63	25								3 <sup>(c)</sup>	3	2	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>					
2013			13	56	24								3	3	1 <sup>(d)</sup>	3 <sup>(b)</sup>	1 <sup>(b)</sup>	2 <sup>(b)</sup>					
2014			13	52	22								3	3	2	3 <sup>(e)</sup>	0	1 <sup>(e)</sup>					

Note: Bold text denotes target reached.

Key: \* Phase out accelerated by full or partial buy-off.

# Denotes fishery closed by byelaw.

<sup>(a)</sup> Fisheries have not operated for a number of years, now formally closed through byelaw.

<sup>(b)</sup> Licences issued but fishermen compensated not to fish in these years.

<sup>(c)</sup> Phase out replaced by new NLO in 2012 permitting the use of 1 net.

<sup>(d)</sup> Phase out replaced by new NLO in 2013 permitting the use of 2 nets.

<sup>(e)</sup> Phase out remains in place, but under new NLO existing licensees able to resume fishing following 10-year buy-off, subject to catch limits.

**Table 4. Buy off arrangements operating on net fisheries in 2014.**

River/Fishery	Method	Period without netting (full season in parentheses)	Brokers/Funding agency
Fowey	seine nets (all)	complete season (2007 to present) (2 March–31 August)	Brokered by: Environment Agency / South West Water plc
Piddle and Frome (Poole Harbour)	seine net (all)	All salmon & sea trout caught to be released (2008 to present) (1 June–31 July)	Brokered by: Environment Agency and funded by the Frome and Piddle Association <sup>[a]</sup>

Notes: Fowey buy-off – fishing from 2 March to 31 May applies to sea trout only.

<sup>[a]</sup> Local arrangements apply in respect of provision of compensation.

**Table 5. Summary of buy off arrangements and local agreements operating on net fisheries, 1993–2014 (X denotes compensation measure applied; O denotes fishery closed or no licences issued/available).**

Year	Fishery																						
	Itchen seine net #	Avon & Stour seine nets \$	Piddle & Frome seine net \$	Exe seine nets	Teign seine nets	Dart seine nets	Tavy seine nets	Tamar seine nets	Lynher seine nets	Fowey seine nets	Camel drift nets	Taw & Torridge seine nets	Lyn fixed engine	Severn fixed engine	Wye fixed engines	Usk Drift nets	Usk fixed engines	Tywi seine nets	Dee seine nets	Dee trammel nets	Ribble drift nets	Leven lave nets	Cumbrian coastal drift nets
1993	X											X											
1994	X											X											
1995	O											X											
1996	O																						
1997	O	X					X	X	X	X													
1998	O	X		X			X	X	X	X													X
1999	O	X		X			X	X	X	X													X
2000	O	X					X	X	X	X					X	X	X						X
2001	O	X					X	X	X	X					X	O	X						X
2002	O	X					X	X	X	X	X	X			X	O	X					X	X
2003	O	X					X	X	X	X	X		X		X	O	X						X
2004	O	X					X	X	X	X	X		O	X	X	O	X						X
2005	O	X					X	X	X	X	X		O		O	O	O				X		O
2006	O	X			X	X	X	X	X	X	X		O		O	O	O			X	X		O
2007	O	X		X			X	X	X	X	X		O		O	O	O			X	X		O
2008	O	X	X	X			X	X	X	X	X		O		O	O	O		X	X	X		O
2009	O	X	X	X			X	X	X	X	X		O		O	O	O		X	X	O		O
2010	O	X	X	X			X	X	X	X	X		O	X	O	O	O		X	O	O		O
2011	O	X	X	X			X	X	X	X	X		O	X	O	O	O		X	O	O		O
2012	O	O	X				X	X	X	X			O	X	O	O	O		X	O	O		O
2013	O	O	X				X	X	X	X			O		O	O	O			O	O		O
2014	O	O	X						O	X			O		O	O	O			O	O		O

Key: # Fishery operated for scientific purposes - all fish released alive in tracking investigation (no compensation agreement).  
\$ Agreement for all salmon caught to be released alive.

### 3. FISHING EFFORT

The regulatory measures outlined above provide overall limits on the 'allowable' fishing effort in England and Wales; this has fallen in recent years as measures have been introduced to regulate exploitation. The amount that both netsmen and anglers actually fish (the 'utilised' effort) also varies due to weather conditions, perceptions about the numbers of fish returning, and other factors. The following tables and figures summarise changes in allowable and utilised effort:

**Net fisheries** – Table 6 and Figure 2 illustrate the long-term decline in the numbers of licences issued for all types of nets and traps over the period since 1971. The rate of decline in the number of fishing days available, covering a shorter time period, has been greater over this more recent period as a result of additional effort restrictions on remaining licensees (Figure 3). Table 7 provides details of the allowable and utilised effort in salmon net fisheries for the latest season. The percentage of available days that are utilised varies markedly. Figure 3 also illustrates the overall changes in allowable and utilised effort, and the percentage of available days utilised by netsmen, over the available time series. The overall trend in utilised effort has been downward, although there have been some recent marked peaks that are thought to be associated with improved catches in these years.

**Rod fisheries** – Numbers of rod licences (annual and short-term) from 1994 are shown in Table 6 and Figure 4. No comparable data are available for earlier years because of changes in licensing arrangements. Regional summaries of the total rod days fished, over available time series, are provided in Table 8 and Figure 5. It should be noted that effort data (days fished) submitted via rod licence returns do not distinguish between times spent fishing for salmon and sea trout.

#### Overview of fishing effort in 2014

There has been a progressive decline in the number of net and fixed engine licences issued, and hence in fishing effort, over the time series. There was a small increase in the number of licences issued in 2014 compared with 2013 (5 more licences issued), although licence numbers remain around the lowest in the time series. The percentage of the available days utilised by netsmen in the different regions in 2014 (regional averages ranging from 25% to 41%) were broadly similar to 2013. Utilised effort increased in some regions (e.g. South West as a result of the resumption of netting on the Tavy and Tamar), but fell in others (e.g. North East). As in previous years, there was marked variation between the levels of utilised effort in individual fisheries, ranging from 86% (Yorkshire coast drift net) to zero, where licences were available but no fishing for salmon took place. Overall, the percentage of available days utilised by netsmen, declined from a little over 34% in 2000 to about 20% in 2009. However, percentage utilisation has increased again more recently (Figure 3), with some of the highest values (32% and 31% in 2011 and 2013 respectively) associated with relatively good catches. This suggests that the take-up of available fishing opportunities is strongly influenced by catch rates.

The number of salmon rod licences issued over the shorter available time series (1994 on) show variable patterns. The number of short term (one-day and eight-day) rod licences issued has shown a modest decline over the period, from a 5-year mean of about 11,000 licences at the start of the period to a 5-year mean of approximately 9,000 recently. There has been greater variation in the number of annual licences issued; these account for the majority of the salmon caught by anglers. Annual licence numbers decreased sharply from over 26,000 in 1994 to about 15,000 in 2001. This was thought to reflect the decline in salmon stocks and the introduction of restrictions on angling, especially those to protect early-run MSW fish, although licence sales were particularly

low in 2001 due to the restrictions on access to many rivers as a result of an outbreak of the 'foot and mouth' livestock disease. Sales of annual licences increased progressively after this date, reflecting Environment Agency efforts to promote angling and to reduce levels of licence evasion through targeted enforcement efforts. Licence sales in the period 2009 to 2012 have been in excess of 26,000, similar to levels at the start of the time period, although were reduced somewhat in the last two years. In 2014, about 23,800 annual licences were issued; these data are provisional.

The number of days fished by anglers closely followed the reduction in rod licence numbers over the period 1994 to 2001. However, while annual licence sales then recovered to the levels at the start of the time series, the number of days fished by anglers has not. The number of days fished fluctuated somewhat from year to year, but without any obvious trend between 2002 and 2011, but has declined in the most recent years. Provisional estimates of rod days fished in 2014 are the lowest in the time series (Figure 4). There is some variation in the pattern of fishing effort between regions (Figure 5), with the decline in days fished being particularly marked in Wales. Conditions for angling in 2014 appear to have been poor; river flows were generally well below the long-term average for much of the summer (July to October inclusive) (Section 9.2). This coincides with most of the main angling months when peak catches would typically be expected. Provisionally, the overall number of days fished by anglers in 2014 was 33% below the average of the previous five years and the lowest in the available time series.



**Table 6. Numbers of rod licences (1994–2014) and net & fixed engine licences (1971–2014) in England and Wales.**

Year	Rod licences		Gear Type					Total net licences
	Short-term	Annual	Gill	Sweep	Hand-held	FE	Combined drift/T net #	
1971			437	230	294	79	75	1040
1972			308	224	315	76	75	923
1973			291	230	335	70	75	926
1974			280	240	329	69	75	918
1975			269	243	341	69	75	922
1976			275	247	355	70	75	947
1977			273	251	365	71	75	960
1978			249	244	376	70	75	939
1979			241	225	322	68	75	856
1980			233	238	339	69	75	879
1981			232	219	336	72	75	859
1982			232	221	319	72	75	844
1983			232	209	333	73	75	847
1984			226	223	354	74	75	877
1985			223	232	375	69	75	899
1986			220	221	369	64	75	874
1987			213	206	352	68	75	839
1988			210	212	284	70	75	776
1989			208	199	282	75	75	764
1990			207	204	292	70	75	773
1991			199	187	264	66	75	716
1992			203	158	267	65	75	693
1993			187	151	259	55	36	652
1994	10,637	26,641	177	158	257	53	30	645
1995	9,992	24,949	163	156	249	47	29	615
1996	12,508	22,773	151	132	232	42	29	557
1997	11,640	21,146	139	131	231	35	27	536
1998	11,364	21,161	130	129	196	35	26	490
1999	10,709	18,423	120	109	178	30	26	437
2000	10,916	19,223	110	103	158	32	25	403
2001	9,434	14,916	113	99	143	33	24	388
2002	10,039	19,368	113	94	147	32	24	386
2003	8,683	21,253	58	96	160	57	5	371
2004	10,628	22,138	57	75	157	65	5	354
2005	10,170	23,870	59	73	148	65	5	345
2006	9,460	22,146	52	57	147	65	5	321
2007	9,065	23,116	53	45	157	66	5	321
2008	9,761	24,139	55	42	130	66	5	293
2009	9,353	27,108	50	42	118	66	4	276
2010	10,024	26,135	51	41	118	66	4	276
2011	10,121	26,870	53	41	117	66	3	277
2012	9,045	26,090	51	34	115	73	3	273
2013	8,264	25,037	49	29	111	62	3	251
2014	7,574	23,794	48	34	109	65	3	256

Notes: Rod short-term licences are for 1 or 8 days; annual licences are valid from the date of issue to 31 March following.

Gill nets include: drift, trammel, sling and coracle nets.

Sweep nets include: seine (draft and draw) and wade nets.

Hand-held nets include: haaf/heave and lave/dip nets.

Fixed engines include: T-nets, J-nets, stop (compass) nets, putcher ranks, traps, weirs and cribs (coops).

East Anglian coastal nets & Southern seine net are not included, as they are targeted primarily at sea trout and catch few salmon.

Table only includes data for gear licences that are fished (i.e. excluding licences that remain available, but which cannot be fished due to compensation arrangements or other similar provisions).

Data for 2014 are provisional.

Key: # Combined drift/T net licences (issued in Northumbria (Northern area)) have been included in the gill net totals.

**Table 7. Allowable and utilised effort for the principal salmon net fisheries in 2014.**

EA Region / NRW	River/ Fishery <sup>[a]</sup>	Method	No. of licences	NLO <sup>[c]</sup>	Days available <sup>[a,f]</sup>	Allowable effort net days <sup>[h]</sup>	Utilised effort		% days utilised	Av. day/lic.
							net days	net tides		
NE	N Coastal (N)	Drift & T	3	0	113	339				
	N Coastal (N)	Drift	9	0	66	594	587	1007	63	49
	N Coastal (N) <sup>[b]</sup>	T [i]	23	0	113	2,599	663	936	26	29
	N Coastal (S)	Drift	0	0	0	0	0	0	0	0
	N Coastal (S) <sup>[b]</sup>	T [i]	0	0	0	0	0	0	0	0
	Y Coastal	Drift	1	0	66	66	57	70	86	57
	Y Coastal <sup>[b]</sup>	T or J <sup>[i]</sup>	29	0	113	3,277	1,098	1478	34	38
	<b>Region total</b>		<b>65</b>			<b>6,875</b>	<b>2,405</b>	<b>3,491</b>	<b>35</b>	
SW	Avon & Stour	Seine	0	0	0	0	0	0	0	0
	Poole Harbour <sup>[g]</sup>	Seine	1	1	44	44	24	24	55	24
	Exe	Seine	3	3	54	162	131	129	81	44
	Teign <sup>[b]</sup>	Seine	3	3	99	297	84	92	28	28
	Dart <sup>[b]</sup>	Seine	3	3	111	333	162	245	49	54
	Camel	Drift	6	6	21	126	31	34	25	5
	Tavy	Seine <sup>[k]</sup>	1	0	65	65	24	29	37	21
	Tamar	Seine <sup>[k]</sup>	3	0	65	195	110	137	56	33
	Lynher	Seine	0	0	0	0	0	0	0	0
	Fowey <sup>[b,g]</sup>	Seine	0	1	131	131	0	0	0	0
	Taw/Torrige	Seine	3	1	53	159	59	61	37	20
	<b>Region total</b>		<b>23</b>			<b>1,512</b>	<b>625</b>	<b>751</b>	<b>41</b>	
Midlands	Severn	Putchers <sup>[d,k]</sup>	4		76	304	101	160	33	25
	Severn	Seine <sup>[k]</sup>	1	0	66	66	24	24	36	24
	Severn	Lave <sup>[k]</sup>	26	15	66	1,716	409	473	24	16
	<b>Region total</b>		<b>31</b>			<b>2,086</b>	<b>534</b>	<b>657</b>	<b>26</b>	
NW	Ribble	Drift	2	2	66	132	62	66	47	31
	Lune	Haaf	11	12	66	792	451	562	57	41
	Lune	Drift	7	7	66	462	197	206	43	28
	Lune	Seine	0	0	0	0	0	0	0	0
	Kent	Lave	4	6	66	396	74	74	19	19
	Leven	Lave	1	2	44	88	45	56	51	45
	Eden & Esk	Haaf <sup>[k]</sup>	58	105	72	7,560	1366	1493	18	24
	Eden & Esk	Coops <sup>[d]</sup>	3		66	198	198	458	100	66
	<b>Region total</b>		<b>86</b>			<b>9,628</b>	<b>2,393</b>	<b>2,915</b>	<b>25</b>	
Wales	Wye	Lave	8	<sup>[e]</sup>	66	528	177	241	34	22
	Tywi <sup>[b]</sup>	Seine	3	3	109	327	233	291	71	78
	Tywi <sup>[b]</sup>	Coracles	8	8	109	872	396	395	45	50
	Taf	Coracles	1	1	44	44	10	9	23	10
	Taf	Wade	1	1	44	44	7	7	16	7
	E/W Cleddau	Compass	6	6	66	396	72	74	18	12
	Nevern <sup>[b]</sup>	Seine	1	1	109	109	24	24	22	24
	Teifi <sup>[b]</sup>	Seine	3	3	109	327	102	109	31	34
	Teifi <sup>[b]</sup>	Coracles	11	12	109	1,308	460	455	35	42
	Dyfi <sup>[b]</sup>	Seine	2	3	109	327	29	29	9	15
	Dysynni	Seine	1	1	66	66	17	22	26	17
	Mawddach	Seine	2	3	66	198	67	80	34	34
	Conwy	Seine	3	3	66	198	46	56	23	15
	Conwy	Basket <sup>[d]</sup>	0		0	0	0	0	0	0
	Dee	Trammel	0	0	0	0	0	0	0	0
Dee	Seine	0	0	0	0	0	0	0	0	
	<b>Wales total</b>		<b>50</b>			<b>4,744</b>	<b>1,640</b>	<b>1,792</b>	<b>35</b>	

Key: <sup>[a]</sup> National spring salmon byelaws apply – all net fisheries closed until June 1.

<sup>[b]</sup> Sea trout fisheries – exempted from national spring salmon byelaws (all salmon caught before 1 June to be released).

<sup>[c]</sup> NLO refers to number of nets allowed under the terms of the net limitation order for that fishery. Where number of licences exceeds NLO, numbers are being reduced as licensees leave the fishery. For coastal mixed stock fisheries a zero NLO means the fishery is being phased out permanently, but for other fisheries the zero limit may only apply for the duration of the NLO.

<sup>[d]</sup> Denotes fishery operates under an historical certificate of privilege.

<sup>[e]</sup> No NLO, but number of licences capped.

<sup>[f]</sup> In calculating the days available, any day, or part day, on which fishing has been allowed is included. Days available have been adjusted to take account of partial buy-off arrangements and the national measures.

<sup>[g]</sup> Buy-off applies for all or part season (see Table 4 for details).

<sup>[h]</sup> Allowable effort is calculated by multiplying the days available by the number of nets permitted under the NLO, except where the number of licences exceeds the NLO, in which case the higher figure is used.

<sup>[i]</sup> Fisheries subject to phase out, but this subject to review in 2017 to see if some level of sustainable fishing by these nets might be retained.

<sup>[k]</sup> Fishery subject to seasonal catch limit.

**Table 8. Total number of rod days fished, as reported in catch returns, 1994–2014.**

Total days	Former Environment Agency Region						NRW Wales	E&W Total
	NE	Thames	Southern	SW	Midlands	NW		
1994	37,937	343	2,446	41,087	13,596	78,176	118,862	292,447
1995	38,724	414	2,696	35,853	14,893	65,601	85,107	243,288
1996	34,726	154	1,928	32,504	13,056	64,454	84,922	231,744
1997	40,345	181	2,332	38,809	14,886	70,222	102,930	269,705
1998	38,229	145	2,095	31,285	11,493	64,248	85,906	233,401
1999	31,676	311	2,018	25,642	7,024	50,667	70,660	187,998
2000	32,319	143	1,771	22,401	5,373	49,255	66,270	177,532
2001	27,485	111	2,117	18,573	4,084	23,320	59,163	134,853
2002	34,423	91	2,462	25,526	4,720	43,278	72,328	182,828
2003	31,030	126	2,663	23,322	5,302	37,567	72,719	172,729
2004	37,677	110	2,344	24,730	4,633	48,174	72,846	190,514
2005	37,355	86	2,096	22,427	5,221	49,698	69,786	186,669
2006	30,441	21	1,602	17,704	4,124	40,782	53,441	148,115
2007	33,292	64	1,816	19,979	3,800	40,828	64,694	164,473
2008	35,633	53	2,132	20,708	4,211	44,499	63,776	171,012
2009	37,366	46	2,046	22,828	4,819	47,509	69,144	183,758
2010	42,061	37	2,652	23,279	5,052	51,774	70,201	195,056
2011	42,982	22	2,873	24,122	5,105	53,340	68,453	196,897
2012	38,349	13	2,284	20,763	3,521	47,352	63,131	175,413
2013	38,785	17	2,709	18,497	4,211	46,163	56,634	167,016
2014	30,421	46	2,189	13,121	3,482	32,889	41,056	123,204
Mean (2009-13)	39,909	27	2,513	21,898	4,542	49,228	65,513	183,628
% change:								
2014 on 2013	-22	+171	-19	-29	-17	-29	-28	-26
2014 on 5-yr mean	-24	+70	-13	-40	-23	-33	-37	-33

Notes: Includes effort targeted at both salmon and sea trout.

Table does not include rod days fished in the Anglian Region, where there is a very low level of effort.

Data for 2014 are provisional.




Figure 2. Numbers of salmon net & fixed engine licences issued in England and Wales, 1971–2014.




Figure 3. Numbers of fishing days available to net and fixed engine fisheries in England and Wales, and number and percentage of available days utilised, 1999–2014.




Figure 4. Numbers of annual and short-term rod licences issued, and the number of rod days fished in England and Wales, 1994–2014.




Figure 5. Numbers of rod days fished (as reported in catch returns), 1994–2014.

## 4. DECLARED CATCHES

The chief indicators of the state of salmon stocks are the catches taken by rod and net fisheries. It should be remembered that the data presented here for 2014 are provisional. Final confirmed data for 2014 will be reported in the Environment Agency's annual compilation of catch statistics, which will be available later in the year (see, for example, Environment Agency, 2014).

**Net and rod fisheries** – The following tables and figures provide estimates for 2014 together with confirmed catches for earlier years:

- Table 9 provides the total declared number and weight of salmon caught by nets & fixed engines and by rods in England and Wales since 1988, and the total catch and total retained catch (excluding fish that have been caught and released) by all methods combined.
- Table 10 gives a regional breakdown of the provisional 2014 rod and net catches (based on the former Environment Agency regions). These data are total catches and include fish that have been caught and released.
- Table 11 and Figure 6 provide time series of regional net and fixed engine catches from 1971 on.
- Table 12 and Figure 7 provide time series of regional rod catches from 1993 on, distinguishing fish caught and released from those caught and retained (data on catch-and-release were not recorded prior to 1993).

**Catches in coastal, estuary and river fisheries** – ICES requests that catch data (fish caught and retained only) are grouped for coastal, estuary and river fisheries. Data for the available time series, since 1988, are presented in Table 13 and Figure 8. Details of the fisheries included in the various categories are provided in the footnotes to the table. The catch for the coastal zone mainly reflects the catch in the north east drift and fixed net fishery. Only two coastal fisheries remained in operation in 2014 and one of these, Anglian, takes very few salmon (Table 11). The catches in each of the categories have been subjected to downward pressures over recent years, in the case of the coastal and estuarine categories due to the substantial reductions in fishing effort, and, in the case of rod fisheries, due to the increasing use of catch and release.

**Catch and release (C&R)** – C&R data were first collected in England and Wales in 1993, and the practice has been used increasingly by salmon anglers in recent years. This is largely a result of voluntary measures, but also reflects the introduction of mandatory C&R on some rivers (details available in Annex 3). Regional C&R rates are provided in Table 12 and Figure 7 and a summary for England and Wales as a whole is given in Table 14 and Figure 9. C&R rates for each major salmon river in England and Wales are published in the annual catch statistics.

**Long-term catch trends** – The annual declared net and fixed engine catch for England and Wales since 1956 is shown in Figure 10; this distinguishes the catch taken in the north east coast fishery from net catches elsewhere. Figure 11 presents the declared rod catch of salmon from 1956, including (since 1993) fish that have been caught and released. It is unclear to what extent fish may be caught and recorded more than once as a result of C&R.

**Undeclared and illegal catches** – The non-reported and illegal catch for England and Wales in 2014 is estimated at about 10 tonnes, which represents approximately 16% of the total weight (including the unreported and illegal catch) of salmon caught and killed. Of the total unreported and illegal catch in 2014 (approximately 2,400 salmon), 52% by number is estimated to have derived from illegal catches, 39% from under-reporting in rod fisheries and 9% from under-reporting in net fisheries. The methodology used to derive these crude estimates is provided in the background report.

**Effect of the national spring salmon measures** – The restrictions imposed as a result of the national measures, since 1999, have affected both net and rod fisheries. Table 15 and Figures 12a (nets) and 12b (rods) show the reduction in the number of fish caught before June. Table 16 and Figure 13 show the numbers of salmon released by weight category (<3.6 kg (8 lbs), 3.6–6.4 kg (14 lbs), and >6.4 kg) and season, since 1998. This illustrates that anglers have been voluntarily releasing an increased proportion of all fish caught after June, and large salmon in particular.

**Age composition of catches** – ICES conducts annual salmon stock assessments on two separate stock components: those fish that mature after one winter at sea (i.e. one-sea-winter fish / 1SW or grilse) and those that mature after two or more winters at sea (i.e. multi-sea-winter / MSW fish). The relative proportions of the different sea-age groups have shown marked variability over time (Figure 14). It is therefore necessary to be able to estimate the relative proportions of 1SW and MSW fish in catches; details of the approaches used are provided in the background report.

- **Nets** – The relative proportions of 1SW and MSW fish in regional net catches in 2014 are provided in Table 17 and available time series are presented in Figures 15 and 16. The longer time series for the North East Region reflects the consistent reporting arrangements that have applied in this fishery since the mid 1960s.
- **Rods** – The estimated age composition of catches for each of the principal salmon rivers in 2014 is provided in Table 18. Of these, 10 rivers (24%) were estimated to contain 50% or more MSW salmon (including fish subsequently released), 21 rivers (51%) had between 25% and 50% MSW salmon and 10 rivers (24%) less than 25% MSW salmon in the rod catch. Trends in the relative proportions of fish in these different categories are presented in Figure 17.

The estimated numbers of 1SW and MSW salmon (including fish released), and the percentage of MSW fish, in regional rod catches over the period since 1992 are provided in Table 19; these data have been corrected for under-reporting – a scaling factor of 1.1 is currently applied. The number and percentage of MSW salmon in regional rod catches are illustrated in Figure 18. A summary of the estimated rod catch of 1SW and MSW salmon for England and Wales as a whole, for the same period, is provided at Figure 19.

The proportion of MSW salmon recorded in the catch was expected to fall after 1999, following the introduction of the measures to restrict netting effort in the early part of the season when MSW salmon comprise the majority of the catch. The apparent lack of such an effect may reflect poor grilse runs or catches, particularly in recent years. Changes in run timing might also be a factor. There are not thought to have been any substantive changes to fishing gears that might have affected their size selectivity over this period.

## Overview of catches in 2014

The total declared salmon catch for 2014 (including those fish released alive by anglers) is provisionally estimated at 83.2 t, representing 20,943 fish, and comprising 43.0 t (11,391 fish) by nets and fixed engines and 40.2 t (9,552 fish) by rods. Of the rod caught fish, 7,368 were released alive (31.4 t), representing 77% of the reported rod catch by number. In total, just over 13,500 fish (51.5 t) were retained in all fisheries. These figures do not take account of catches of salmon which go unreported (including those taken illegally), and it is estimated that there may have been a total of 10 t of additional fish caught in 2014.

The total declared catch by nets and fixed engines in 2014 was 37% below the catch recorded in 2013, 31% below the average of the previous five years and among the lowest in the time series. There has been a marked decline in net catches over the past 15-20 years as a consequence of increased regulatory controls and the phasing out of some fisheries.

The policy to phase out salmon fisheries predominantly exploiting mixed stocks, where the capacity to manage individual river stocks is compromised, has had a major effect on catches. The largest phase out has occurred in the north east coast fishery. This was enhanced by a partial buy out in 2003, which reduced the number of drift net licences from 69 in 2002 to 16 (an immediate reduction of 77%). The ongoing phase out has resulted in the number of drift net licences continuing to fall; this currently stands at 13. Despite this, the north east coast fishery still accounts for the majority of the England and Wales net catch. In the past five years, the fishery has accounted for between 86% and 94% of the total net catch.

The provisional declared rod catch in 2014 (including released fish) was over 50% down on the average of the previous 5 years. Long-term trends in rod catch (Figure 11) indicate a progressive decline from the peak in the mid-1960's to the early 2000's. This was followed by a general improvement in the rod catch between 2004 and 2011, suggesting some degree of reversal in the declining trend, when catches, including fish caught and released, were typically above the long-term average. However, there has been a progressive decline in catches since 2012 and the provisional rod catch for 2014 is the lowest in the entire time series. Low flows and poor conditions for angling likely contributed to this low catch (Section 9.2). It should also be noted that rod catch trends on individual rivers have varied from much more severe declines to substantial recoveries. The percentage of rod caught fish released by anglers has increased progressively since such data were first recorded in 1993; it is provisionally estimated that 77% of rod caught fish were released in 2014.

Rod catches of 1SW salmon show greater year to year variability than those of MSW fish in numerical terms (Figure 19). Since the early 1990s, catches of 1SW salmon have ranged from a high of over 24,200 to just 5,600. Catches in the period 2004 to 2011 were generally higher than those in the early part of the time series. However, there has been a sharp downturn in the 1SW rod catch in the last three years, with the provisional declared catch in 2014 being the lowest in the time series. Rod catches of MSW salmon have demonstrated somewhat smaller changes (range 3,100 to 10,900). While there has been a decrease in MSW catches in the most recent years, MSW catches appear to be trending positively over the period as a whole. MSW salmon have comprised 45% of the total rod catch, on average, in the last four years, compared with an average of 25% in the preceding period back to 1992.



## **Assessment of national catch trend**

The annual assessment of the status of salmon stocks in the north east Atlantic carried out by the ICES North Atlantic Salmon Working Group, requires the best available time series of nominal catch data (i.e. fish retained) for each country. Figure 20 provides the current best estimate of the total catches of 1SW and MSW salmon in England and Wales, for the period since 1971. These data have been adjusted to take account of non-reported and illegal catches, and exclude the estimated number of Scottish origin fish taken in the north east coast fishery. Further details on the procedures used in deriving these estimates are provided in the background report.

The data indicate that catches of salmon in England and Wales (fish caught and killed only) have declined by over 80% from the early 1970s to the present time. There was a particularly marked decline in catch around 1990, which is consistent with the general perception of a decrease in the marine survival for many stocks around the North Atlantic at about this time. The overall decrease has been greater in the non-maturing (i.e. potential MSW) fish, which have declined by approaching 90% between these periods, whilst the maturing (i.e. potential 1SW) component has declined by about 80%. However, there has been a marked increase in the proportion of MSW salmon in the catch in the last four years (Figure 20).

**Table 9. Declared number and weight of salmon caught by nets & fixed engines and by rods in England and Wales, 1988–2014.**

Year	Nets & Fixed Engines		Rods (inc. released fish)		Total caught		Total retained	
	No.	Wt (t)	No.	Wt (t)	No.	Wt (t)	No.	Wt (t)
1988	77,317	271.1	32,846	123.6	110,163	394.8	110,163	394.8
1989	68,940	239.3	14,728	56.6	83,668	295.9	83,668	295.9
1990	71,827	277.8	14,849	60.3	86,676	338.1	86,676	338.1
1991	37,675	144.6	13,974	55.5	51,649	200.1	51,649	200.1
1992	33,849	130.4	10,737	40.2	44,586	170.5	44,586	170.5
1993	56,566	202.3	14,059	51.1	70,625	253.4	69,177	248.1
1994	66,457	241.9	24,891	94.0	91,348	335.9	88,121	323.7
1995	67,659	245.7	16,008	61.0	83,667	306.7	80,478	294.6
1996	32,680	125.7	17,444	71.5	50,124	197.2	46,696	183.2
1997	31,459	107.2	13,047	48.4	44,506	155.6	41,374	141.8
1998	25,179	84.7	17,109	59.1	42,288	143.9	36,917	122.9
1999	34,167	124.4	12,505	49.8	46,672	174.2	41,107	150.0
2000	50,998	182.7	17,596	67.5	68,594	250.2	60,953	218.8
2001	43,243	153.3	14,383	56.8	57,626	210.1	51,307	184.2
2002	38,279	133.2	15,282	60.4	53,561	193.6	45,669	161.0
2003	17,219	69.2	11,519	48.5	28,738	117.7	22,206	89.0
2004	16,581	59.1	27,332	104.5	43,913	163.6	30,559	111.4
2005	16,811	60.9	21,418	85.8	38,229	146.7	26,162	96.5
2006	13,578	50.5	19,509	72.1	33,087	122.6	22,056	79.8
2007	10,922	37.9	19,984	71.6	30,906	109.5	19,914	67.1
2008	8,647	30.2	23,512	83.7	32,159	113.9	19,036	63.7
2009	7,505	29.3	15,563	62.0	23,068	91.3	13,910	54.0
2010	22,615	72.9	25,153	89.4	47,768	162.3	32,695	108.7
2011	26,193	101.2	23,199	98.5	49,392	199.7	34,575	135.8
2012	8,484	31.0	18,450	81.1	26,934	112.1	14,926	58.0
2013	18,176	67.2	14,920	62.2	33,096	129.4	22,608	84.1
2014	11,391	43.0	9,552	40.2	20,943	83.2	13,502	51.5
Mean (2009-2013)	16,595	60.3	19,457	78.6	36,052	139.0	23,743	88.1

Note: Data for 2014 are provisional.

**Table 10. Provisional regional declared number and weight of salmon caught by nets (retained fish only) and rods (including catch-and-release), 2014.**

Former EA Region / NRW	Net catch		Rod catch		Total catch	
	No.	Weight (kg)	No.	Weight (kg)	No.	Weight (kg)
North East	10,210	38,377	3,957	17,979	14,167	56,356
Anglian	7	19	0	0	7	19
Southern <sup>[a]</sup>	0	0	323	1,107	323	1,107
South West	291	955	729	2,564	1,020	3,519
Midlands	177	816	190	942	367	1,758
North West	479	1,942	2,399	9,371	2,878	11,313
Wales	227	905	1,954	8,261	2,181	9,166
Unknown	0	0	0	0	0	0
<b>E&amp;W Total</b>	<b>11,391</b>	<b>43,014</b>	<b>9,552</b>	<b>40,223</b>	<b>20,943</b>	<b>83,237</b>

Key: <sup>[a]</sup> Rod catches for Southern Region have previously been owners returns, but declared catch reported since 2012.

**Table 11. Declared number of salmon caught by nets and fixed engines, 1971–2014.**

Year	Previous Environment Agency Region						NRW Wales	E&W Total
	NE	Anglian <sup>(a)</sup>	Southern	SW	Midlands	NW		
1971	60,353		186	11,827	3,629	4,989	9,008	89,992
1972	51,681		317	13,146	4,467	3,941	9,633	83,185
1973	62,842		455	12,637	3,887	4,939	9,006	93,766
1974	52,756		346	8,709	3,152	6,282	8,883	80,128
1975	53,451		384	14,736	3,833	5,251	11,107	88,762
1976	15,701		195	11,365	3,194	5,348	7,712	43,515
1977	52,888		212	7,566	2,593	5,312	6,492	75,063
1978	51,630		163	6,653	2,327	7,321	7,426	75,520
1979	43,464		282	7,853	1,404	3,723	4,552	61,278
1980	45,780		137	9,303	3,204	3,769	6,880	69,073
1981	69,113		233	11,391	4,014	5,048	9,050	98,849
1982	50,167		94	6,341	1,738	3,944	4,481	66,765
1983	77,277		163	8,718	2,699	8,489	4,834	102,180
1984	59,295		157	8,489	3,376	7,957	3,947	83,221
1985	57,356		251	9,876	2,423	2,559	3,465	75,930
1986	63,425		461	11,548	3,300	6,682	5,031	90,447
1987	36,143		505	14,530	2,963	5,052	4,535	63,728
1988	50,849		477	11,799	3,511	5,671	5,010	77,317
1989	41,453	4	83	10,684	4,364	7,294	5,058	68,940
1990	51,530	9	43	5,892	4,397	5,579	4,377	71,827
1991	25,429	34	25	2,897	1,747	4,499	3,044	37,675
1992	20,144	11		5,521	2,117	3,123	2,927	33,843
1993	41,800	4		5,017	950	5,460	3,324	56,555
1994	46,554	3		6,437	2,321	6,143	4,995	66,453
1995	53,210	5		3,251	2,588	5,566	3,039	67,659
1996	18,581	3		5,093	1,608	4,464	2,931	32,680
1997	21,922	0		2,466	1,282	3,161	2,628	31,459
1998	18,265	3		1,759	1,074	1,778	2,300	25,179
1999	26,833	6		1,605	989	2,387	2,347	34,167 <sup>(b)</sup>
2000	43,354	0		2,171	973	3,496	1,004	50,998 <sup>(b)</sup>
2001	36,115	0		1,794	1,027	3,310	997	43,243 <sup>(b)</sup>
2002	30,980	112		1,404	1,190	3,318	1,275	38,279 <sup>(b)</sup>
2003	10,435	24		1,444	1,540	2,801	975	17,219 <sup>(b)</sup>
2004	11,017	53		1,295	769	2,477	970	16,581 <sup>(b)</sup>
2005	8,987	15		572	938	5,178	1,121	16,811 <sup>(b)</sup>
2006	7,566	15		477	864	3,977	679	13,578 <sup>(b)</sup>
2007	7,091	7		211	676	2,324	613	10,922 <sup>(b)</sup>
2008	6,241	9		385	871	981	160	8,647 <sup>(b)</sup>
2009	5,395	3		285	883	846	93	7,505 <sup>(b)</sup>
2010	19,982	1		506	238	1,665	223	22,615 <sup>(b)</sup>
2011	24,429	6		377	171	915	295	26,193 <sup>(b)</sup>
2012	7,318	2		261	210	578	115	8,484 <sup>(b)</sup>
2013	16,649	2		290	131	877	227	18,176 <sup>(b)</sup>
2014	10,210	7		291	177	479	227	11,391 <sup>(b)</sup>
Mean (2009–2013)	14,755	3		344	327	976	191	16,595
% change:								
2014 on 2013	-39			0	+35	-45	0	-37
2014 on 5-yr mean	-31			-15	-46	-51	19	-31

Note: Data for 2014 are provisional.

Key: <sup>(a)</sup> Returns not required before 1989. It is unusual for salmonids positively identified as salmon to be caught in this sea trout fishery in any numbers; some reported fish may have been misidentified in some years. Hence, no period means are reported.

<sup>(b)</sup> Includes a small number of fish caught & released (various regions).

**Table 12. Declared number of salmon caught by rods and the number and percentage of salmon released, 1993-2014.**

Year	Previous Environment Agency Region						NRW Wales	E&W Total #
	NE	Thames	Southern <sup>(a)</sup>	SW	Midlands	NW		
<b>Number caught</b>								
1993	1,696	2	84	2,806	336	5,055	4,080	14,059
1994	1,939	11	432	5,213	555	8,840	7,901	24,891
1995	2,201	13	302	2,554	442	6,348	4,146	16,006
1996	2,514	34	384	2,681	643	5,720	5,468	17,444
1997	2,445	2	149	2,372	312	4,144	3,622	13,047
1998	2,941	0	366	2,919	186	6,359	4,325	17,109
1999	2,670	1	253	1,881	185	4,133	3,369	12,493
2000	3,600	0	316	2,487	327	6,814	4,049	17,596
2001	3,733	0	405	1,396	273	4,209	4,351	14,383
2002	3,967	0	531	1,737	195	5,532	3,312	15,282
2003	3,507	0	225	1,266	333	3,547	2,632	11,519
2004	6,788	0	609	2,799	319	10,022	6,648	27,332
2005	5,933	0	438	1,725	430	8,446	4,408	21,418
2006	5,774	0	331	1,802	356	6,771	4,355	19,509
2007	4,872	0	466	2,071	280	7,151	5,136	19,984
2008	5,634	0	711	2,686	294	8,065	6,122	23,512
2009	4,421	0	391	1,648	213	5,532	3,356	15,563
2010	7,947	2	590	2,628	235	8,074	5,676	25,153
2011	8,373	0	606	2,402	362	6,672	4,784	23,199
2012	6,465	0	364	2,022	249	4,609	4,740	18,450
2013	6,469	0	271	1,085	332	3,539	3,224	14,920
2014	3,957	0	323	729	190	2,399	1,954	9,552
<b>Number released</b>								
1993	191	1	36	262	17	668	273	1,448
1994	322	0	69	745	36	1,253	802	3,227
1995	555	7	83	526	32	1,393	593	3,189
1996	732	25	88	510	57	1,332	684	3,428
1997	797	1	107	586	30	1,131	480	3,132
1998	1,037	0	222	1,077	31	2,019	979	5,371
1999	1,348	1	137	898	65	1,795	1,203	5,447
2000	1,888	0	247	1,152	103	2,816	1,264	7,470
2001	1,855	0	397	635	128	1,779	1,347	6,143
2002	2,257	0	528	920	73	2,534	1,346	7,658
2003	2,265	0	225	746	153	1,859	1,172	6,425
2004	3,612	0	609	1,572	174	4,672	2,487	13,211
2005	3,426	0	438	1,130	271	4,376	2,310	11,983
2006	3,283	0	331	1,342	210	3,450	2,285	10,959
2007	2,545	0	466	1,406	145	3,838	2,517	10,922
2008	2,831	0	711	1,825	155	4,360	3,153	13,035
2009	2,533	0	391	1,080	119	3,236	1,736	9,096
2010	4,714	2	587	1,795	133	4,807	2,974	15,012
2011	5,232	0	604	1,678	222	3,904	2,766	14,406
2012	3,995	0	358	1,454	185	2,774	3,186	11,952
2013	4,444	0	266	870	227	2,320	2,331	10,458
2014	2,927	0	318	602	148	1,850	1,523	7,368

**Table 12. continued**

<b>Number retained</b>								
1993	1,505	1	48	2,544	319	4,387	3,807	12,611
1994	1,617	11	363	4,468	519	7,587	7,099	21,664
1995	1,646	6	219	2,028	410	4,955	3,553	12,817
1996	1,782	9	296	2,171	586	4,388	4,784	14,016
1997	1,648	1	42	1,786	282	3,013	3,142	9,915
1998	1,904	0	144	1,842	155	4,340	3,346	11,738
1999	1,322	0	116	983	120	2,338	2,166	7,046
2000	1,712	0	69	1,335	224	3,998	2,785	10,126
2001	1,878	0	8	761	145	2,430	3,004	8,240
2002	1,710	0	3	817	122	2,998	1,966	7,624
2003	1,242	0	0	520	180	1,688	1,460	5,094
2004	3,176	0	0	1,227	145	5,350	4,161	14,121
2005	2,507	0	0	595	159	4,070	2,098	9,435
2006	2,491	0	0	460	146	3,321	2,070	8,550
2007	2,327	0	0	665	135	3,313	2,619	9,062
2008	2,803	0	0	861	139	3,705	2,969	10,477
2009	1,888	0	0	568	94	2,296	1,620	6,467
2010	3,233	0	3	833	102	3,267	2,702	10,141
2011	3,141	0	2	724	140	2,768	2,018	8,793
2012	2,470	0	6	568	64	1,835	1,554	6,498
2013	2,025	0	5	215	105	1,219	893	4,462
2014	1,030	0	5	127	42	549	431	2,184
<b>% of fish released</b>								
1993	11		43	9	5	13	7	10
1994	17		16	14	6	14	10	13
1995	25		27	21	7	22	14	20
1996	29		23	19	9	23	13	20
1997	33		72	25	10	27	13	24
1998	35		61	37	17	32	23	31
1999	50		54	48	35	43	36	44
2000	52		78	46	31	41	31	42
2001	50		98	45	47	42	31	43
2002	57		99	53	37	46	41	50
2003	65		100	59	46	52	45	56
2004	53		100	56	55	47	37	48
2005	58		100	66	63	52	52	56
2006	57		100	74	59	51	52	56
2007	52		100	68	52	54	49	55
2008	50		100	68	53	54	52	55
2009	57		100	66	56	58	52	58
2010	59		99	68	57	60	52	60
2011	62		100	70	61	59	58	62
2012	62		98	72	74	60	67	65
2013	69		98	80	68	66	72	70
2014	74		98	83	78	77	78	77
Mean total catch – including fish caught & released (2009-2013)	6,735		444	1,957	278	5,685	4,356	19,457
% change:								
2014 on 2013	-39		+19	-33	-43	-32	-39	-36
2014 on 5-yr mean	-41		-27	-63	-32	-58	-55	-51

Key: # Totals include some fish of unknown region of capture.

<sup>[a]</sup> Prior to 2012, the Rivers Test and Itchen (Southern Region) were based upon owner's returns; data from 2012 are declared catches.

Notes: The 2014 figures are based on angler's catch returns processed up to 23 February 2015. Data for 2014 are provisional.

**Table 13. Declared weight and percentage of salmon caught (retained fish only) in coastal, estuarine and riverine fisheries in England & Wales, 1988–2014.**

Year	Coastal		Estuarine		Riverine		Total
	Wt (t)	%	Wt (t)	%	Wt (t)	%	Wt (t)
1988	218.1	55	53.0	13	123.6	31	394.8
1989	159.3	54	80.0	27	56.6	19	295.9
1990	212.4	63	65.5	19	60.3	18	338.1
1991	105.9	53	38.7	19	55.6	28	200.1
1992	90.7	53	39.6	23	40.2	24	170.5
1993	158.8	64	43.4	18	45.9	18	248.1
1994	183.5	57	58.4	18	81.9	25	323.8
1995	200.3	68	45.4	15	48.9	17	294.6
1996	83.3	45	42.3	23	57.5	31	183.2
1997	80.5	57	26.7	19	34.6	24	141.8
1998	65.2	53	19.4	16	38.2	31	122.9
1999	101.0	67	23.1	15	26.0	17	150.0
2000	156.6	72	25.4	12	36.9	17	218.8
2001	128.6	70	24.2	13	31.3	17	184.2
2002	107.9	67	24.4	15	28.7	18	161.0
2003	42.0	47	26.6	30	20.4	23	89.0
2004	39.2	35	19.4	17	52.8	47	111.4
2005	32.2	33	28.3	29	36.0	37	96.5
2006	29.5	37	20.7	26	29.6	37	79.8
2007	23.9	36	13.4	20	29.8	44	67.1
2008	21.7	34	8.1	13	34.0	53	63.7
2009	20.2	37	8.6	16	25.2	47	54.0
2010	63.8	59	8.8	8	36.2	33	108.7
2011	93.1	69	6.4	5	36.3	27	135.8
2012	26.1	45	4.6	8	27.2	47	58.0
2013	61.5	73	5.6	7	17.0	20	84.0
2014	38.4	74	4.4	8	8.8	17	51.5
Mean (1993–98)	128.6	59	39.3	18	51.2	23	219.1
Mean (1999–02)	123.5	69	24.3	14	30.7	17	178.5
Mean (2003–14)	41.0	49	12.9	15	29.4	35	83.3

Notes: Coastal catches in 2014 from North East coast nets and Anglian coastal nets, but previously included River Parrett putcher rank (last fished 1999), River Usk drift nets (1997) & putcher rank (1999), SW Wales coastal wade (1995) & seine nets (1997), River Ogwen seine nets (2000), River Seiont/Gwyrfaï seine nets (1997), River Dwyfawr seine nets (1999), N. Caernarvonshire seine nets (1996), River Clwyd sling (drift) nets (1997) and the SW Cumbria drift nets (2003). Riverine catches in 2014 from rod catches and River Eden coops; River Conwy basket trap (also operated in freshwater) was last fished in 2002. Estuarine fisheries include all other nets and fixed engines not mentioned above. Data for 2014 are provisional.

**Table 14. Declared number, weight and percentage of salmon released by rods, and declared number and weight of salmon released by nets in England & Wales, 1993-2014.**

Year	Salmon released by rods			Salmon released by nets	
	Number released	Weight (t)	% of declared catch	Number	Weight (t)
1993	1,448	5.26	10		
1994	3,227	12.19	13		
1995	3,189	12.11	20		
1996	3,428	13.99	20		
1997	3,132	13.77	24		
1998	5,371	20.98	31		
1999	5,447	23.87	44	118	0.4
2000	7,470	30.70	42	171	0.7
2001	6,143	25.50	43	176	0.4
2002	7,658	31.80	50	234	0.9
2003	6,425	28.20	56	107	0.5
2004	13,211	51.70	48	143	0.5
2005	11,983	49.80	56	84	0.4
2006	10,959	42.50	56	72	0.3
2007	10,922	42.00	55	70	0.3
2008	13,035	49.80	55	88	0.3
2009	9,096	37.00	58	62	0.3
2010	15,012	53.38	60	61	0.2
2011	14,406	62.40	62	411	1.5
2012	11,952	53.89	65	56	0.2
2013	10,458	45.26	70	30	0.1
2014	7,368	31.44	77	73	0.2

Notes: Many of the salmon released by nets are fish caught pre June, which, since 1999, are required to be released.  
A proportion of the salmon released by nets have previously been due to an agreement between the Environment Agency and netmen fishing the estuary of the River Avon (Hants); this fishery ceased to operate in 2012  
There was no requirement for net caught salmon to be released prior to 1999.  
Data for 2014 are provisional.

**Table 15. Declared number and percentage of salmon caught by nets and rods taken before (<) and from (≥) 1 June, 1989 on.**

Year	Net catch (including released fish)				Rod catch (including released fish)			
	Number			%	Number #			%
	< 1 June	≥ 1 June	Total	< 1 June	< 1 June	≥ 1 June	Total	< 1 June
1989	4,742	64,198	68,940	6.9	3,199	11,529	14,728	21.7
1990	7,339	64,488	71,827	10.2	2,397	12,290	14,687	16.3
1991	3,637	34,038	37,675	9.7	2,240	11,496	13,736	16.3
1992	2,497	31,352	33,849	7.4	1,012	9,725	10,737	9.4
1993	1,630	54,936	56,566	2.9	865	13,194	14,059	6.2
1994	4,824	61,633	66,457	7.3	2,609	22,282	24,891	10.5
1995	4,888	62,771	67,659	7.2	2,141	13,865	16,006	13.4
1996	2,913	29,767	32,680	8.9	2,691	14,753	17,444	15.4
1997	1,528	29,931	31,459	4.9	1,335	11,278	12,613	10.6
1998	832	24,335	25,167	3.3	712	15,275	15,987	4.5
1999	116	34,043	34,159	0.3	920	11,211	12,131	7.6
2000	19	50,979	50,998	0.04	760	16,496	17,256	4.4
2001	47	43,196	43,243	0.11	708	13,675	14,383	4.9
2002	32	38,247	38,279	0.08	815	14,250	15,065	5.4
2003	42	17,177	17,219	0.24	1,037	10,373	11,410	9.1
2004	35	16,546	16,581	0.21	1,168	25,777	26,945	4.3
2005	29	16,782	16,811	0.17	1,652	19,239	20,891	7.9
2006	17	13,561	13,578	0.13	1,618	17,891	19,509	8.3
2007	14	10,908	10,922	0.13	908	18,733	19,641	4.6
2008	17	8,630	8,647	0.20	1,068	22,444	23,512	4.5
2009 <sup>[a]</sup>	1	7,504	7,505	0.01	925	14,638	15,563	5.9
2010 <sup>[a]</sup>	1	22,614	22,615	0.00	682	23,811	24,493	2.8
2011 <sup>[b]</sup>	367	25,826	26,193	1.40	1,255	21,383	22,638	5.5
2012	59	8,425	8,484	0.70	1,175	17,025	18,200	6.5
2013	30	18,146	18,176	0.17	1,236	13,541	14,777	8.4
2014	47	11,417	11,464	0.41	955	8,597	9,552	10.0
Mean (1994-98)	2,997	41,687	44,684	6.7	1,898	15,491	17,388	10.9
Mean (1999-14)	55	21,500	21,555	0.3	1,055	16,818	17,873	5.9

Notes: National measures to protect 'spring' salmon introduced on April 15 1999 - required compulsory catch and release of all rod caught salmon prior to June 16, and closed most net fisheries prior to June 1. Those net fisheries still allowed to operate before June (mainly targeted at sea trout) are required to release all salmon alive.

Rod catch data only include fish for which date of capture recorded; data may differ from total catch (Table 10).

Data for 2014 are provisional.

Key: # Excludes fish for which no capture date recorded.

<sup>[a]</sup> No requirement to record net-released fish on new logbooks, so pre June catch under-estimated.

<sup>[b]</sup> The increase in the pre-June catch from 2011 reflects the fact that salmon caught and released by T&J nets operating in the NE Region were not recorded over the period 1999-2010.



**Table 16. Declared number of salmon caught by rods, and number and percentage of fish released, by weight category and season, 1998 on.**

Season Wt. category (kg)	April to June			July to August			September to October			April to October		
	<3.6	3.6–6.4	>6.4	<3.6	3.6–6.4	>6.4	<3.6	3.6–6.4	>6.4	<3.6	3.6–6.4	>6.4
<b>Number caught</b>												
1998	523	753	111	3782	857	222	5767	2045	562	10,072	3,655	896
1999	354	864	262	1283	627	203	3667	2209	879	5,303	3,699	1,345
2000	388	771	206	2495	818	240	5813	3111	896	8,695	4,700	1,342
2001	205	971	203	1758	1041	200	4290	2536	724	6,253	4,548	1,127
2002	377	1014	300	2033	767	173	4434	2728	775	6,844	4,508	1,247
2003	282	817	241	885	839	188	2879	2400	862	4,046	4,056	1,292
2004	516	832	241	3374	1587	283	11124	6120	1212	15,014	8,539	1,736
2005	546	1454	327	2007	1198	169	8048	4941	974	10,601	7,593	1,470
2006	567	1505	269	1422	779	110	9176	3593	766	11,165	5,877	1,145
2007	565	931	161	2936	1897	233	7876	3445	707	11,377	6,273	1,101
2008	719	1,381	215	3,367	2,213	288	8,908	4,028	1,018	12,994	7,622	1,521
2009	500	849	172	2,163	1,933	221	4,955	3,096	802	7,618	5,878	1,195
2010	441	469	117	3740	1418	215	11284	4986	1099	15,465	6,873	1,431
2011	643	1,426	364	2,606	2,777	574	6,831	5,255	1,567	10,080	9,458	2,505
2012	597	1,395	512	2,504	2,750	558	4,476	3,762	1,185	7,577	7,907	2,255
2013	437	1,200	486	1,644	1,146	228	5,202	3,130	1,006	7,283	5,476	1,720
2014	374	856	201	1,213	1,019	175	2,794	2,072	599	4,381	3,947	975
<b>Number released</b>												
1998	136	113	20	643	197	40	2,076	900	253	2,855	1,210	313
1999	209	570	194	295	163	61	1,430	994	466	1,934	1,727	721
2000	221	532	148	499	229	72	2,325	1,431	502	3,045	2,192	722
2001	119	602	138	422	302	52	1,673	1,141	420	2,214	2,045	610
2002	241	659	213	488	207	57	2,084	1,473	488	2,813	2,339	758
2003	214	629	193	239	235	64	1,382	1,392	595	1,835	2,256	852
2004	283	576	143	1074	501	116	5,154	2,962	707	6,511	4,039	966
2005	464	1105	265	715	439	67	4,240	2,661	598	5,419	4,205	930
2006	499	1234	239	583	304	54	4,496	2,048	498	5,578	3,586	791
2007	436	666	142	1181	726	109	4,253	1,981	448	5,870	3,373	699
2008	507	948	170	1547	874	116	4,827	2,307	622	6,881	4,129	908
2009	378	630	148	957	743	104	2,925	1,963	549	4,260	3,336	801
2010	339	367	104	1743	604	107	6751	3141	802	8,833	4,112	1,013
2011	481	1,038	298	1,380	1,289	301	4,242	3,351	1,092	6,102	5,678	1,691
2012	449	1,046	443	1,391	1,371	334	2,960	2,502	871	4,800	4,919	1,648
2013	367	996	456	874	619	137	3,553	2,292	794	4,794	3,907	1,387
2014	323	737	190	791	601	104	2,228	1,656	510	3,342	2,994	804
<b>Percentage (%) released</b>												
1998	26	15	18	17	23	18	36	44	45	28	33	35
1999	59	66	74	23	26	30	39	45	53	36	47	54
2000	57	69	72	20	28	30	40	46	56	35	47	54
2001	58	62	68	24	29	26	39	45	58	35	45	54
2002	64	65	71	24	27	33	47	54	63	41	52	61
2003	76	77	80	27	28	34	48	58	69	45	56	66
2004	55	69	59	32	32	41	46	48	58	43	47	56
2005	85	76	81	36	37	40	53	54	61	51	55	63
2006	88	82	89	41	39	49	49	57	65	50	61	69
2007	77	72	88	40	38	47	54	58	63	52	54	63
2008	71	69	79	46	39	40	54	57	61	53	54	60
2009	76	74	86	44	38	47	59	63	68	56	57	67
2010	77	78	89	47	43	50	60	63	73	57	60	71
2011	75	73	82	53	46	52	62	64	70	61	60	68
2012	75	75	87	56	50	60	66	67	74	63	62	73
2013	84	83	94	53	54	60	68	73	79	66	71	81
2014	86	86	95	65	59	59	80	80	85	76	76	82

Notes: 1998 Pre national byelaw.

1999 National byelaw requiring compulsory catch and release before 16 June introduced on 15 April.

2000 First full year of national catch and release byelaw.

Only includes data for fish where weights reported; totals may differ from declared catches (Table 10).

Data for 2014 are provisional.

**Table 17. Provisional declared number and percentage of small (<3.6kg) and large (>3.6kg) salmon caught by net fisheries in England & Wales, 2014 (excluding released fish).**

EA Region/NRW	Small salmon (1SW)		Large salmon (MSW)		Total
	(<3.6 kg)	%	(>3.6 kg)	%	
North East	5,564	54	4,646	46	10,210
South West	193	66	98	34	291
Midlands	43	24	134	76	177
North West	223	47	256	53	479
Wales	104	46	123	54	227
<b>Total</b>	<b>6,127</b>	<b>54</b>	<b>5,257</b>	<b>46</b>	<b>11,384</b>

**Table 18. Provisional declared number and percentage of 1SW (grilse) and MSW salmon caught by selected rod fisheries (including fish caught and released), 2014.**

EA Region / NRW	River	No. 1SW	%	No. MSW	%
NE	Coquet	152	57	116	43
	Tyne	1081	43	1451	57
	Wear	600	66	306	34
Southern	Itchen	144	76	46	24
	Test	87	66	44	34
SW	Hants Avon	14	30	33	70
	Frome	17	44	22	56
	Exe	48	70	21	30
	Teign	44	77	13	23
	Dart	10	71	4	29
	Tavy	24	73	9	27
	Tamar	60	61	38	39
	Lynher	20	57	15	43
	Fowey	64	86	10	14
	Camel	56	80	14	20
	Taw	27	25	82	75
	Torridge	11	31	25	69
	Lyn	5	41	7	59
Midlands	Severn	44	23	146	77
NW	Ribble	354	63	208	37
	Lune	250	71	101	29
	Kent	97	88	13	12
	Leven	41	89	5	11
	Irt	21	81	5	19
	Ehen	125	76	39	24
	Derwent	191	70	83	30
	Eden	220	52	204	48
	Border Esk	221	63	130	37
Wales	Wye	90	21	335	79
	Usk	146	38	243	62
	Ogmore	14	70	6	30
	Tywi	92	52	86	48
	Tawe	27	69	12	31
	Taf	16	39	25	61
	E & W Cleddau	25	66	13	34
	Teifi	119	52	112	48
	Dyfi	9	56	7	44
	Mawddach	28	67	14	33
	Ogwen	12	92	1	8
	Conwy	59	86	10	14
	Dee	151	55	126	45
<b>E&amp;W Total</b>		<b>4,816</b>	<b>54</b>	<b>4,180</b>	<b>46</b>

Note: Data only included for fish for which weight data provided on catch return and do not include all rivers; these data therefore differ from the total reported catch (Table 10).

**Table 19. Estimated number of 1SW and MSW salmon (corrected for under-reporting) and the percentage composition of MSW salmon caught by rods (including fish caught and released), 1992–2014.**

Year	Environment Agency Region										NRW Wales		E&W Total	
	NE		Southern		SW		Midlands		NW		1SW	MSW	1SW	MSW
	1SW	MSW	1SW	MSW	1SW	MSW	1SW	MSW	1SW	MSW				
1992	1,085	723	235	29	3,186	476	112	175	4,029	945	2,282	1,074	10,927	3,422
1993	966	729	465	82	3,216	706	145	192	5,245	999	4,788	1,197	14,825	3,905
1994	1,173	660	277	156	4,172	1,043	217	339	7,162	1,680	5,609	2,291	18,611	6,169
1995	1,270	1,082	218	65	1,914	860	71	402	5,380	1,102	2,769	1,491	11,622	5,002
1996	1,246	1,405	262	97	1,674	1,116	90	603	4,620	1,228	3,431	2,287	11,322	6,736
1997	1,325	1,084	120	30	1,932	483	54	266	3,780	667	2,382	1,021	9,593	3,551
1998	2,226	909	378	24	2,543	501	66	131	5,975	699	3,548	843	14,736	3,107
1999	1,586	1,351	206	72	1,386	683	70	132	3,589	955	2,278	1,175	9,115	4,368
2000	2,188	1,618	292	56	2,270	441	200	139	6,507	807	3,196	816	14,653	3,877
2001	2,628	1,478	344	61	1,275	261	90	210	3,936	694	3,638	1,149	11,911	3,853
2002	2,924	1,440	520	64	1,452	459	92	123	5,233	852	2,550	1,093	12,771	4,031
2003	2,353	1,505	151	74	947	446	117	249	3,121	780	1,766	1,129	8,455	4,183
2004	5,222	2,245	528	81	2,633	446	123	228	9,790	1,234	5,927	1,386	24,223	5,620
2005	5,481	2,088	306	132	1,404	494	151	322	7,804	1,487	3,588	1,261	18,734	5,784
2006	4,637	1,715	256	76	1,388	595	145	247	5,810	1,639	3,593	1,198	15,829	5,470
2007	3,798	1,431	382	84	1,615	656	171	136	6,725	1,029	4,110	1,267	16,801	4,603
2008	4,651	1,547	633	78	2,245	710	106	217	7,724	1,147	5,387	1,347	20,746	5,046
2009	3,686	1,346	157	95	1,326	477	74	157	4,686	1,346	2,323	1,163	12,252	4,584
2010	6,119	2,623	498	88	2,486	335	106	153	7,194	1,687	5,027	1,103	21,430	5,989
2011	4,422	4,788	420	183	1,882	760	105	293	4,564	2,775	3,066	2,126	14,460	10,925
2012	3,528	3,584	273	128	1,219	1,005	68	206	2,877	2,193	2,198	3,016	10,162	10,132
2013	3,978	3,138	140	158	778	416	76	289	2,790	1,103	1,828	1,719	9,590	6,822
2014	2,153	2,200	256	100	463	339	48	161	1,738	901	953	1,197	5,610	4,897
Mean (2009–2013)	4,347	3,096	298	130	1,538	599	86	220	4,422	1,821	2,888	1,825	13,579	7,690
% change:														
2014 on 2013	-46	-30	+82	-37	-40	-18	-37	-44	-38	-18	-48	-30	-41	-28
2014 on 5-year mean	-50	-29	-14	-24	-70	-43	-44	-27	-61	-51	-67	-34	-59	-36

**Percentage MSW**

Year	Environment Agency Region										NRW Wales		E&W Total	
	NE		Southern		SW		Midlands		NW		1SW	MSW	1SW	MSW
	1SW	MSW	1SW	MSW	1SW	MSW	1SW	MSW	1SW	MSW				
1992	40	11	13	61	19	32	24							
1993	43	15	18	57	16	20	21							
1994	36	36	20	61	19	29	25							
1995	46	23	31	85	17	35	30							
1996	53	27	40	87	21	40	37							
1997	45	20	20	83	15	30	27							
1998	29	6	16	66	10	19	17							
1999	46	26	33	65	21	34	32							
2000	43	16	16	41	11	20	21							
2001	36	15	17	70	15	24	24							
2002	33	11	24	57	14	30	24							
2003	39	33	32	68	20	39	33							
2004	30	13	14	65	11	19	19							
2005	28	30	26	68	16	26	24							
2006	27	23	30	63	22	25	26							
2007	27	18	29	44	13	24	22							
2008	25	11	24	67	13	20	20							
2009	27	38	26	68	22	33	27							
2010	30	15	12	59	19	18	22							
2011	52	30	29	74	38	41	43							
2012	50	32	45	75	43	58	50							
2013	44	53	35	79	28	48	42							
2014	51	28	42	77	34	56	47							
Mean (2009–2013)	42	30	28	72	29	39	36							

Note: Data for 2014 are provisional.





Figure 6. Declared number of salmon caught by nets and fixed engines, 1971–2014. (Note: y-axes not to same scale.)



**Figure 7. Declared number of salmon caught by rods and the number of salmon released (upper, blue part of histogram), 1993-2014. (Note: y-axes not to same scale.)**




Figure 8. Percentage of the declared total catch of salmon (caught and retained only) taken in coastal, estuarine and riverine fisheries, 1988–2014.






Figure 9. The number and percentage of the declared salmon catch released by anglers, 1993–2014.



**Figure 10. Declared number of salmon caught by nets and fixed engines in England & Wales and the percentage of the catch taken in the north east coast fishery, 1956-2014.**



**Figure 11. Declared number of salmon caught by rods in England & Wales, 1956-2014.**




Figure 12. Declared number and percentage of salmon caught by (a) nets and (b) rods before 1 June, 1989–2014.






Figure 13. Percentage of rod caught fish released by anglers by weight category, 1998–2014.







**Figure 14. Variation in the proportions of 1SW and older salmon returning to the Rivers Dee, Tamar, Lune and Tyne over available time series.**




**Figure 15. Estimated percentage (%) of salmon >8lb (3.6kg) caught in net and fixed engine fisheries (excluding NE Region), 1999–2014.**




**Figure 16. Estimated percentage (%) of salmon >8lb (3.6kg) caught in the north east coast net fishery (as declared by netmen), 1965–2014.**




**Figure 17. Estimated percentage of principal salmon rivers with  $\geq 50\%$ , 25-49% and  $\leq 24\%$  of MSW salmon in the declared rod catch, 1997–2014.**



**Figure 18. Estimated number (histogram) and percentage (solid line) of MSW salmon caught by rods, 1992–2014. (Note: y-axes not to same scale.)**



**Figure 19. Estimated total number (corrected for under-reporting) of 1SW and MSW salmon caught by rod fisheries in England and Wales (including fish caught and released), 1992–2014.**



**Figure 20. Estimated total catch of 1SW and MSW salmon in England and Wales (fish caught and killed only), 1971–2014, as used in the ICES PFA assessment.**

## 5. CATCH PER UNIT EFFORT (CPUE)

Since catch levels are influenced strongly by the level of fishing effort, catch per unit effort (CPUE) data are commonly used as well as the declared catch in order to help evaluate the status of stocks. However, the relationship between CPUE and abundance can be influenced by confounding factors in both rod and net fisheries. It should also be remembered that net and rod fisheries operate sequentially (the net fisheries exploit the returning fish first), and over different time periods (fishing seasons). Rod fisheries are active over a longer period and typically extend into the early autumn after net fisheries have ceased. Thus, changes in patterns of run-timing may also impact on CPUE values in the different fisheries.

- **Nets** – Regional CPUE data for net fisheries for the period since 1997 are presented in Table 20. These data are based on the number of tides fished by netsmen, except in the North East Region where the number of days fished is used. In order to provide comparable time series, the data only include fishing gears that have operated in a consistent manner over the period. Plots of the standardised CPUE Z-scores for the various regions and for net fisheries overall (expressed as a 2-year moving average) are provided in Figure 21.
- **Rods** – Regional CPUE data for rod fisheries for the period since 1997 (expressed as the number of salmon caught per 100 days fished) are presented in Table 21. Plots of the standardised CPUE Z-scores for the various regional rod fisheries and the overall rod CPUE for England and Wales are provided in Figure 22 for the same period. Individual CPUE data for all the major salmon rivers in England and Wales are reported in the annual catch statistics reports (e.g. Environment Agency, 2014). The trends in rod CPUE for the different regions show a reasonable degree of coherence and available evidence from selected rivers where we have estimates of returning stock size, as well as CPUE, suggests rod CPUE values provide a reasonable indicator of stock abundance (Figure 23).

### Overview of CPUE in 2014

The overall CPUE for nets and fixed engines in 2014 was 16% lower than the previous 5-year mean (2009-2013). CPUE declined in all regions, ranging from a 13% fall in the former North East Region to a 37% fall in the former South West Region. Normalised CPUE values (Z-scores) for the various former regions and an overall average (Figure 21) indicate that CPUE, and by inference abundance, peaked during the period 2000-02, then declined steadily until 2009 (which had the lowest CPUE of the time series), before increasing again between 2010 and 2011. Overall CPUE in 2014 fell from 2013 and was close to the long-term average. An earlier analysis of net CPUE and river flow suggests above average flows in July (when a high proportion of the net catch typically occurs) tend to result in reduced CPUE values.

Rod CPUE fell in 2014 compared with 2013 in all former regions, except Southern Region where there was a substantial upturn in CPUE, and fell by 13% overall (Table 21). While regional CPUE estimates were generally lower, values remained within reported ranges, albeit at the lower end of these ranges. This was despite the particularly low rod catch. Normalised CPUE values (Z-scores) for rod fisheries (Figure 22) suggest a largely positive trend in CPUE, and by inference abundance (Figure 23), over the available time series, although there has been something of a downturn in the last two years. Overall rod CPUE in 2014 was close to the long-term average.

**Table 20. Mean catch per unit of effort (CPUE) for salmon net fisheries, 1997–2014.**

Year	Environment Agency Region					NRW Wales	E&W (incl. NE Drift nets, excl. Midlands)
	NE Drift nets (June-August)	NE	SW	Midlands	NW		
1997	6.48	4.40	0.70	0.23	0.63	0.07	1.23
1998	5.92	3.81	1.25	0.24	0.46	0.08	1.17
1999	8.06	4.88	0.79	0.31	0.52	0.20	1.35
2000	13.06	8.11	1.01	0.33	1.05	0.18	2.19
2001	10.34	6.83	0.71	0.33	0.71	0.16	1.77
2002	8.55	5.59	1.03	0.53	0.90	0.23	1.66
2003	7.13	4.82	1.24	0.60	0.62	0.11	1.43
2004	8.17	5.88	1.17	0.36	0.69	0.11	1.65
2005	7.23	4.13	0.60	0.60	1.28	0.09	1.35
2006	5.60	3.20	0.66	0.51	0.82	0.09	1.04
2007	7.24	4.17	0.33	0.51	0.75	0.05	1.14
2008	5.41	3.59	0.63	0.64	0.34	0.06	0.96
2009	4.76	3.08	0.53	0.64	0.51	0.04	0.89
2010	17.03	8.56	0.99	0.26	0.47	0.09	2.08
2011	19.25	9.93	0.63	0.14	0.34	0.10	2.25
2012	6.80	5.35	0.69		0.31	0.21	1.36
2013	11.06	8.22	0.54		0.39	0.08	1.89
2014	10.30	6.12	0.43		0.31	0.07	1.42
Mean (2009–2013)	11.78	7.03	0.68	0.34	0.40	0.10	1.69
No. fisheries	2	4	4	1	6	6	20
% change (2014 on 5-year mean)	-13	-13	-37		-24	-27	-16

Notes: Fisheries were selected on the basis that they were fished consistently during the period. Data are expressed as catch per licence-tide, except for the North East, for which data are recorded as catch per licence-day. From 2012, the fishery operating in the Severn (Midlands Region) has been limited by a catch limit (cap); the Midlands data have therefore been removed from the combined E&W total for the whole time series. Data for 2014 are provisional.

**Table 21. Mean catch per unit of effort (CPUE) for salmon rod fisheries in each Region, 1997–2014.**

Year	Environment Agency Region						NRW Wales	England & Wales
	NE	Thames	Southern	SW	Midlands	NW		
1997	5.0	0.6	3.1	5.2	1.7	5.3	2.6	4.0
1998	6.5	0.0	5.9	7.5	1.3	8.6	3.9	6.0
1999	7.4	0.3	3.1	6.3	2.1	7.4	3.5	5.5
2000	9.2	0.0	5.2	8.8	4.9	11.7	4.4	7.9
2001	11.3	0.0	11.0	6.6	5.4	15.4	5.5	8.7
2002	9.4	0.0	18.3	6.0	3.5	10.0	3.6	6.8
2003	9.7	0.0	8.8	4.7	5.2	8.3	2.9	5.7
2004	14.7	0.0	18.8	9.6	5.5	17.4	6.6	11.4
2005	12.4	0.0	12.7	6.2	6.6	13.9	4.5	9.0
2006	14.2	0.0	15.6	8.7	6.6	13.3	5.9	10.1
2007	11.7	0.0	18.0	8.7	5.7	14.2	6.0	9.6
2008	12.7	0.0	21.8	10.9	5.8	15.3	7.3	10.5
2009	9.5	0.0	13.7	5.7	3.6	9.3	3.6	6.6
2010	16.7	2.8	17.1	9.9	4.3	14.1	6.5	10.2
2011	17.5	0.0	14.5	9.4	6.5	11.4	6.0	10.9
2012	15.4	0.0	17.3	9.2	6.3	9.1	6.5	10.6
2013	16.7	0.0	10.0	5.9	7.9	7.7	5.7	8.9
2014	13.0	0.0	14.8	5.6	5.5	7.3	4.8	7.8
Mean (2009–2013)	15.1	0.6	14.5	8.0	5.7	10.3	5.7	9.5
% change:								
2014 on 2013	-22		+48	-5	-31	-5	-16	-13
2014 on 5-yr mean	-14		+2	-31	-5	-29	-16	-18

Notes: Based only on catch returns for which effort data have been reported. CPUE is expressed as number of salmon (including released fish) caught per 100 days fished. Data for 2014 are provisional.




Figure 21. Normalised catch per unit of effort (CPUE) (z-score) for salmon net fisheries, 1997–2014.





Figure 22. Normalised catch per unit of effort (CPUE) (z-score) for salmon rod fisheries, 1997–2014.





**Figure 23. The relationship between mean rod CPUE and mean stock size for the Rivers Fowey, Dee and Lune, 1997–2013.**

## 6. EXPLOITATION RATES

Care is required in trying to draw general conclusions about current stock status from catches alone. The actual relationship between catch and stock abundance depends upon exploitation rates (i.e. the proportion of the salmon population taken in the catch - both retained fish and those released), although it is important to remember that fishing effort and catchability (the proportion of the stock taken per unit of fishing effort) can be influenced by factors such as river flow, angler activity and changes in run timing. Exploitation rates can be estimated where there is a fishery-independent measure of the salmon run, such as that obtained from fish counters and traps (Table 24 and Figure 28), and these data can then be compared against the catch (both total catch and retained fish) to estimate exploitation rates (Table 22 and Figure 24). These show varying trends, but the 'true' exploitation rates (i.e. fish retained) show a marked decline in most rivers over the available time series, due largely to the increasing use of catch-and-release.

### Overview of exploitation rates in 2014

Exploitation rates in rod fisheries in 2014 were mainly below those in 2013, with the majority of values well below the average of the previous five years. This may have been affected by the relatively poor runs of fish in many rivers (Section 7) and the low flows and poor conditions for angling in many catchments (Section 9.2). Viewed over available time series, exploitation rates on some rivers (e.g. Tyne and Leven) appear to have been increasing, while others (e.g. Lune) have been decreasing. However, the 'true' exploitation rates (i.e. fish retained) show a marked decline over the available time series in almost all rivers. This is largely attributable to C&R, which has increased from 10% to almost 80% over the past 2 to 3 decades, with mandatory catch and release now applying on some rivers. The exploitation rates for the net fisheries where estimates have been possible have either been reduced to zero or been greatly reduced, largely reflecting a major reduction in effort.

### Assessment of national trend in exploitation

Estimates of aggregated national exploitation rates, split by sea-age class, are required for use in the ICES annual assessment of stock status to estimate numbers of returning fish. The procedures used in deriving these estimates are described in the background report. The overall trends in national exploitation rate derived from this process are provided in Figure 25. These indicate that exploitation rates have fallen from about 50% for 1SW fish and 35-40% for MSW fish at the start of the period to a little over 15% and a little under 10%, respectively, in recent years due to the measures taken to control both legal and illegal fisheries. The decline in exploitation rates has been more pronounced since the early 1990s, but has levelled out more recently.

**Table 22. Estimated exploitation rates (%) for selected rod and net fisheries, 1988-2014. Data for rod fisheries distinguish exploitation rates for all fish (All: including fish released) and for retained fish only (Ret.) (Rates are corrected for under-reporting).**

Year	Rod Fisheries												Net Fisheries																			
	Region/NRW				Southern				SW				NW				Wales				NW											
	River		Tyne (e)		Test		Itchen		Hampshire/Avon		Frome (b)		Tamar		Fowey		Kent		Leven		Dee (b)		Dee (b)		Kent		Leven		Lune		Dee	
Wild/Hatchery		W		All		Ret.		W		All		Ret.		W		All		Ret.		W		All		Ret.		W		All		Ret.		
1988																																
1989																																
1990																																
1991																																
1992																																
1993																																
1994																																
1995																																
1996																																
1997																																
1998																																
1999																																
2000																																
2001																																
2002																																
2003																																
2004																																
2005																																
2006																																
2007																																
2008																																
2009																																
2010																																
2011																																
2012																																
2013																																
2014																																
Mean (2009-2013)																																
% change																																
2014 on 2013																																
2014 on 5-yr mean																																


Notes: It is unclear to what extent total rod exploitation rate (All) has been affected by catch and release and the repeat capture of fish; no correction factor has been applied. The entire catch from net fisheries is assumed killed.

Data for 2014 are provisional.


(a) Data based on Game & Wildlife Conservation Trust counter at East Stoke, and supplied courtesy of GWCT.

(b) Data derived from mark recapture experiment.

(c) Tyne values are provisional; work is ongoing with Newcastle University to further refine RSEs.



**Figure 24. Estimated exploitation rates (%), for selected rod and net fisheries in England & Wales, 1988–2014. For rod fisheries the figures display exploitation rates for all fish caught – ie. including fish released (blue dots) and fish killed (red triangles). Note that estimates for the Dee rods have been split by sea age class (1SW and MSW); all other estimates are combined for all ages.**



**Figure 25. Estimated national exploitation rates for 1SW and MSW salmon caught in England and Wales (fish caught and killed only), including non-reported catch, 1971–2014, as used in the ICES PFA assessment.**

# REPORT ON STATUS OF STOCKS IN 2014

## 7. STOCK MONITORING

The Environment Agency and Natural Resources Wales monitor both stocks and fishery performance in most rivers supporting salmon stocks, in England and Wales respectively. This includes operating counters, undertaking surveys of juvenile fish and collecting fishery statistics. These data provide the basis for assessing stock status and informing management decisions. In addition to protecting the abundance of stocks, managers need to maintain the diversity of stocks in terms of their biological characteristics. Measures of stock diversity potentially encompass a wide range of biological characteristics, but those of greatest significance for the management of stocks are the population structure within the river, the river-age of the emigrating smolts and the run-timing and sea-age composition of the returning adult stock. Such data tend to be derived from a small number of 'indicator' rivers. Further details on the various monitoring programmes are provided in the background report.

### Juvenile surveys (salmon fry and parr)

A programme of juvenile salmonid monitoring is carried out to identify spatial variation in juvenile populations and temporal trends in their abundance. The habitat at all sites is assessed such that the abundance of the juvenile salmon population at any site can be compared with standard reference conditions. A classification scheme is also applied such that the proportion of sites falling into different salmon abundance classes (Classes A to F) provides a measure of the health of the juvenile salmon populations for each river. Figure 26 presents the proportion of sites in each catchment that fall into the top three categories (Classes A to C) over the period 2009 to 2014. Thus, for catchments shaded red, less than 25% of sites fall within this category, while for those shaded green more than 75% of sites are at or above average. Overall, the majority (57%) of sites surveyed over the period were in the lowest two classes (Classes E or F). The proportion of sites surveyed that fell within Classes A to C varied between regions from 9% to 39%. Work is continuing to revise and update the procedures for collecting and interpreting juvenile data.

Figure 27 presents annual estimates of the overall percentage of sites falling within classes A to C viewed over the available time series. There is no clear trend in the data. However, the percentage of sites in classes A to C has increased in the last three years and the value for 2014 (38%) is among the highest in the time series.

### Upstream counts of adult salmon

Electronic fish counters or traps are operated on a number of catchments to provide estimates of the upstream run of adult salmon and sea trout. Where it is possible to separate the species, the counts are adjusted to provide estimates of the numbers of returning salmon. For some rivers (e.g. River Tyne) the time-consuming validation procedures mean data may not be available for the latest year. A number of facilities have also ceased to operate in recent years as a result of changes in operating procedures, counter / trap damage or due to budgetary restrictions. Available time series, including those that have been recently discontinued, are presented in Table 23 and Figure 28.

The majority of the available counts and returning stock estimates of adult fish entering freshwater in 2014 were below the average of the previous five years. For two rivers (Frome and Dee) the 2014 values were the lowest in the time series, while run estimates were among the lowest in

the available time series for some other rivers. The time series for different catchments indicate differing trends; some suggest something of an improvement over the last 10 years (e.g. Test, Itchen and Tees), while others indicate a recent decline (e.g. Caldey, Lune and Hampshire Avon).

### **Tagging investigations**

Tagging studies have often been employed to monitor stocks and to evaluate the outcome of different management initiatives, although tagging effort has fallen in recent years. In 2014, around 2,000 wild salmon smolts were microtagged and released in England and Wales to assess levels of marine survival; all these fish were also adipose fin-clipped. A further 64,000 hatchery parr and 9,000 wild parr were marked with adipose fin clips; the wild parr were also fitted with PIT tags. In addition, about 60 adult salmon were tagged for the assessment of returning stocks or in conjunction with the use of radio and acoustic tags in behaviour studies. Details of the tagged and marked salmon released each year around the whole North Atlantic are compiled annually by ICES. Details of the fish tagged in England and Wales in 2014 are provided in Table 24.

### **Marine survival**

Evidence from monitored rivers around the North Atlantic indicates that the survival of salmon during the marine phase of their life-cycle has declined in recent decades. Marine survival estimates, measured as percentage return rates, over an extended period are shown in Table 25 for the River Corrib (Ireland), River Bush (Northern Ireland) and River North Esk (Scotland) (data from ICES, 2014). Shorter time series for the Rivers Dee (Wales), Tamar and Frome (Table 25 and Figure 29) indicate similar low levels of marine survival in recent years. It was not possible to monitor adult returns on the Tamar in 2014, or to undertake any smolt tagging. There are therefore no updates possible for the Tamar time series. However, the programme resumed in 2015.

For both the Dee and Frome, the return rate (marine survival) of 1SW fish in 2014 (from the 2013 smolt cohort) was lower than that in 2013 and values for both rivers were among the lowest in the respective time series, indicating continued poor survival of 1SW fish. The poor return of 1SW fish is also reflected in the low proportion of 1SW fish in the 2014 catch. An estimate of survival for 2SW salmon in 2014 (from the 2012 smolt cohort) was only available for the Frome; this indicated a small increase on 2013, but was well within the range of recently observed values. Analysis of data for the River Dee has previously indicated a clear correlation between survival rates of particular smolt cohorts and the condition of the adult salmon returning to the river derived from these cohorts. This suggests a direct link between the ability of salmon to feed and grow at sea and their subsequent survival.

**Table 23. Validated counts and run estimates of salmon smolts and adults in selected monitored rivers, 1986–2014.**

Year	Smolts		Adults																								
	Southern		SW		NE		Thames		Southern		Hampshire Avon		SW		NW		Wales										
	Test (a)	Run estimate	Frome (a)	Test (a)	Tyne (b)	RSE1	Tees (a)	Thames (c)	Test (a)	RSE1	Itchen	RSE1	Frome (a)	RSE1	Tamar (e)	Fowey (f)	Lune	RSE1	Kent	RSE1	Leven	Caldew	T (h)	Dee	RSE2		
1986								81																			
1987								41																			
1988								288																			
1989								91																			
1990								63																			
1991								36																			
1992								247																			
1993								259																			
1994								143																			
1995								162																			
1996								122																			
1997								25																			
1998								6																			
1999								35																			
2000								53																			
2001								9																			
2002								22																			
2003								18																			
2004								7																			
2005								0																			
2006								0																			
2007								1																			
2008								9																			
2009								4																			
2010								4																			
2011								4																			
2012								15																			
2013								3																			
2014								n/a																			
Mean (2009–2013)								6																			

**Key to methods:**  
 (a) Smolt run estimates from 2010 are from a new trapping location further upstream, so shouldn't be compared directly with the earlier time series.  
 (b) Tyne RSEs have been updated based on video validation, but remain provisional pending work to further develop analytical methods for count specification.  
 (c) = adult salmon count.  
 (d) RSE1 = returning stock estimate (validated count + catch below counter).  
 (e) RSE2 = returning stock estimate (mark/recapture estimate).  
 (f) Index of run only - based on adult trap in barrage. Trap not operated after 2010; new counter now in place but provides combined salmon & sea trout count.  
 (g) Data adjusted for multiple entry (re-entry rate of 6.6% in 2002). Data relate to spawning year, i.e. 12 month period from March to February. Trap no longer operative from 2014.  
 (h) Due to counter malfunction, estimates for 2011–12 based on relationship between rod catch and RSE for the period 1990–2010.  
 (i) Slight under-estimate due to counter malfunction during May/June.  
 (j) Estimates informed by return rate of PIT tagged fish in addition to adult counter.  
 (k) Due to significant resistivity counter downtime, estimates based on a correlation between rod exploitation rate and validated counter estimates (from 2006–2012).



**Table 24. ICES Compilation of microtag, fin clip and external tag releases**

Marking Agency		Age	Life Stage	H/W	Stock Origin	Primary Tag or Mark	Number marked	Code or Serial	Secondary Tag or Mark	Release date	Release Location
Marking season: 2014 Country: UK (England and Wales)											
		Totals		Origin		Primary Tag or Mark		Other internal <sup>(a)</sup>		Total	
				Microtag		External Mark		Adipose Clip			
		Hatchery juvenile				1,981		64,121		64,121	
		Wild juvenile						9,052		11,081	
		Adult				630		58		688	
		Total fish marked				1,981		73,173		75,890	
Marking Agency	Age	Life Stage	H/W	Stock Origin	Primary Tag or Mark	Number marked	Code or Serial	Secondary Tag or Mark	Release date	Release Location	
EA North East	Various	Adult	W	Tyne	Floy tag	111	Green 1577 - 2124 & Blue 2233	None	Nov - Dec 2014	Tyne & tributaries	
Natural Resources Wales	Various	Smolt	W	Dee (Worthenbury)	CWT	64	01/42/97	Adipose	May 2014	Dee - Worthenbury	
Natural Resources Wales	Various	Smolt	W	Dee (Ceiriog)	CWT	1,917	01/42/88 and 01/42/34	Adipose	Apr-May 2014	Dee - Ceiriog	
Natural Resources Wales	Various	Adult	W	Dee	Floy tag	519	Various (blue and blue/green)	None	March-Oct 2014	Dee - Chester	
Natural Resources Wales	1+	Parr	H	Dee	Adipose clip	3,072		None	March 2014	Dee - Celyn	
Natural Resources Wales	0+	Fed Fry	H	Dee	Adipose clip	14,972		None	April 2014	Dee - Hirnant	
Natural Resources Wales	1+	Parr	H	Dee	Adipose clip	780		None	May 2014	Dee - main river	
Natural Resources Wales	1+	Parr	H	Dee	Adipose clip	2,340		None	June 2014	Dee - main river	
Natural Resources Wales	1+	Parr	H	Dee	Adipose clip	3,000		None	July 2014	Dee - Celyn	
Natural Resources Wales	1+	Parr	H	Dee	Adipose clip	3,000		None	Aug 2014	Dee - Celyn	
Natural Resources Wales	0+	Parr	H	Conwy	Adipose clip	11,300		None	Sept 2014	Conwy - upper	
Natural Resources Wales	0+	Parr	H	Conwy	Adipose clip	3,501		None	Oct 2014	Conwy - Roman Bridge	
Natural Resources Wales	0+	Parr	H	Conwy	Adipose clip	3,446		None	Nov 2014	Conwy - Llugwy	
Natural Resources Wales	1+	Smolt	H	Wye	Adipose clip	4,320		None	May-Jun 2014	Wye - Nant Gwyn Pond	
Natural Resources Wales	1+	Smolt	H	Wye	Adipose clip	3,800		None	May 2014	Wye - Bigsweir Pond	
Natural Resources Wales	1+	Smolt	H	Wye	Adipose clip	6,539		None	May 2014	Wye - Redbrook Pond	
Natural Resources Wales	1+	Smolt	H	Wye	Adipose clip	3,251		None	May 2014	Wye - Caradoc Pond	
Natural Resources Wales	1+	Smolt	H	Taff	Adipose clip	800		None	May 2014	Taff	
Natural Resources Wales	Various	Adult	H	Taff	PIT tag	21		None	Jan 2014	Taff - Penarth (broodstock)	
Natural Resources Wales	Various	Adult	H/W	Wye	PIT tag	27		None	Feb 2014	Wye (broodstock)	
Natural Resources Wales	Various	Adult	H/W	Wye	PIT tag	10		None	Nov 2014	Wye (broodstock)	
Cefas	Various	Parr	W	Itchen	Adipose clip	176	DC00xxxxxx	PIT tag	08/Oct/2014	Itchen	
Cefas/GWCT	Various	Parr	W	Frome	Adipose clip	8,876	3DD.003BCxxxxx	PIT tag	Sept 2014	Frome	
Cefas	Various	Smolt	W	Frome	Acoustic tag	48		None	April 2014	Frome	

Notes: <sup>(a)</sup> Includes PIT and radio/acoustic tags.

**Table 25. Estimated survival of wild smolts (%) to return to homewaters (prior to coastal fisheries) for index rivers in the UK and Ireland (from ICES, 2014 and Environment Agency/Natural Resources Wales/Cefas/GWCT data) for 1984 to 2013 smolt years.**

Smolt migration year	Ireland				UK (N. Ireland)				UK (England and Wales)												
	River Corrib		Burishoole		River Bush <sup>(a)</sup>		River North Esk <sup>(b)</sup>		Dee <sup>(c)</sup>		Tamar		1SW		95% CL		MSW		Frome <sup>(d)</sup>		
	1SW	2SW	1SW	1SW	1SW	1SW	MSW	1SW	MSW	1SW	95% CL	MSW	95% CL	1SW	95% CL	MSW	95% CL	1SW	MSW		
1984	26.2	2.0	7.8			6.0	4.0														
1985	18.9	1.8	7.9			13.6	5.4														
1986			8.7		31.3																
1987	16.6	0.7	12.0		35.1	10.4	3.9														
1988	14.6	0.7	10.1		36.2																
1989	6.7	0.7	3.5		25.0	6.6	4.2														
1990	5.0	0.6	9.2		34.7	6.0	3.1														
1991	7.3	1.3	9.5		27.8	7.6	3.1														
1992	7.3		7.6		29.0	10.9	6.5														
1993	10.8	0.1	9.5		14.5	6.1	6.3	3.6	2.5	2.2											
1994	9.8	1.4	9.4		27.1	10.9	3.6	1.2	1.2	1.3											
1995	8.4	0.1	6.8		8.4	3.8	2.7	1.8	0.4	0.7											
1996	6.3	1.2	9.2		31.0	5.9	2.7	1.7	2.1	1.3											
1997	12.7	0.8	8.2		19.8	7.2	4.2	2.9	3.4	1.9											
1998	5.5	1.1	5.3		13.4	2.6	1.4	2.4	3.7	3.6											
1999	6.4	0.9	8.1		16.5	6.8	3.8	8.3	12.4	11.8											
2000	9.4		9.0		10.1	6.0	2.8	1.1	0.9	0.8											
2001	7.2	1.1	7.6		12.4	4.7	2.9	5.1													
2002	6.0	0.5	6.5		11.3	2.2	2.0	1.4	0.7	0.9				3.6	2.1	1.4		5.6	1.7		
2003	8.3	2.1	8.3		6.8			1.7	0.4	0.4				6.1	2.0	1.8		4.8	0.9		
2004	6.3	0.8	5.8		6.8			1.1	1.0	0.5				6.0	2.3	1.5		5.3	2.9		
2005			5.3		5.9	6.7	2.8	1.6	0.5	0.4				6.4	1.6	1.2					
2006	1.2	0.9	13.0		14.0	3.3	3.4	1.2	1.5	0.9				3.8	1.3	5.3		5.1	2.2		
2007	0.9		8.4		8.3	5.0	4.0	1.1	0.9	0.7				7.6	3.8	3.3		5.7	1.3		
2008	1.7	1.0	8.2		4.0	6.4		2.0	1.3	1.5				1.6	0.9	0.9		3.1	1.6		
2009	6.0		8.9		5.9	9.0	8.7	2.1	1.1	1.0				8.2	2.1	1.9		7.7	2.6		
2010	2.9		7.5		4.0			1.9	0.7	1.3				3.4	1.5	5.0		8.6	2.4		
2011	2.4		10.8		2.7			1.9	0.3	0.5				1.1	1.6	1.9		1.5	1.8		
2012	1.5		9.4		10.8			4.9	0.3	0.5				2.5	1.4	1.9		3.2	2.1		
2013								1.8													
Mean (2008–12)	2.9	1.0	9.0		5.5	7.7	8.7	0.8	0.8	0.8				3.4	2.5	2.5		4.8	2.1		
Mean (2003–12)	3.5	1.2	8.6		6.9	6.1	4.7	0.8	0.8	0.8				4.7	2.5	2.5		5.0	2.0		

Key: <sup>(a)</sup> Based on microtagging, corrected for tagging mortality.

<sup>(b)</sup> Based on tagging with Carlin tags, not corrected for tagging mortality

<sup>(c)</sup> Based on microtagging with a 90% tag retention rate, not corrected for tagging mortality.

<sup>(d)</sup> Data based on Game & Wildlife Conservation Trust monitoring facilities at East Stoke, and supplied courtesy of GWCT; slightly revised methodology applied in latest year.

Notes: Data for 2013 smolt migration year are provisional.







Figure 26. Juvenile salmon abundance indices for each catchment, presented as percentage of surveys in classes A-C only, 2009–2014.



**Figure 27. Overall percentage of juvenile survey sites in England and Wales in classes A to C, 2003–2014.**



**Figure 28.** Counts from electronic counters (C) and monitoring traps (T), and returning stock estimates (RSE) (based on trapping and tagging, or validated counts plus catch below counter) for selected salmon stocks in England and Wales, 1988-2014. (Regression lines are indicative only and based on data from 1995 on (red squares); earlier data for some rivers indicated as blue diamonds. Note that the y-axes scales differ.)



**Figure 29. Estimated survival ( $\pm$  95 CLs where available) of wild smolts (%) to return to homewaters (prior to coastal fisheries) for (a) 1SW and (b) MSW salmon for the Rivers Dee, Tamar and Frome.**

## 8. ASSESSMENT OF STOCK STATUS

The status of individual river stocks in England and Wales is evaluated annually against stock conservation limits (CLs) and management targets (MTs) in line with the requirements of ICES and NASCO. A national assessment of the status of the salmon resource in England and Wales is also undertaken annually, using the Pre-fishery Abundance (PFA) and National Conservation Limit Models (Potter *et al.*, 2004), and reported to ICES to assist with the development of management advice for the distant water fisheries. Full details of these assessment approaches are provided in the background report.

### Status of river stocks in 2014

Egg deposition estimates for 2014 have been calculated for each of the 64 main salmon rivers in England and Wales and values, expressed as the proportion of the CL attained, are provided in Table 26 and illustrated in Figure 30.

Only 12 rivers (19%) were provisionally assessed as meeting their conservation limit (CL) in 2014, the lowest in the time series (Table 27). Thirty rivers (47%) were below 50% of their CL in 2014, the highest level in the time series. The marked downturn in levels of CL attainment in the last two years (Figure 31) follows a period of relatively good CL attainment between 2004 and 2012 when levels of egg deposition were typically higher than those in the preceding decade.

River-to-river variation in the proportion of the CL attained in 2014 (Figure 30) indicates that rivers where spawning escapement was below the CL were widely distributed throughout England and Wales. A higher proportion of rivers in northern England exceeded their CL; this is consistent with the pattern in recent years.

In 2014, additional egg deposition resulting from fish that were caught and released is estimated at about 15 million eggs (assuming 80% survival to spawning, 50% females and an average of 5,000 eggs per female). This represents about 8% of the total estimated egg deposition in England and Wales in 2014.

### Compliance with the management objective

The 'management objective' for salmon stocks in England and Wales is that they should meet or exceed their CLs in at least four years out of five, on average. Compliance with this objective takes trends in egg deposition into account, and has been calculated for all 64 principal river stocks in England and Wales for 2014 and forecast for 2019 (Table 26 and Figure 32).

The latest compliance assessment indicates that none of the principal rivers across England and Wales were classified as 'not at risk' in 2014 – having a high probability ( $p > 95\%$ ) of achieving the management objective (i.e. exceeding the CL in 4 years out of 5, on average). The same is forecast to apply in 2019. In 2014, 21 rivers (33%) were classified as 'at risk' - having a low probability ( $p < 5\%$ ) of achieving the management objective. This is the joint lowest number in the time series and represents a reduction on 2013, when 31 rivers were classified as 'at risk'. Fourteen rivers are forecast to be 'at risk' in 2019. The majority of the rivers in England and Wales in 2014 (61%) are classified as 'probably at risk' ( $5\% < p < 50\%$  of achieving the management objective); this rises to 70% in 2019. Only 4 rivers are classified as 'probably not at risk' ( $50\% \leq p < 95\%$ ). These compliance figures are summarised, separately, for rivers in England and Wales below:

## Rivers in England

Stock status category	Probability of meeting the management objective	2014		2019	
		Number of rivers	%	Number of rivers	%
<b>Not at risk</b>	>95%	0	0	0	0
<b>Probably not at risk</b>	50–95%	4	10	4	10
<b>Probably at risk</b>	5–50%	28	67	29	69
<b>At risk</b>	<5%	10	24	9	21

## Rivers in Wales

Stock status category	Probability of meeting the management objective	2014		2019	
		Number of rivers	%	Number of rivers	%
<b>Not at risk</b>	>95%	0	0	0	0
<b>Probably not at risk</b>	50–95%	0	0	1	5
<b>Probably at risk</b>	5–50%	11	50	16	73
<b>At risk</b>	<5%	11	50	5	23

For rivers in England (Figure 33a) there has been a general decrease in the proportion of rivers regarded as 'at risk' over the past 11 years, and this is predicted to continue. However, while the percentage of rivers classified as 'not at risk' was relatively stable, at about 20%, over the early part of the time series, there has been a progressive decline in the rivers in this category in the last four years with none assessed as 'not at risk' in 2014, and this is predicted to continue in 2019. There has also been a recent decline in the number of rivers classified as 'probably not at risk' with the majority of rivers (67% in 2014) falling in the 'probably at risk' category. The 2014 assessment suggests that the majority (90%) of English rivers will fall in the 'probably at risk' and 'at risk' categories in 2019.

For Wales (Figure 33b), a higher proportion of rivers have fallen in the 'at risk' category over the time series and very few rivers have been classed as 'not at risk'. In 2014, all the rivers are classified as either 'at risk' (50%) or 'probably at risk' (50%). The predicted trend suggests that 96% of rivers will fall in the same lowest two categories in 2019, albeit with the majority classed as 'probably at risk'.

The latest assessment thus indicates that the majority of salmon stocks in England and Wales remain in a depleted state. While for a number of rivers, catch derived estimates of spawner numbers may not adequately take account of factors such as fishing conditions and angling effort, the poor assessment outcome in 2014 is consistent with the low numbers of returning fish seen on many of the counted rivers (Section 7).



## Assessment of pre-fishery abundance (PFA) for England and Wales

Each year, ICES makes an assessment of the status of the salmon stocks in the Northeast Atlantic (NEAC) area as a basis for advising managers and providing catch advice for the distant water fisheries. A key part of this assessment is the estimation of the pre-fishery abundance (PFA) of all NEAC stocks, which is defined as the number of fish alive in the sea on January 1 in their first sea winter. This is split between maturing (potential 1SW) and non-maturing (potential MSW) fish. The PFA estimates for the period since 1971 provide our best interpretation of what the available catch and effort data tell us about changes in the status of the total national stock of salmon over the past four to five decades.

The estimated PFA of salmon from England and Wales has declined by about 45% from the early 1970s to the present time (Figure 34). The decrease has been somewhat greater in the non-maturing (i.e. potential MSW) component of the PFA, which has declined by a little over 50%, whilst the maturing (i.e. potential grilse) component has declined by a little over 40%. It should be noted that these trends mask conflicting changes in individual river stocks. Many rivers have experienced more serious declines but these are obscured by the very substantial improvements and recovery in others. The results also suggest that there was a marked decline in PFA around 1990, which is consistent with the general perception of a decrease in the marine survival for many stocks around the North Atlantic at about this time. [NB the model cannot provide an estimate of PFA of potential MSW fish for the most recent year].

The estimated number of salmon returning to rivers in England and Wales (prior to exploitation in homewater fisheries) shows a similar downward trend to the PFA (Figure 35), although the decrease is less marked due to the reduction in net exploitation in distant water fisheries. Thus, numbers of returning fish are estimated to have declined by about 40% between the early 1970s and the present time. For much of the period, the decline in returning MSW fish has been greater than that of the 1SW (grilse) returns, as with the PFA. However, a higher proportion of MSW fish has been observed in recent years and the percentage reduction in returning fish between the start and the end of the times series is now slightly greater for 1SW fish. The difference between the estimated numbers of returning fish and those surviving to spawn has reduced progressively over the time series and the total spawning escapement has remained reasonably consistent over the time period (Figure 35). This reflects the marked reduction in homewater net and rod fisheries, including the increasing use of catch and release. Although there was an upturn in numbers of returning fish and spawners in 2010 (Figure 35), these have both declined in subsequent years. The estimated numbers of returning fish and spawners in 2014 were both the lowest in the time series.

**Table 26. Conservation Limits (CL) and the percentage of the CL attained for the principal salmon rivers in England and Wales, 2005-2014. Current compliance against the management objective and predicted compliance in 2019 are shown in the right hand columns (see background report for details).**

EA Region/ NRW River	Accessible wetted area (ha)	CL eggs/ 100m <sup>2</sup>	CL eggs (x10 <sup>6</sup> )	Mgmt Target eggs (x10 <sup>6</sup> )	2014 egg deposition (x10 <sup>6</sup> )	Percentage of Conservation Limit attained (%) <sup>(a)</sup>										Current compliance <sup>(a)</sup> 2014	Predicted compliance <sup>(a)</sup> in 2019
						2005	2006	2007	2008	2009	2010	2011	2012	2013	2014		
<b>NE</b>																	
Coquet	144	218	3.14	5.75	3.90	462	280	266	337	185	370	325	366	228	124	PaR	PaR
Tyne <sup>(c)</sup>	542	208	11.25	22.49	28.51	360	386	245	218	214	519	518	277	436	253	PNaR	PNaR
Wear	232	250	5.80	11.51	17.71	238	204	211	236	181	383	460	321	526	305	PNaR	PNaR
Tees	620	240	14.90	16.94	0.52	20	24	33	55	14	14	21	50	23	3	AR	AR
Esk-Yorks	86	236	2.02	2.45	1.35	93	107	96	133	44	120	105	89	100	67	PaR	PaR
<b>Southern</b>																	
Test	138	246	3.40	3.85	2.45	82	76	48	107	65	60	70	68	73	72	PaR	PaR
Itchen	69	234	1.63	2.01	1.98	62	63	45	87	41	114	105	97	75	122	PaR	PNaR
<b>SW</b>																	
Avon-Hants	369	175	6.48	7.60	2.59	26	76	74	44	49	32	50	43	89	40	PaR	PaR
Stour	142	149	2.12	2.20	0.11	5	16	16	10	10	7	11	9	19	5	AR	AR
Piddle	18	177	0.31	0.40	0.08	51	71	63	95	58	102	135	50	33	27	AR	PaR
Frome	88	171	1.50	2.22	0.77	98	142	111	161	102	179	239	93	57	52	PaR	PaR
Axe	83	175	1.45	1.76	0.23	10	31	12	74	51	25	58	77	26	16	PaR	PaR
Exe	282	253	7.14	13.11	2.96	154	113	186	305	192	221	341	279	70	41	PaR	PaR
Teign	98	251	2.47	3.85	2.52	99	102	123	304	96	133	175	207	123	102	PaR	PaR
Dart	137	218	2.98	3.94	0.40	47	55	76	107	53	96	93	143	37	13	AR	AR
Avon-Devon	35	202	0.70	0.92	0.61	117	162	148	120	57	151	122	127	50	88	PaR	PaR
Erme	20	180	0.37	0.48	0.05	13	129	15	63	47	87	86	66	76	13	PaR	PaR
Yealm	11	212	0.24	0.27	0.07	54	55	39	48	56	80	64	57	49	29	AR	AR
Plym	29	188	0.55	0.68	0.19	17	78	42	71	15	54	91	43	24	35	PaR	PaR
Tavy	68	201	1.37	1.71	0.62	101	88	89	102	68	152	84	102	64	45	AR	PaR
Tamar	293	395	11.56	14.92	8.93	121	108	75	186	104	139	104	126	74	77	PaR	PaR
Lynher	29	233	0.68	0.98	0.50	144	111	180	131	149	266	104	162	150	73	PaR	PaR
Fowey	42	207	0.86	1.27	1.56	289	254	234	266	206	345	196	153	261	181	PNaR	PaR
Camel	56	176	0.98	1.92	0.63	277	309	316	301	170	462	241	142	158	65	PaR	PaR
Taw	274	211	5.78	9.60	6.02	92	133	277	221	158	134	287	199	52	104	PaR	PaR
Torrige	198	207	4.10	5.10	1.96	66	65	65	128	58	80	68	131	58	48	PaR	PaR
Lyn	27	359	0.97	1.67	0.58	118	195	154	305	90	227	291	166	85	60	PaR	PaR
<b>Midlands</b>																	
Severn	898	143	12.85	16.63	11.28	178	140	87	120	88	88	161	113	156	88	PaR	PaR
<b>NW</b>																	
Ribble	351	242	8.49	10.48	4.90	123	113	117	146	100	121	103	91	60	58	AR	AR
Wyre	67	70	0.47	0.57	0.08	13	18	30	94	48	32	45	39	15	16	AR	PaR
Lune	423	280	11.84	17.24	8.76	202	151	237	194	176	175	132	88	95	74	AR	AR
Kent	68	223	1.52	3.03	1.05	413	345	337	136	148	326	275	160	138	69	PaR	AR

Table 26. continued

Leven	46	182	0.83	1.15	1.42	81	61	114	95	42	175	109	60	119	170	PNaR
Crake	16	194	0.32	0.42	0.13	117	72	47	28	57	104	152	119	91	40	PaR
Duddon (& Lickle)	26	121	0.31	0.75	0.41	351	380	370	783	357	427	540	343	316	133	PaR
Esk	20	181	0.37	0.85	0.50	251	278	592	249	78	64	176	112	156	134	PaR
Irt	35	198	0.69	0.99	0.27	179	200	170	158	81	192	192	143	164	40	PaR
Ehen	41	230	0.94	1.76	1.46	227	213	303	335	216	513	371	220	279	155	PNaR
Calder	13	261	0.33	0.48	0.13	85	124	46	158	79	204	174	124	91	41	PaR
Derwent	213	185	3.93	7.08	3.09	357	215	354	309	211	306	271	152	144	78	AR
Eden	688	200	13.75	19.09	6.62	212	152	133	148	139	128	115	105	65	48	AR
Esk-Border <sup>(d)</sup>	306	255	7.79	10.77	4.75	147	114	120	161	111	194	189	123	66	61	PaR
<b>Wales</b>																
Wye	1,721	224	38.57	44.33	15.94	58	45	40	59	35	25	50	79	79	41	AR
Usk	407	248	10.11	14.77	9.04	124	189	226	211	97	99	137	222	122	89	PaR
Taff & Ely	146	219	3.19	3.43	0.49	9	9	22	15	14	25	34	25	30	15	AR
Ogmore	61	180	1.10	1.36	0.28	34	30	55	74	48	110	52	47	17	26	PaR
Tawe	88	211	1.85	2.36	0.38	87	97	96	98	58	114	55	34	37	21	AR
Tywi	500	226	11.30	15.33	3.89	108	129	152	146	85	178	120	78	84	34	AR
Taf	90	189	1.70	2.60	0.88	136	130	84	115	68	268	122	78	58	52	PaR
E&W Cleddau	87	179	1.55	1.81	0.54	32	29	67	89	69	61	44	55	37	35	PaR
Teifi	326	265	8.65	12.18	5.28	125	142	137	151	81	235	155	116	90	61	PaR
Rheidol	31	222	0.68	0.84	0.20	70	53	60	124	31	37	62	46	66	29	AR
Nevern	19	259	0.48	0.60	0.43	63	140	107	88	33	91	57	60	81	89	PaR
Dyfi	179	235	4.21	5.67	0.38	75	125	97	123	56	93	125	70	24	9	AR
Dysinni	31	216	0.68	0.73	0.04	6	6	8	13	13	11	0	30	18	5	PaR
Mawddach	57	242	1.37	2.10	1.16	149	43	169	148	99	236	199	199	73	84	PaR
Dwyrdd	9	201	0.19	0.38	0.14	286	353	155	357	72	52	116	44	56	73	PaR
Glaslyn	25	191	0.48	0.66	0.50	36	57	96	146	106	78	122	107	193	103	PNaR
Dwyrfawr	33	258	0.86	1.00	0.23	35	46	26	53	15	79	43	27	19	27	PaR
Seiont	21	226	0.48	0.94	0.10	375	342	229	219	99	214	127	158	49	20	AR
Ogwen	24	362	0.87	1.71	0.33	396	153	302	365	216	347	244	231	112	38	AR
Conwy	63	185	1.17	1.85	0.78	186	171	207	212	153	331	200	164	107	67	PaR
Clwyd	84	237	1.99	3.38	0.27	24	40	173	212	104	233	175	138	33	14	PaR
Dee	617	248	15.30	16.52	12.30	83	101	82	93	106	80	85	87	79	80	PaR
E & W Total			265.97	365.41	186.22											

Key to compliance assessments: **NaR** Not at risk **PNaR** Probably not at risk **PaR** Probably at risk **AR** At risk

Key: <sup>(a)</sup> Estimates include eggs contributed by rod-released fish.

<sup>(b)</sup> Basis for current and predicted compliance explained in Background Report (see text for details).

<sup>(c)</sup> Provisional salmon counts now used on the Tyne to estimate egg deposition.

<sup>(d)</sup> Prior to 1 April 2005, Border Esk egg deposition estimates were based only on English rod catch and likely to be underestimates.

Notes: Some entries in this table have been updated from that presented in previous reports as a result of river-specific refinements and corrections.

On some rivers, catch returns from fishery owners (rather than declared catches) or data from counters/traps have been used to derive estimates of egg deposition where these are considered to provide the most complete record of the returning stock.


Data for 2014 are provisional.

**Table 27. Number and percentage of salmon river stocks above their Conservation Limit (CL), between 50% and 100% of the CL, and less than 50% of the CL, 1993–2014.**


Year	>CL		50-100% CL		<50% CL	
	No.	%	No.	%	No.	%
1993	33	54	13	21	15	25
1994	41	65	14	22	8	13
1995	26	41	21	33	16	25
1996	31	49	16	25	16	25
1997	21	33	25	39	18	28
1998	30	47	23	36	11	17
1999	19	30	24	38	21	33
2000	26	41	25	39	13	20
2001 <sup>[a]</sup>	21	36	17	29	20	34
2002	27	42	20	31	17	27
2003	19	30	17	27	28	44
2004	41	64	15	23	8	13
2005	32	50	17	27	15	23
2006	38	59	14	22	12	19
2007	33	52	16	25	15	23
2008	43	67	15	23	6	9
2009	22	34	26	41	16	25
2010	39	61	16	25	9	14
2011	42	66	13	20	9	14
2012	34	53	17	27	13	20
2013	20	31	27	42	17	27
2014	12	19	22	34	30	47
Average % 1993–2014		47		30		24

Key: <sup>[a]</sup> No CL possible for 6 rivers due to impact of foot and mouth disease.


Notes: Data for 2014 are provisional.




**Figure 30. Pie charts for individual rivers for which Conservation Limits (CLs) have been set showing the % of the CLs attained in 2014.**



**Figure 31. Percentage of salmon river stocks exceeding their Conservation Limit (CL), between 50% and 100% of the CL, and less than 50% of the CL, 1993-2014.**



**Figure 32. Status of river catchments in 2014 assessed against the management objective (i.e. that the CL is met or exceeded in at least 4 years out of 5, on average).**



**Figure 33. Percentage of principal salmon rivers in each risk category, assessed against the management objective, for 2004–2014, and as predicted for 2019 for rivers in (a) England and (b) Wales.**






Figure 34. Estimated Pre Fishery Abundance (PFA) of salmon from UK (England & Wales), 1971-2014, as derived from the ICES-NEAC PFA model, 2014.




Figure 35. Estimated numbers of returning and spawning salmon for UK (England & Wales), 1971-2014, as derived from the ICES-NEAC PFA model, 2014, together with the national conservation limit (derived from the sum of river-specific CLs).

## **9. FACTORS AFFECTING STOCKS, FISHERIES AND CATCHES**

### **9.1 Management measures**

Viewed against historical data, current stock estimates and catches provide ongoing cause for concern and the conservation of salmon remains a priority. As a result, a number of measures aimed at better management of this valuable resource have been implemented or strengthened in England and Wales in recent years. The following provides a brief overview:

- The number of licences issued for nets and fixed engines in all parts of England and Wales has continued to decline as a result of measures taken to reduce levels of exploitation and the declining commercial viability of some fisheries. Overall, the number of net licences has decreased by 75% since 1971.
- The national spring salmon measures introduced in 1999 have reduced the proportion of the net catch taken before June from a 5-year average of 6.7% in the mid-1990's to 0.3%, on average, from 1999; these latter fish are all required to be released. These measures were renewed for another ten years in December 2008.
- Several net fisheries are being (or have been) phased out because they exploit migratory salmonids returning to more than one river (i.e. mixed stock fisheries). Arrangements have also been made to reduce netting effort in some fisheries by either compensating netsmen not to fish for a particular period (buy-offs), or through voluntary agreement to return salmon alive.
- In Ireland, action by the Government and fisheries authorities in 2007 imposed new restrictions on their fisheries and ended legal coastal drift netting for salmon. This change is estimated to have resulted in up to 5,000 more grilse returning to homewaters, particularly to rivers in the south and west of England and Wales.
- The national spring salmon measures have also affected rod fisheries. The proportion of the rod catch taken before June fell from a mean of 10.9% over the period 1994–1998 to an average of 5.9% for the period since 1999, and these fish are now required to be released.
- Non-statutory restrictions on methods and fishing areas imposed by fishery owners and angling associations include weekly and seasonal bag limits, and there are ongoing efforts to promote catch and release (C&R). As a consequence, the proportion of salmon released by anglers has increased steadily from 10% in 1993 to at or above 60% in the last five years (77%, provisionally, in 2014, the highest in the time series). Tracking studies suggest that, if handled appropriately, the majority (~85%) of released salmon go on to spawn successfully.

### **9.2 Other factors**

Other, non-regulatory, factors may also contribute to changes in stocks and catches, for example, the condition of returning fish, weather conditions, water quality, extreme flow events and the market prices of wild and farmed fish. Further information on these factors is provided in the background report. The following provides brief details of factors pertinent to 2014:

## **The effect of river flows on angler effort and catches**


For rod fisheries, river flow is a key factor affecting angler effort. In 2014, flows were generally below the long-term average in March and April and again for much of the summer and early autumn (July to October inclusive). River flows were particularly low in July and September (Figure 36). The summer and early autumn represents an important period for most rod fisheries and relatively low flows at this time are likely to have affected runs of fish and provided conditions that were unfavourable for angling, particularly for 1SW salmon, since these only start to return to rivers in the summer months. This would also have had an effect on fishing effort and contributed to the marked reduction in the number of days fished by anglers in 2014, which was 33% below the average of the previous five years (Table 8 and Figure 4).

Monthly rod catch data for the majority of the rivers featured in Figure 36, expressed in the same format as the flow data, as a percentage of the long-term average, are presented in Figure 37. This excludes the River Cynon, which has no catch of salmon, and includes the catch for the whole River Tyne rather than just the South Tyne tributary. The long-term average for the rod data has only been extended back as far as 1999, which is when the national measures were introduced imposing compulsory catch and release in the early part of the season. Fishing patterns are likely to have been different prior to this time. The monthly rod catch data have also been restricted to the period February to October, since for most rivers fishing seasons do not extend outside this period.


Median monthly rod catches in 2014 were well below the long-term average for all months of the year. The particularly low catches for February and March need to be treated with some caution since there is relatively little fishing at this time and this is restricted to only some rivers. However, it is evident that catches were typically poor in all months and were particularly low in July and September, coinciding with the periods of particularly low flow. It should be remembered that differing proportions of 1SW and MSW fish in the runs and the timing of the return migrations will also have an impact on catch rates, in addition to river flows. Thus, the slightly better catches in April and May will probably reflect the relatively better returns of MSW salmon, many of which return earlier in the year, while the poorer catches later in the year likely reflect the very low numbers of returning 1SW salmon (Figure 19). Taken overall, conditions for salmon angling in 2014 appear to have been poor and rod catches were exceptionally low (Figure 11).

## **First sale price of salmon**


The first sale price of salmon has potential implications for fishing effort and the economic viability of those net fisheries that target these fish. The average monthly price of wild salmon varies seasonally, reflecting both availability and the size of fish. Figure 38 provides an indication of trends in the first sale price of both wild and farmed salmon since 1978. The data (from Mawle, in prep) are provided for a single month, August, and have been adjusted by the Retail Prices Index. Further discussion on these price changes is provided in the background report. Wild salmon comprised just 0.2% of the salmon sold through Billingsgate fish market in 2014.



**Figure 36. Monthly mean river flows (cubic metres per second) in 2014 for 12 rivers (South Tyne, Itchen, Avon, Exe, Taw, Severn, Wye, Cynon, Teifi, Dee, Lune and Eden) in England & Wales, expressed as a percentage of the long-term average on each river for the same month.**



**Figure 37. Monthly rod catches in 2014 for 11 rivers (Tyne, Itchen, Avon, Exe, Taw, Severn, Wye, Teifi, Dee, Lune and Eden) in England & Wales, expressed as a percentage of the long-term average on each river for the same month.**



**Figure 38. The average price of farmed and wild Scottish salmon sold in August at Billingsgate, London, 1978 to 2014, adjusted to 2014 prices (adapted from Mawle, in prep.).**

## 10. EXISTING AND EMERGING THREATS TO SALMON POPULATIONS

Further information on the various factors impacting on salmon stocks in England and Wales is reported in the NASCO Implementation Plan and in the annual progress reports to NASCO. These reports are available at: [http://www.nasco.int/implementation\\_plans\\_cycle2.html](http://www.nasco.int/implementation_plans_cycle2.html). Some additional information is also available in the background report. The following provides a brief update on one issue:

### Red Vent Syndrome

The occurrence of salmon returning to rivers in England and Wales with swollen and/or bleeding vents has been noted since 2004. The condition, referred to as Red Vent Syndrome (RVS), has continued to be observed since this time, and has been subject to ongoing monitoring. Monitoring programmes on salmon 'index' rivers provide the most consistent measure of the incidence of RVS. Since 2007, this consistency has been improved through the introduction of a system whereby symptoms have been classified according to their apparent severity (with samplers referring to a set of standard photographs and descriptions to assist their judgement). Time series of RVS incidence in returning fish are presented in Table 28 for the Rivers Tyne, Tamar, Dee, Lune and Caldew (a tributary of the River Eden). However, no sampling was possible at two of these sites in 2014 and sampling effort was substantially reduced at another.

Fish affected by RVS show a degree of recovery in freshwater and appear to be able to spawn successfully.

**Table 28. Percentage of returning salmon showing signs of Red Vent Syndrome in monitored rivers in England and Wales, 2004–2014**

River	Tyne #	Tamar	Dee	Lune	Caldew #
Region/NRW	NE	SW	N. Wales	NW	NW
Sample source	Upper river broodstock	Lower river trap	Lower river trap	Lower river trap	Sub-catchment trap
% incidence of RVS in returning fish					
2004			0.4		
2005			3.2	0	
2006			9.2	1.4	
2007	1.4	60.2	29.9	23.1	5.3 <sup>[a]</sup>
2008	0.8	45.3	20.9	24.7	0.3 <sup>[a]</sup>
2009	3.4	41.5	28.2	21.2	10.2
2010	5.3	57.1	23.7	18.8	5.1
2011	3.8	45.6	10.9	16.3	6.4
2012	5.2	26.1	13.2	0 <sup>[a]</sup>	6.1
2013	10.1	44.5 #	20.5	41.6	0.8 <sup>[a]</sup>
2014	7.5	n/a <sup>[b]</sup>	25.3	9.5 #	n/a <sup>[b]</sup>

Note: Except where indicated (#), these estimates are based on fish sampled over a common (June–October) period and have been weighted according to monthly run totals. Three of the traps (not the Caldew) are located at or close to head-of-tide.

<sup>[a]</sup> Minimum values.

<sup>[b]</sup> Trap not operational in 2014.

## 11. REFERENCES

- Environment Agency, 2014. Salmonid and freshwater fisheries statistics for England and Wales, 2013, 42 pp. <https://www.gov.uk/government/collections/salmonid-and-freshwater-fisheries-statistics>
- ICES. 2014. Report of the Working Group on North Atlantic Salmon. ICES CM 2014/ACOM:09, 431 pp.
- Mawle, G.W. (in prep.). The impact of negative publicity for farmed salmon on the price of wild salmon in the United Kingdom.
- Potter, E.C.E., Crozier, W.W., Schön, P-J., Nicholson, M.D., Prévost, E., Erkinaro, J., Gudbergsson, G., Karlsson, L., Hansen, L.P., Maclean, J.C., Ó Maoiléidigh, N. and Prusov S. 2004. Estimating and forecasting pre-fishery abundance of Atlantic salmon (*Salmo salar* L.) in the north-east Atlantic for the management of mixed stock fisheries. ICES Journal of Marine Science 61: 1359-1369.

# **ANNEX 1. NASCO's request for scientific advice from ICES in 2015**

## **1. With respect to Atlantic salmon in the North Atlantic area:**

- 1.1 provide an overview of salmon catches and landings by country, including unreported catches and catch and release, and production of farmed and ranched Atlantic salmon in 2014<sup>1</sup>;
- 1.2 report on significant new or emerging threats to, or opportunities for, salmon conservation and management<sup>2</sup>;
- 1.3 provide a review of examples of successes and failures in wild salmon restoration and rehabilitation and develop a classification of activities which could be recommended under various conditions or threats to the persistence of populations<sup>3</sup>;
- 1.4 provide a compilation of tag releases by country in 2014;
- 1.5 identify relevant data deficiencies, monitoring needs and research requirements.

## **2. With respect to Atlantic salmon in the North-East Atlantic Commission area:**

- 2.1 describe the key events of the 2014 fisheries<sup>4</sup>;
- 2.2 review and report on the development of age-specific stock conservation limits;
- 2.3 describe the status of the stocks;
- 2.4 provide catch options or alternative management advice for 2015/16-2017/18 fishing seasons, with an assessment of risks relative to the objective of exceeding stock conservation limits, or pre-defined NASCO Management Objectives, and advise on the implications of these options for stock rebuilding<sup>5</sup>;
- 2.5 advise on options for taking into account the recent genetic analysis that suggests there was a significant contribution of North American origin stocks to historic mixed-stock fisheries in Faroese waters for the provision of catch advice<sup>6</sup>;
- 2.6 update the Framework of Indicators used to identify any significant change in the previously provided multi-annual management advice; and
- 2.7 advise on what data would enhance the development of the catch options.

## **3. With respect to Atlantic salmon in the North American Commission area:**

- 3.1 describe the key events of the 2014 fisheries (including the fishery at St Pierre and Miquelon)<sup>4</sup>;
- 3.2 update age-specific stock conservation limits based on new information as available;
- 3.3 describe the status of the stocks;
- 3.4 provide catch options or alternative management advice for 2015-2018 with an assessment of risks relative to the objective of exceeding stock conservation limits, or pre-defined NASCO Management Objectives, and advise on the implications of these options for stock rebuilding<sup>5</sup>;
- 3.5 update the Framework of Indicators used to identify any significant change in the previously provided multi-annual management advice;



- 3.6 considering the available contemporary data on stock origin of salmon in the Labrador fisheries, estimate the catches by stock origin and describe their spatial and temporal distribution; and
- 3.7 considering the available contemporary data on stock origin of salmon in the Saint-Pierre et Miquelon fishery, estimate the catches by stock origin and describe their spatial and temporal distribution.

**4. With respect to Atlantic salmon in the West Greenland Commission area:**

- 4.1 describe the key events of the 2014 fisheries<sup>4</sup>;
- 4.2 describe the status of the stocks<sup>7</sup>;
- 4.3 provide catch options or alternative management advice for 2015-2017 with an assessment of risk relative to the objective of exceeding stock conservation limits, or pre-defined NASCO Management Objectives, and advise on the implications of these options for stock rebuilding<sup>5</sup>;
- 4.4 update the Framework of Indicators used to identify any significant change in the previously provided multi-annual management advice; and
- 4.5 considering the available contemporary data on stock origin of salmon in the West Greenland fishery, estimate the catches by stock origin and describe their spatial and temporal distribution.

**Notes:**

- 1. *With regard to question 1.1, for the estimates of unreported catch the information provided should, where possible, indicate the location of the unreported catch in the following categories: in-river; estuarine; and coastal. Numbers of salmon caught and released in recreational fisheries should be provided.*
- 2. *With regard to question 1.2, ICES is requested to include reports on any significant advances in understanding of the biology of Atlantic salmon that is pertinent to NASCO, including information on any new research into the migration and distribution of salmon at sea and the potential implications of climate change for salmon management.*
- 3. *With regards to question 1.3, NASCO is particularly interested in case studies highlighting successes and failures of various restoration efforts employed across the North Atlantic by all Parties/jurisdictions and the metrics used for evaluating success or failure.*
- 4. *In the responses to questions 2.1, 3.1 and 4.1, ICES is asked to provide details of catch, gear, effort, composition and origin of the catch and rates of exploitation. For homewater fisheries, the information provided should indicate the location of the catch in the following categories: in-river; estuarine; and coastal. Information on any other sources of fishing mortality for salmon is also requested.*
- 5. *In response to questions 2.4, 3.4 and 4.3, provide a detailed explanation and critical examination of any changes to the models used to provide catch advice and report on any developments in relation to incorporating environmental variables in these models.*

6. *In response to question 2.5, this should include consideration of the implications of the new genetic results with regard to the factors previously identified by ICES as requiring management decisions for the finalisation of the risk framework for the provision of catch advice for the Faroes fishery (i.e. annual or seasonal catch advice, sharing agreement, choice of management units to consider and specified management objectives).*
7. *In response to question 4.2, ICES is requested to provide a brief summary of the status of North American and North-East Atlantic salmon stocks. The detailed information on the status of these stocks should be provided in response to questions 2.3 and 3.3.*

## ANNEX 2. Net Limitation Orders applying to salmon net fisheries in England & Wales

EA Region / NRW	Area	Net Limitation Order	End date	Welsh rivers in Wales 'all areas' NLO	NLO licence provision	
					Type	Number
Anglian	Coastal	Anglian Coast 2005	2015		Drift net & non-drift net	0
North East	Coastal	North East Coast 2012	2022		T and J nets	0
					Drift net - Northumbria and Yorkshire	0
North West	North	River Lune Estuary 2010	2020		Drift	7
	North	River Lune Estuary 2010	2020		Haaf	12
	North	River Ribble Estuary 2006	2017		Drift (hang or whammel) nets	2
	North	River Kent Estuary 2013	2023		Lave net	6
	North	River Leven Estuary 2013	2023		Lave net	2
	North	Solway Firth 2007	2017		Heave or Haaf net	105
Southern	Solent & S Downs	Southern Region 2008	2018		Seine	1
South West	Cornwall	River Camel 2013	2018		Draft, seine, drift or hang net	6
	South Wessex	Christchurch Harbour 2012 (Hants Avon & Stour)	2022		Draft or seine net	0
	South Wessex	Poole Harbour 2012 (Piddle & Frome)	2017		Seine net	1
	Devon	River Dart 2005	2015		Draft or seine net	3
	Devon	Exe Estuary 2011	2021		Draft nets	3
	Cornwall	River Fowey 2007	2018		Draft or seine net	1
	Cornwall	River Lynher 2014	2024		Draft or seine net	0
	Cornwall	River Tamar 2014	2024		Draft or seine net	0
	Cornwall	River Tavy 2014	2024		Draft or seine net	0
	Cornwall	Rivers Taw and Torridge 2012	2022		Draft or seine net	1
Midlands		River Severn			Draft or seine net	3
					Lave net	15
Wales	All areas	Wales 2009	2017	Nevern	Draft or seine net	1
				Taf	Coracle net	1
				Taf	Wade net	1
				Dyfi	Draft or seine net	3
				Dysynni	Draft or seine net	1
				Glaslyn & Dwyrdd	Draft or seine net	0
				Mawddach	Draft or seine net	3
				Conwy	Draft or seine net	3
				Cleddau	Compass nets	6
				Teifi	Coracle net	12
				Teifi	Draft or seine net	3
				Tywi	Draft or seine net	3
				Tywi	Coracle net	8
				Wales	North	River Dee 2004
Trammel nets	0					

Notes: Table does not include Historical Installation fisheries which operate under Certificates of Privilege or the private lave net fishery on the River Wye.

Net Limitation Orders for the seine and lave net fisheries on the River Severn (Midlands Region) are in preparation. Some fisheries are also subject to seasonal catch limits.

## ANNEX 3. Byelaws applying to salmon rod fisheries in England and Wales

EA Region / NRW	River	Salmon Season (inclusive dates)	*Method Restrictions	*Bag limits/Catch and Release etc.	Effective from (date); expires (date)
NE	Aln	1.2 -31.10			
	Coquet	1.2 -31.10			
	Tyne	1.2 -31.10			
	Wear	1.2 -31.10			
	Tees	1.2 -31.10			
	Esk (Yorks.)	6.4 -31.10			
	Ouse (Yorks.)	6.4 -31.10			
Anglian	Region	1.3 -28.9			
Thames	Thames	1.4 -30.9		2 salmon bag limit a day	
SW	Avon (Hants.)	1.2 -31.8	Artificial fly only before 15/5		
	Piddle	1.3 -31.8	Artificial fly only before 15/5		
	Frome	1.3 -31.8	Artificial fly only before 15/5		
	Axe	15.3 -31.10	No shrimp, prawn, worm or maggot. Fly only after 31/7 below Axbridge		
	Exe	14.2 -30.9 (trial extension to 14.10)	No worm or maggot	Fly only and mandatory catch and release during trial extension period.	
	Teign	1.2 -30.9	No worm or maggot before 1/6		
	Dart	1.2 -30.9	No worm or maggot. No shrimp/prawn etc. below Staverton Bridge.		
	Avon (Devon)	15.4 -30.11	No worm or maggot		
	Plym	1.4 -15.12	No worm, maggot, shrimp or prawn after 31/8		
	Tavy	1.3 -14.10	No worm, maggot, shrimp or prawn after 31/8		
	Tamar	1.3 -14.10	No worm, maggot, shrimp or prawn after 31/8		
	Lynher	1.3 -14.10	No worm, maggot, shrimp or prawn after 31/8		
	Fowey	1.4 -15.12			
	Camel	1.4 -15.12			
	Taw	1.3 -30.9	No shrimp, prawn, worm or maggot. Fly only 1/4 to 31/5	Numbers for Taw, Torridge in brackets: 2 (2) salmon a day, 3 (2) a week and 10 (7) a season, (2 salmon limit before June 1st) & return of all salmon > 70 cm after Aug 1st.	
	Torridge	1.3 -30.9	No shrimp, prawn, worm or maggot. Fly only 1/4 to 31/5		
Lyn	1.2 -31.10	No worm or maggot before 1/6			
Yealm	1.4 -15.12	No worm, maggot, shrimp or prawn after 31/8			
Midlands	Severn	1.2 -7.10	No float fishing with lure or bait		
Wales	Wye	3.3 -17.10 (a)	Fly only 1.9 to 17.10. No bait all season	Mandatory C&R all season	Commenced June 2012; expires 2021
	Usk	3.3 -17.10	Fly only 3.3 - 1.6. Fly & Spin 15.9 - 17.10		
	Taff & Ely	20.3 -17.10	Fly & Spin 20.3 -15.4 & 1.10 -17.10	Mandatory C&R all season	Commenced June 2012; expires 2018
	Ogmore	20.3 -17.10	Fly & Spin 20.3 -15.4 & 1.10 -17.10		

<b>EA Region / NRW</b>	<b>River</b>	<b>Salmon Season (inclusive dates)</b>	<b>*Method Restrictions</b>	<b>*Bag limits/Catch and Release etc.</b>	<b>Effective from (date); expires (date)</b>
Wales (cont)	Afan	20.3 -17.10	Fly & Spin 20.3 -15.4 & 1.10 -17.10		
	Neath	20.3 -17.10	Fly & Spin 20.3 -15.4 & 1.10 -17.10		
	Tawe	20.3 -17.10	Fly & Spin 20.3 -15.4 & 1.10 -17.10		
	Loughor	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Tywi	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon. C&R 8.10 to 17.10	
	Taf	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon. C&R 8.10 to 17.10	
	E+W.Cleddau	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon. C&R 8.10 to 17.10	
	Nevern	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon	
	Teifi	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon	
	Aeron	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon	
	Ystwyth	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon	
	Rheidol	1.4 -17.10	Fly & Spin 7.10 -17.10	Daily bag of 2 salmon & 4 sea trout, weekly bag of 5 salmon	
	Dyfi	20.3 -17.10 (some sections to 31.10)	Fly & Spin 20.3 -15.4 & 7.10 -31.10	Catch & Release salmon and sea trout 18.10 to 31.10	
	Dysynni	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10	Catch & Release salmon and sea trout 18.10 to 31.10	
	Mawddach	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Artro	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Dwyrhyd	20.3 -17.10 (some sections to 31.10)	Fly & Spin 20.3 -15.4 & 7.10 -31.10	Catch & Release salmon and sea trout 18.10 to 31.10	
	Glaslyn	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Dwyfawr	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Llyfni	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Gwyrfai	20.3 -17.10	Fly & Spin 20.3 -15.4 & 7.10 -17.10		
	Seiont	20.3 -15.11	Fly & Spin 20.3 -15.4 & 7.10 -15.11	Catch & Release salmon and sea trout 18.10 to 15.11	
	Ogwen	20.3 -17.10 (some sections to 31.10)	Fly & Spin 20.3 -15.4 & 7.10 -31.10	Catch & Release salmon and sea trout 18.10 to 31.10	
	Conwy	20.3 -17.10 (some sections to 31.10)	Fly & Spin 20.3 -15.4 & 7.10 -31.10	Catch & Release salmon and sea trout 18.10 to 31.10	
	Clwyd	20.3 -17.10	Fly & Spin 20.3 - 1.6, Fly only 1.10 - 17.10		
	Dee	3.3 -17.10	Fly only 3.3 - 1.6, Fly & Spin 1.10 - 17.10		

<b>EA Region / NRW</b>	<b>River</b>	<b>Salmon Season (inclusive dates)</b>	<b>*Method Restrictions</b>	<b>*Bag limits/Catch and Release etc.</b>	<b>Effective from (date); expires (date)</b>
NW	Ribble	1.2 -31.10		Byelaw (NW-28/02/07) no more than two salmon may be killed between 16.6 and 31.10	28.02.2007 – 27.02.2017
	Wyre	1.2 -31.10			
	Lune	1.2 -31.10		Byelaw (NW-14/02/00) no more than four salmon may be killed during the season.	26.11.2009 – 26.11.2019
	Kent	1.2 -31.10			
	Leven	1.2 -31.10		Byelaw (NW-08/05/03) requiring release of all salmon after capture.	09.05.2003 – 07.05.2013
	Crake	1.2 -31.10		Byelaw (NW-08/05/03) requiring release of all salmon after capture.	09.05.2003 – 07.05.2013
	Duddon	1.2 -31.10			
	Esk (Cumb.)	1.2 -31.10			
	Irt	1.2 -31.10			
	Calder	1.2 -31.10			
	Ehen	1.2 -31.10			
	Derwent	1.2 -31.10			
	Ellen	1.2 -31.10			
	Eden	15.1 -14.10		Byelaw (NW-13.11.07) no more than two salmon may be killed between 16.6 and 31.10	13.11.2007 – 12.11.2017
				Byelaw (NW-13.11.07) prohibits retention of female salmon 10.9 to 31.10	13.11.2007 – 12.11.2017
	Esk (Border)	1.2 -31.10		Byelaw (NW-13.11.07) no more than two salmon may be killed between 16.6 and 31.10	13.11.2007 – 12.11.2017
				Byelaw (NW-13.11.07) prohibits retention of female salmon 10.9 to 31.10	13.11.2007 – 12.11.2017
Others	1.2 -31.10 <sup>(b)</sup>				

Notes: <sup>(a)</sup> Season 3.3 to 25.10 Rivers Irfon, Ithon and main River Wye upstream of Llanwrthwl Bridge

<sup>(b)</sup> Rivers Annas, Bleng, Mite.

\* National spring salmon byelaws apply.

Natural Resources Wales – variations apply to Anglesey and the Lleyn Peninsula (check local byelaws).

Always check local byelaws before fishing.





Front cover images (clockwise from top left)

- 1 – Leaping salmon
- 2 – Drift netting
- 3 – Salmon being released by angler
- 4 – Haaf netsman

Photographs courtesy of the Environment Agency

Centre for Environment, Fisheries & Aquaculture Science  
Lowestoft Laboratory  
Pakefield Road  
Lowestoft  
Suffolk  
NR33 0HT  
[www.cefasc.co.uk](http://www.cefasc.co.uk)

Tel: 01502 562244  
Twitter: @cefascgovuk

Environment Agency  
National Customer Contact Centre  
PO Box 544  
Rotherham  
S60 1BY  
[www.Gov.uk](http://www.Gov.uk)

Tel: 03708 506 506 (Mon-Fri, 8am – 6pm)  
Email: [enquiries@environment-agency.gov.uk](mailto:enquiries@environment-agency.gov.uk)  
Facebook: <https://www.facebook.com/environmentagency>  
Twitter: <https://twitter.com/envagency>

Cyfoeth Naturiol Cymru /Natural Resources Wales  
Cambria House  
29 Newport Road  
Cardiff  
CF24 0TP.  
[www.naturalresourceswales.gov.uk](http://www.naturalresourceswales.gov.uk)

Tel: Customer contact centre 0300 065 3000  
Email: [enquiries@naturalresourceswales.gov.uk](mailto:enquiries@naturalresourceswales.gov.uk)  
Facebook: [www.facebook.com/NatResWales](http://www.facebook.com/NatResWales)  
Twitter: @NatResWales

