



Local Tobacco Control Profiles Indicators at a glance November 2016

Notes:

- Value cells are shaded **red**, **amber** or **green** where the England value can be benchmarked against a goal.
- In the change columns, prev refers to the change in England value compared to the previous data point; base refers to the change in England value compared to the baseline value.
- Changes are only shown if they are statistically significant. An upwards arrow (↑, ↑ or ↑) represents a significant increase in the indicator value, a downwards arrow (↓, ↓ or ↓) represents a significant decrease. A sideways arrow (↔) is displayed if there has been no significant change.
- The arrows are coloured **green** and **red** for those indicators where a change can be described as improving or worsening respectively.

Domain: smoking prevalence in adults				
Indicator	Period	Value	Unit	Change from previous
Smoking Prevalence in adults - current smokers (APS)	2015	16.9	%	↓
Smoking Prevalence in adults – ex-smokers (APS)	2015	34.5	%	↔
Smoking Prevalence in adults - never smoked (APS)	2015	48.6	%	↑
Smoking prevalence in adults in routine and manual occupations - current smokers (APS)	2015	26.5	%	↓
Smoking prevalence in adults in routine and manual occupations – ex-smokers (APS)	2015	30.4	%	↔
Smoking prevalence in adults in routine and manual occupations - never smoked (APS)	2015	43.1	%	↑
Smoking Prevalence in adults - current smokers (IHS)	2014	18.0	%	↓
Smoking Prevalence in adults - ex-smokers (IHS)	2014	33.9	%	↑
Smoking Prevalence in adults - never smoked (IHS)	2014	48.1	%	↔
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	28.0	%	↔
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014	30.8	%	↔
Smoking prevalence in adults in routine and manual occupations - never smoked (IHS)	2014	41.2	%	↔
Smoking prevalence in adults - current smokers (GPPS)	2013/14	17.1	%	
Smoking prevalence in adults – ex-smokers (GPPS)	2013/14	27.6	%	
Smoking prevalence in adults - never smoked (GPPS)	2013/14	55.3	%	
Smoking prevalence in adults - current smokers (QOF)	2014/15	18.6	%	↓
Smoking prevalence in adults with serious mental illness (SMI)	2014/15	40.5	%	
Smoking status at time of delivery	2015/16	10.7	%	↓
Domain: smoking prevalence in young people				
Indicator	Period	Value	Unit	Change from previous
Smoking prevalence age 15 years - regular smokers (SDD survey)	2014	8	%	↔
Smoking prevalence age 15 years - occasional smokers (SDD survey)	2014	5	%	↓
Smoking prevalence at age 15 - current smokers (WAY survey)	2014/15	8.2	%	

Smoking prevalence at age 15 - regular smokers (WAY survey)	2014/15	5.5	%	
Smoking prevalence at age 15 - occasional smokers (WAY survey)	2014/15	2.7	%	
Use of e-cigarettes at age 15 years (WAY survey)	2014/15	18.4	%	
Use of other tobacco products at age 15 years (WAY survey)	2014/15	15.2	%	
Smoking prevalence at ages 11-15 - regular smokers (modelled estimates)	2009 - 12	3.1	%	
Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates)	2009 - 12	1.4	%	
Smoking prevalence at age 15 - regular smokers (modelled estimates)	2009 - 12	8.7	%	
Smoking prevalence at age 15 - occasional smokers (modelled estimates)	2009 - 12	3.9	%	
Smoking prevalence at ages 16-17 - regular smokers (modelled estimates)	2009 - 12	14.7	%	
Smoking prevalence at ages 16-17 - occasional smokers (modelled estimates)	2009 - 12	5.8	%	
Domain: smoking related mortality				
Indicator	Period	Value	Unit	Change from previous
Smoking attributable mortality	2012-14	274.8	DSR per 100,000	↓
Deaths from lung cancer	2013-15	58.7	DSR per 100,000	↓
Deaths from chronic obstructive pulmonary disease	2013-15	52.6	DSR per 100,000	↑
Smoking attributable deaths from heart disease	2012-14	29.7	DSR per 100,000	↓
Smoking attributable deaths from stroke	2012-14	9.3	DSR per 100,000	↔
Potential years of life lost due to smoking related illness	2012-14	1407.3	DSR per 100,000	
Domain: smoking related ill health				
Indicator	Period	Value	Unit	Change from previous
Premature births (<37 weeks gestation)	2012-14	77.6	Crude rate per 1,000	↑
Low birth weight of term babies	2014	2.9	%	↔
Hospital admissions for asthma (under 19 years)	2014/15	216.1	DSR per 100,000	↑
Smoking attributable hospital admissions	2014/15	1671.2	DSR per 100,000	↑
Cost per capita of smoking attributable hospital admissions	2011/12	38.0	£ per capita	
Emergency Hospital admissions for COPD	2014/15	415.0	DSR per 100,000	↑
Lung cancer registrations	2012-14	79.8	DSR per 100,000	↔
Oral cancer registrations	2012-14	14.2	DSR per 100,000	↔
Domain: impact of smoking				
Indicator	Period	Value	Unit	Change from previous
Indicative tobacco sales figures (£ millions)	2013	15446.1	£	
Illicit tobacco: tax gap (£ millions)	2014/15	1200	£	↔

Illicit tobacco: share of cigarette sales (%)	2014/15	10	%	↔
Fatalities from accidental fires ignited by smoking materials	2014/15	41	%	↔
Accidental fires ignited by smoking related materials	2014/15	7	%	↓
Attitudes to smoking in 15 year olds - 'smoking causes harm to others' (WAY survey)	2014/15	91	%	
Domain: smoking quitters				
Indicator	Period	Value	Unit	Change from previous
No. setting a quit date	2015/16	5091.8	Crude rate per 100,000	↓
Successful quitters at 4 weeks	2015/16	2598.0	Crude rate per 100,000	↓
Successful quitters (CO validated) at 4 weeks	2015/16	1854.2	Crude rate per 100,000	↓
Completeness of NS-SEC recording by Stop Smoking Services	2015/16	91.1	%	↑
Cost per quitter	2015/16	479.0	£ per capita	