

A1a.4 Fish and Shellfish

A1a.4.1 Introduction

This section describes the distribution and ecology of fish and commercially important shellfish species in UK waters, with reference, where information is available, to spawning and nursery grounds. The section draws on previous fishery studies commissioned for the SEA programme and a range of other technical reports and scientific publications. Information on the seasonal distribution of commercially important fish and shellfish is available from several sources, with the most reliable being the routine research vessel surveys undertaken by European research laboratories, co-ordinated by the International Council for the Exploration of the Sea (ICES). These annual surveys target major commercial species but they also record information on the distribution and abundance of the non-target components of the catch. The recently published Fish Atlas of the Celtic Sea, North Sea and Baltic Sea (Heessen *et al.* 2015) synthesises results from these international research surveys, incorporating data from over 70,000 hauls carried out between 1977-2013.

A1a.4.2 UK context

Over 330 species of fish have been recorded on the UK continental shelf and the deep water regions which lie to the north and west of Scotland are home to many more. The following sections give a brief overview of a selection of abundant and widespread species in UK waters, grouped into pelagic (open water), demersal (bottom dwelling), elasmobranch (sharks and rays), diadromous (migrate between the sea and freshwater to spawn) and shellfish species.

A1a.4.2.1 Pelagic species

Mackerel (*Scomber scombrus*) are widely distributed around the north-east Atlantic where they tend to shoal in large schools in waters down to 1,000m deep. They usually grow to between 35 and 45cm long although they can reach 60cm. They feed on pelagic crustaceans and other zooplankton and small fish. Mackerel are fast growing and are sexually mature by three years of age (Gordon 2006). They are long-lived; individuals may live for 20-22 years (Navarro *et al.* 2012). Spawning is pelagic and the spawning season prolonged. Eggs are shed in large batches (a 200g female may produce 211,000 eggs per batch (von Damme & Thorsen 2014)) and studies of spawning patterns reveal there to be two main mackerel stocks in UK waters: a western stock and a North Sea stock (Coull *et al.* 1998). Mixing between the two stocks has increased since the 1980s, through migration of western fish into the southern and northern North Sea (Jansen & Gialson 2013).

Atlantic herring (*Clupea harengus*) are widespread throughout the north-east Atlantic, although they reach the southern limit of their range just south of the UK (Dickey-Collas *et al.* 2015). A number of different spawning grounds exist around the UK, with spawning occurring at different times (although the majority takes place in late summer). Spawning usually takes place at depths of between 15-40m, when herring deposit their sticky eggs on coarse sand and gravel. Spawning may take place in coastal waters or open sea, as long as waters are well-mixed, and temperature and salinity tolerance are wide (Pörtner & Peck 2010). The dependency of herring on these specific substrates makes the species potentially susceptible to disturbance at these sites and largely limits herring distribution to the shelf region. Shoals congregate and spawn simultaneously at traditional spawning grounds, resulting in the formation of an, "egg carpet," which may extend over an area of one hectare (Blaxter & Hunter 1982). Each female produces a single batch of eggs every year from the age of about 3 years, although the number, size and weight of eggs will vary between spawning populations. Eggs hatch after 1-3 weeks, depending on the water temperature, and pelagic larvae will drift towards important nursery grounds (Dickey-Collas *et al.* 2015). As the length of the larval development is dependent on food

availability, herring spawned later in the year when food availability has declined may not undergo metamorphosis until the following spring. These larvae are likely to drift long distances in this time (Fässler *et al.* 2011). Young herring occur in dense shoals in inshore waters, and are often found in mixed shoals with varying proportions of sprats (*Sprattus sprattus*) (Dickey-Collas *et al.* 2015).

Sprats are widespread along Atlantic coasts and are usually found in shallow water close to shore, where they can tolerate low salinities. They tend to range in length from 8-12cm and are a short-lived species, the abundance of which is heavily dependent on the strength of the recruiting year class (Gordon 2006). They feed on a range of planktonic crustaceans. Spawning mainly occurs in the summer months, near the coast or up to 100km out to sea, at depths of 10-20m (Gordon 2006). During spawning, 6,000-14,000 pelagic eggs can be produced and this fecundity, along with the rapid life-cycle, means sprat have a very short population doubling time of less than 15 months. Juvenile sprat are generally found in dense schools in shallow, coastal waters (Dickey-Collas *et al.* 2015).

Horse mackerel (*Trachurus trachurus*) is a schooling fish, particularly abundant to the south and west of the UK. Horse mackerel around the UK are split into two stocks; a western stock and a North Sea stock. The North Sea stock spends most of the year in the central North Sea, Skagerrak and Kattegat, but migrates to the southern North Sea in the summer to spawn (FishBase website¹). Strong influxes of Atlantic water leads to periods of increased abundance in the northern North Sea (Iversen *et al.* 2002). Adults form large shoals in coastal areas with sandy sediments, where they feed on fish, cephalopods and crustaceans. Horse mackerel is notable for an irregular pattern of recruitment, with extremely strong year classes being produced at lengthy intervals. The most recent strong year classes were 1982 and 2001, with the 1982 class (three times larger than the 2001 class) strong enough to dominate the stock for years (Ellis 2015a).

Argentines are deep water salmoniformes found over muddy sediment at depths of between 55-550m, although most commonly at depths greater than 200m (Wheeler 1978, Heessen 2015a). Two species of argentine, the greater (*Argentina silus*) and lesser (*Argentina sphyraena*) are present in the north-east Atlantic. The greater argentine is larger and tends to be found in deeper water, closer to the edge of the continental shelf than the lesser argentine. They feed on bottom living worms and molluscs and also predate on pelagic fish, crustaceans and squid at night. They spawn between March and September, producing pelagic eggs and larvae.

Blue whiting (*Micromesistius poutassou*) is a meso-pelagic species, usually found in shoals 30-400m from the surface in water between 150-3,000m deep. Shoals move towards the surface at night. They are widely distributed around the north-east Atlantic, typically reach lengths of 25-30cm and live for 5-7 years (Gordon 2006). They feed primarily on small crustaceans such as euphausiids.

A1a.4.2.2 Demersal species

Gadoids

Gadoids are important components of the fish community of the north-east Atlantic, with 12 (of 31 known) species found in the region (Nelson 2006). The Atlantic cod (*Gadus morhua*) can be found from the shoreline down to depths of 600m and is widely distributed around European

¹ <http://www.fishbase.org/summary/1365>

coasts. It can reach lengths of 190cm, but more commonly of between 50-80cm and, although heavily exploited, has the potential to live to 15 years or older (Hislop *et al.* 2015). Cod are omnivores, feeding on a variety of invertebrates and fish. Sexual maturity is reached between 4-5 years and spawning occurs over the continental shelf between January and April. Cod show a preference to spawn in waters with temperatures between 5-7°C and high salinities, over coarse sand with a low tidal flow (González-Irusta & Wright 2015). Larval abundance peaks at fronts and juveniles remain pelagic until they reach a length of 5-7cm. Adult cod aggregate in loose shoals and generally remain within the continental shelf area (Hislop *et al.* 2015).

The haddock (*Melanogrammus aeglefinus*) is found around north-east Atlantic coasts, over rock, sand or gravel bottoms at 10-450m depth, usually between 80-200m (Albert 1994). It can reach 50-75cm long and feeds on small benthic invertebrates and fish. It is known to reach ages of up to 20 years. There is some evidence of a winter migration of adult haddock from the North Sea to north-western Scotland (Hislop *et al.* 2015). Haddock is a batch spawner, with the season typically extending from February to May, with the north-east coast of Scotland, the seas around Shetland, the north-west coast of Scotland, the Minch and the north-west Irish Sea and Celtic Sea usual spawning areas. Eggs are pelagic, and modelling suggests that a significant number of the eggs spawned off the west coast are transported into the North Sea (Heath & Gallego 1997).

Whiting (*Merlangius merlangus*) are widespread around European coasts at depths of 10-200m over sandy or muddy ground. They typically grow to 30-40cm in length and may reach 20 years of age, although 7 or 8 is more common (Gordon 2006). Their diet comprises mainly crustaceans and fish, with a greater proportion of fish as they get older. Spawning can take place as early as January in the southernmost areas of its distribution and as late as July in more northerly areas. Whiting spend their first 2-3 months near the surface, often associating with *Cyanea* jellyfish blooms (Hay *et al.* 1990), after which they adopt a demersal way of life.

Saithe (*Pollachius virens*) are most abundant at depths of between 125-200m around north-east Atlantic coastlines, usually entering coastal waters in spring and migrating back to deeper sea in winter (Hislop *et al.* 2015). They grow to 60-90cm and have a diet of fish and small crustaceans. Saithe reach maturity between 4-6 years and individuals aged 25 years have been reported (Gordon 2006). They spawn in winter and spring, later in the year for populations further north. Pollack (*Pollachius pollachius*) live inshore over rocky ground at depths of up to 200m, around north-east Atlantic coasts. They can grow to between 60 and 80cm, although individuals of up to 130cm have been found. The pollack feeds primarily on fish, with cephalopods and crustaceans also forming part of its diet. Juvenile shoals are common inshore but the adults shoal only during the spawning period, which takes place in winter and spring at about 100m depth (Whitehead *et al.* 1986).

A number of smaller gadoid species such as poor cod (*Trisopterus minutus*), Norway pout (*Trisopterus esmarkii*) and bib (*Trisopterus luscus*) can be very abundant in places and may be ecologically important as prey for other species.

Flatfish

Plaice (*Pleuronectes platessa*) live to depths of 200m, mainly on soft sediments. Spawning occurs in water temperatures of approximately 6°C to produce pelagic eggs. Larvae move to coasts, and sandy beaches and estuarine regions act as nursery grounds, before older individuals venture out to deeper waters (Gordon 2006). Plaice live on mixed substrates at depths up to 200m (although generally in much shallower waters), with older individuals generally found in deeper water (Whitehead *et al.* 1986). Plaice have a complicated life cycle, with each life stage having a specific set of habitat requirements. Larvae and juveniles rely on

transport by currents to move them from spawning grounds to nursery areas, a habit that adults retain by making use of tidal transport during seasonal migrations between spawning and feeding grounds (Goldsmith *et al.* 2015).

Dab (*Limanda limanda*) and long rough dab (*Hippoglossoides platessoides*) are spring and summer spawners which mature at 2-3 years to produce pelagic eggs and larvae. Dab larvae were the most abundant to be found in the North Sea and Irish Sea during ichthyoplankton surveys. Dab are typically found in shallower water, where they feed on small benthic invertebrates (Amara *et al.* 1998). The long rough dab tends to be found in deeper waters, up to 500m, over muddy substrates. Other important flatfish include the lemon sole (*Microstomus kitt*), mainly on coarser sediments to 200m, and the sole (*Solea solea*), especially on finer sandy and muddy seabeds to around 120m, including estuarine areas. Both are widespread in British waters, but sole is much more abundant in the southern half of the British Isles than in the north as it is generally confined to warmer waters (Rijnsdorp *et al.* 2015).

Other species

Two similar species of monkfish, white-bellied (*Lophius piscatorius*) and black-bellied (*L. budegassa*) are typically found in northern UK waters ranging from shallow, inshore waters down to depths of up to 1,100m. Spawning is thought to take place in deep water, with each female thought to produce just one batch of eggs (in a large, buoyant and gelatinous ribbon) in winter and spring (Laurenson *et al.* 2008). Juvenile monkfish descend to the seabed after 3-4 months spent in the water column and are generally found in shallower water than adults. Female monkfish do not mature until they are at least seven years old and so the species is particularly vulnerable to overfishing. Monkfish are ambush predators, enticing prey (typically fish, cephalopods and crustaceans) towards their mouths with a lure that extends from the top of their head (Fariña *et al.* 2008). Monkfish are generally viewed as typical demersal fish; however, large females, perhaps undertaking spawning migrations, are regularly caught in pelagic trawls in deep water between the UK and Norway (Hislop *et al.* 2001).

Several species of gurnard are abundant in UK waters. The grey gurnard (*Eutrigla gurnardus*) is very abundant in shallow, sandy areas and migrates inshore during summer (Wheeler 1978). Spawning takes place between January and June, with juveniles moving into deeper water as they mature. Red gurnards (*Aspitrigla cuculus*) tend to be found in shallow water and spawn over the summer months. Gurnards are predators, feeding on a range of fish, crustaceans and benthic invertebrates. The rays of their pectoral fins are modified into sensory organs with which they detect prey (Wheeler 1978).

The scorpionfish include bullrout (*Myoxocephalus scorpius*), sea scorpions (*Taurulus bubalis*) and pogge (*Agonus cataphractus*). These species tend to favour coarse sediments in shallow waters and may enter estuaries and river mouths (Power & Attrill 2002). They feed on crustaceans and small fish and pogge have sensory barbels to detect prey (Wheeler 1978). Most scorpionfish will spawn between October and April, with exact periods dependent on the species. They produce benthic eggs which they deposit on a secure holdfast (Ellis 2015b).

Dragonets are typically found inshore on sand and gravel sediments. Of the three main species found in UK waters, the common dragonet (*Callionymus lyra*) is the most abundant (Russell 1976). This species spawns in depths of less than 50m from early spring to August, producing pelagic eggs and larvae (Russell 1976). The other species found in UK waters are the spotted (*Callionymus maculatus*) and reticulated (*Callionymus reticulatus*) dragonets. Dragonets feed primarily on polychaetes, crustaceans and molluscs (King *et al.* 1994). Male dragonets are colourful fish with complex courtship behaviours. *C. lyra* develop iridescent stripes and an elongated snout, which is useless for feeding. Spawning occurs in pairs, with matched males and females ascending together through the water column and, if courtship has proved

satisfactory, releasing eggs and milt in synchrony on the descent. The starved male is likely to die following the spawning season (Daan 2015a).

Sandeels (Ammodytidae, principally *Ammodytes* spp. and *Hyperoplus* spp.) are shoaling species which lie buried in the sand at night and feed in mid-water during daylight (Winslade 1974). Spawning usually takes place between November and February, on sandy sediments. The eggs (up to 15,000 from a large fish (Gauld & Hutcheon 1990)) are demersal and are laid in sticky clumps on sandy substrates. Larvae remain pelagic for between 2-5 months after which they are thought to over-winter buried in the sand (Sparholt 2015). Boulcott & Wright (2011) discovered evidence of regional variation in fecundity of the lesser sandeel, *Ammodytes marinus*, with fecundity of individuals in areas of the central North Sea being greater than that of individuals of the north-east coast of the UK. There is little movement between spawning and feeding grounds, and so fishing activity may have a direct effect on spawning (Sparholt 2015). As well as being a major component of the industrial fishery, sandeels are an important food item for predatory fish and seabirds. There are five species of sandeels in the North Sea, of which *A. marinus* is the most abundant and comprise 90% of commercial landings (Marine Scotland website²).

There are 19 species of goby found in UK waters (Ellis & Rogers 2015). They are found in inshore waters and estuaries where they feed on a wide range of food from planktonic organisms to crustaceans. Gobies spawn on the seabed in summer, with males guarding eggs that have been left under rocks or in shells.

Bass (*Dicentrarchus labrax*) are attracted to warm water discharges and so are common inshore, close to the mouths of rivers, particularly around the southern coasts of the UK (Jennings & Ellis 2015). Bass move inshore to spawn from March to June and form large shoals during this migration, making them a target for fisheries.

The pipefish family includes the pipefish and seahorses. There are six species of pipefish and two seahorses found in north-western European waters. The short-snouted (*Hippocampus hippocampus*) and long-snouted (*Hippocampus guttulatus*) seahorses may be found around UK coasts, although are most common along the south coast of England. They are cryptic species and inhabit seagrass meadows in which they are well camouflaged. Seahorses are notable as the only animals that exhibit a genuine male pregnancy, with the female depositing fertilised eggs into the male brood-pouch, where they become embedded into the tissue. Fry are released after a gestation period of 3-5 weeks, during which they will obtain their nutritional requirements through a placenta-like structure. Breeding takes place over the summer months in UK waters (Wheeler 1978). Juveniles are pelagic, where predation levels are high, but those that survive youth will settle at an appropriate demersal habitat. Pipefish are found around the UK coast and can survive in a wide range of habitats from coastal seagrass meadows to oceanic waters. The snake pipefish (*Entelurus aequoreus*) underwent a remarkable increase in abundance in the early 2000s in UK waters, although since the 2008, the abundance has declined just as sharply (Daan 2015b).

A1a.4.2.3 Elasmobranchs

The most abundant sharks found in UK waters are the lesser and greater spotted dogfish (*Scyliorhinus canicula* and *Scyliorhinus stellaris*), the spurdog (*Squalus acanthias*) and the tope (*Galeorhinus galeus*). Dogfish show a wide but patchy distribution in the North Sea. They feed

² <http://www.gov.scot/Topics/marine/marine-environment/species/fish/sandeels>


on crustaceans, cephalopods and fish (Ellis *et al.* 1996) and are egg layers, with the peak of breeding in June and July. Tope are also widespread and juveniles are often found in large bays and estuaries. They are long-lived, reaching an age of at least 36 years (Peres & Vooren 1991). They are viviparous and young are generally born during the summer, after a year-long gestation period. Large open water species such as the porbeagle (*Lamna nasus*) may occasionally occur around the coast of the UK. Waters to the west of Scotland, beyond the continental shelf, are home to a large number of deep-water shark species. Many of these species remain mysterious, with little known about their biology and behaviour.

There are thought to be 27 species of skate within the north-east Atlantic (Ellis *et al.* 2015), of which many are present in UK waters. Among the most widespread are the thornback ray (*Raja clavata*) and the cuckoo ray (*Raja naevus*). The starry ray (*Amblyraja radiata*), the blonde ray (*Raja brachyuran*), the small-eyed ray (*Raja microocellata*), the undulate ray (*Raja undulata*) and the spotted ray (*Raja montagui*) are regionally abundant. The common skate (*Leucoraja batis*), listed as “critically endangered” on the IUCN Red List, is also present, but rare.

Basking shark

The basking shark (*Cetorhinus maximus*) is the largest fish in the North Atlantic and the second largest in the world. It is widespread and feeds by filtering plankton from the water with its gill rakers. The shark is commonly seen at the surface in the summer months, particularly around the western coasts of the UK, but it is less clear where they spend the winter. Research (Sims *et al.* 2003) suggests that they make extensive migrations both vertically and horizontally to locate high concentrations of plankton that will often be associated with fronts, and that they principally migrate north to south during the winter months along the continental shelf of Europe (Sims *et al.* 2005a, b). However, in the summer of 2007 one of two sharks tagged in the Irish Sea was found to migrate more than 9,500km over a period of ca. 2.5 months to a point off Newfoundland, Canada, reaching depths of over 1,200m on the way (Gore *et al.* 2008). Basking shark sightings in UK waters are typically centred on three main areas: the Minch, the Isle of Man and Cornwall (Solandt & Chassin 2014, Witt *et al.* 2012).

Figure A1a.4.1 – UK basking shark sightings reported by the general public in 2013 (left) and total sightings between 1987-2013 (right)



Source: Solandt & Chassin (2014)

A1a.4.2.4 Diadromous species

Diadromous species are those which migrate between marine and freshwater as part of their lifecycle.

Lampreys are eel-like, jawless fish. Both the river lamprey (*Lampetra fluviatilis*) and the sea lamprey (*Petromyzon marinus*) migrate up rivers to spawn (anadromy) and spend the larval stage buried in muddy substrates in freshwater. Both species need clean gravel for spawning, and silt or sand for the burrowing juveniles (JNCC website³). Once metamorphosis takes place, the adults migrate to the sea where they live as a parasite on various species of fish (Kloppmann 2015a). Sea lampreys are thought to venture further out to sea and spawn in lower reaches of the rivers than the river lampreys (Kloppmann 2015a). The sea lamprey is uncommon in the UK, while the river lamprey or lampren is widespread with substantial populations in some rivers and streams. The main populations of both species are found in the Bristol Channel and adjacent offshore waters.

The allis and twaite shads (*Alosa alosa* and *A. fallax*) and the Atlantic salmon (*Salmo salar*) also display an anadromous lifecycle. The shads are clupeids, or herring-like fish. They are rare, but are most abundant around the west coast and the southern North Sea, where they feed in estuaries before moving upstream to spawn between April and July. A water temperature of 12-20°C is thought to trigger this upstream migration (Dickey-Collas *et al.* 2015). Significant

³ <http://jncc.defra.gov.uk/protectedsites/sacselection/species.asp?FeatureIntCode=S1095>

spawning populations of the twaite shad are still found in the UK, while the allis shad, displays little spawning activity in the UK. Juveniles are thought to remain in freshwater for up to two years, before returning to the sea (Maitland & Hatton-Ellis 2003).

Salmonids, including Atlantic salmon (*Salmo salar*) and sea trout (*Salmo trutta*) undertake extensive migrations out to sea to feed, before returning to “home” rivers to spawn, which takes place in the late autumn to winter in UK rivers. Spawning takes place in shallow excavations (redds), in shallow gravelly areas in clean rivers and streams. After a period of 1-6 years the young salmon migrate downstream to the sea as smolts. Salmon have a homing instinct and spawn in the river of their birth after 1-4 years at sea (Heessen & Daan 2015).

Extensive research on the freshwater phase of the salmon life cycle has revealed much about the factors affecting juvenile production, but much less is known about the salmon’s life at sea. Atlantic salmon leave their home rivers in spring and early summer as smolts, and migrate towards feeding areas in the Nordic Seas and West Greenland (Guerin *et al.* 2014). Juvenile fish, including herring, sandeel and blue whiting form an important part of the diet of smolts during oceanic feeding (Haugland *et al.* 2006). New information is improving our understanding of smolt migration routes and food resources (Malcolm *et al.* 2015, Malcolm *et al.* 2010).

Recapture data from tagged adult salmon strongly suggest that the oceanic homing migration, the opposite of the smolt migration, is independent of currents with homing fish often moving along the shortest distance from tagging site to the coast (Hansen *et al.* 1993). Returns of salmon to western rivers are confined largely to the summer months. In contrast, sea trout appear to remain within nearshore waters rather than undergoing extensive migrations leading to concerns about their greater risk of exposure to sea lice infections from salmon farms in these areas (Gillibrand *et al.* 2005).

Salmonids play a critical role in the life cycle of the freshwater pearl mussel *Margaritifera margaritifera*. The freshwater pearl mussel is long lived with records of individuals over 100 years old (Bauer 1992). The larval stage (or glochidia) of the mussel is inhaled by juvenile Atlantic salmon and brown or sea trout where it attaches to the gills and encysts. Encysted larvae live and grow in the hyper-oxygenated environment on the gills before dropping off in the following spring, an infection apparently without a negative influence either on the lifecycle or the health of the fish (Makhrov & Bolotov 2011).

The European eel (*Anguilla anguilla*) spends most of its life in freshwater or inshore coastal waters, before migrating across the Atlantic to the Sargasso Sea to spawn in late summer (McCleave & Arnold 1999). The larvae drift north-east with the Gulf Stream and return to European coastal waters during the spring where they transform into transparent elvers (glass eels). Glass eels gather in river estuaries and wait for the river water to reach 10-12°C, before swimming upstream and migrating into inland waters. Peak migration takes place on the increasing tides in April and May, with predators such as porbeagle shark taking advantage (Béguer-Pon *et al.* 2012). Eels which successfully reach fresh water acquire green and brown pigments and become yellow eels. Yellow eels spend between 2 and 20 years in rivers and other inland waters although there are records of longer lived animals. Mature fish migrate seawards as silver eels and are thought to migrate westwards at depth to the Sargasso Sea, where they spawn and die. Experimental studies suggest eels are able to use innate magnetic compass orientation to orient themselves relative to magnetic North and, in conjunction with sea temperatures, navigate the long distances required to undertake these migrations (Durif *et al.* 2013).

Very large declines in eel populations since the 1970’s have been reported throughout Europe. Bark *et al.* (2007) concluded that eel stocks in some English west coast rivers are probably still

at or near to carrying capacity, with male-dominated populations. In other rivers, particularly those towards the south-east of England, current and historical data indicate declining female-dominated stocks. Nevertheless, the number of new glass eels entering European rivers declined to 10% and by some figures, to just 1% of former levels, although recruitment seems to have undergone a slight increase over the last 3-4 years (ICES website⁴).

A1a.4.2.5 Shellfish

The benthic fauna of the UK waters is rich and diverse. An important component of this benthic fauna is a collection of molluscs and crustaceans loosely referred to as shellfish. This section considers shellfish of commercial importance, with other species considered in detail in the Benthos section (Section A1a.3). Detail on the fisheries for these species may be found in Section A1h.13.

Crustaceans

The Norway lobster (*Nephrops norvegicus*), commonly known as *Nephrops*, lives in burrows dug into muddy and sandy sediments, at depths between 20-800m. They range in body length from 8-24cm. *Nephrops* feed mainly on detritus, small crustaceans and worms and are most active at night (Marine Scotland website⁵). Eggs hatch in spring or summer after being carried by females for 9 months. The relative inactivity of females during this period, when they remain hidden in burrows, means that males are more heavily exploited in the fishery through most of the year (Marine Scotland website⁵). There is considerable variation in the life-histories of *Nephrops* at different locations. In part, this is linked to sediment type, with higher population densities found at sandier sites, resulting in a reduction in the rate of growth and maximum size (Marine Scotland website⁵). *Nephrops* is more abundant in northern UK waters, although significant populations exist in the Irish and Celtic Seas and on the Fladen Ground.

The brown (or edible) crab (*Cancer pagurus*) is most abundant on rocky grounds, where it hides in holes and crevices. The crab is generally found in shallow water close to shorelines, particularly along the east coast and the south-west of England, although it can be found in water as deep as 100m (FAO website⁶). It is both an active predator and a scavenger. The species spawns between November and February, during which time the females remain in deeper waters offshore (Edwards 1979). Other crabs that are abundant around UK coasts include the green crab (*Carcinus maenas*) and velvet crab (*Necora puber*). The European lobster (*Homarus gammarus*) is found from the shoreline to depths of 150m, usually on a hard substrate such as rock or hard mud, growing to lengths of 60cm. Lobsters are most active at night, remaining in crevices during the day. Females lay eggs in July and carry them for 10 or 11 months (FAO website⁶). The related crawfish *Palinurus elephas* is another valuable crustacean but catches are much reduced since the 1970s. It is most abundant on hard bottoms off the extreme south-west of England and Wales and the west of Scotland. Crawfish feed on a variety of benthic organisms but are thought to have a preference for echinoderms such as urchins. Long distance migrations are a feature of many crabs and lobsters, the edible crab, European lobster, crawfish and spider crab (*Maja squinado = brachydactyla*) (Edwards 1979)

⁴ <http://www.ices.dk/news-and-events/news-archive/news/Pages/Latest-ICES-advice-on-European-Eel---stocks-remain-critical.aspx>

⁵ <http://www.gov.scot/Topics/marine/marine-environment/species/fish/shellfish/nephrops>

⁶ <http://www.fao.org/fishery/species/search/en>

A number of valuable shrimp species are found around the UK. The three most important are the brown shrimp (*Crangon crangon*), the pink shrimp (*Pandalus montagui*) and the deep-water shrimp (*Pandalus borealis*). The brown and pink shrimps are typically found in shallow waters, in bays and estuaries along the east coast of England. The brown shrimp generally favours areas with soft, sandy sediments, in which it can burrow, while the pink shrimp is more common over hard substrates. Eggs are carried by females over the winter months, before hatching in spring (Lee & Ramster 1981). The deep-water shrimp is larger and longer lived than the other species. It has a more northerly distribution in UK waters and the North Sea represents the southern edge of its range. It is a detritivore, found as deep as 900m, typically over muddy sediments and in areas of slow moving water where detritus accumulates (Lee & Ramster 1981).

Molluscs

The most commercially valuable molluscs are scallops (*Pecten maximus*). Scallops are found predominantly to the south and west of the UK on sandy, muddy, shell and gravel substrates, down to depths of over 100m. They occupy depressions in the sediment and are able to escape danger by swimming using jet propulsion (Chapman 2004). Their shells are lined with eyes and sensory tentacles, allowing them to detect light levels and even to form rudimentary images (Marine Scotland website⁷). Scallops are filter feeders, sieving the water for phytoplankton and suspended detritus. They first spawn in autumn, at two years old. Older scallops also spawn in spring (Mason 1983). Queen scallops (*Aequipecten opercularis*) are a much smaller shellfish, without the upward facing flat valve characteristic of *P. maximus*. Habitats and distributions of the two species are similar, but queen scallops are able to live on harder gravel and shell substrates (Chapman 2004) and have a much greater ability to swim.

Cockles (*Cerastoderma edule*) live on inter-tidal beaches of sand, muddy sand and fine gravel, where they burrow into the sediment. They use a siphon tube to feed on material suspended in the water column and can be found in very high densities (several 100/m²) (Chapman 2004). Cockles mature after 2 years and spawn in spring, with each female producing up to a million eggs. The eggs spend about a month in the water column, meaning that settlement of spat and subsequent recruitment can be highly variable (Chapman 2004).

Mussels (*Mytilus edulis*) are suspension feeders generally found attached to hard substrates within the inter-tidal zone, although they also attach to reefs and man-made structures in shallow waters (Chapman 2004). Mussels reach maturity after one year and each female can release over 5 million larvae (Bannister 1998, cited by Chapman 2004), with spawning taking place in late spring. The settlement of spat is influenced by a range of factors, including tidal currents, water temperature and predation and so can be variable (Bannister 1998, cited by Chapman 2004). Other bivalve molluscs common in UK waters include the horse mussel (*Modiolus modiolus*) (although these are little exploited) and razor clams (*Ensis arcuatus* and *Ensis siliqua*).

The most harvested gastropod molluscs in UK waters are whelks (*Buccinum undatum*) and periwinkles (*Littorina littorea*). Whelks are carnivorous, mobile species found close to the coast along rocky shores and on soft sediments. They spawn in November, with eggs attaching to the seabed (Chapman 2004). Winkles are herbivorous, typically found attached to rocks in the inter-tidal zone. They spawn between January and July, releasing planktonic egg capsules (Chapman 2004).

⁷ <http://www.gov.scot/Topics/marine/marine-environment/species/fish/shellfish/scallop>

A1a.4.2.6 Spawning and nursery grounds

Most fish display external fertilisation and therefore need to aggregate in large groups to coordinate spawning. Figures A1a.4.2 – A1a.4.5 show the spawning sites of 13 selected fish species and *Nephrops* around the UK coast. These maps are based on a wide-ranging study by Coull *et al.* (1998). This study was added to by a larger survey of spawning and nursery sites conducted by Ellis *et al.* (2010, 2012). This survey covered more fish species and has been used as a reference source for this section. However, as an illustration, Coull *et al.* (1998) provides an excellent visual representation. In some instances (particularly herring, in which spawning is very site specific) they may be broad in their overview and areas indicated should be regarded as areas in which spawning may occur, rather than well-defined zones. Table A1a.4.1 indicates the main spawning periods of selected fish species around the UK. These periods are estimates, based mainly on plankton trawls and spawning periods are likely to vary with location. This is particularly notable in herring, which has a number of separate spawning sites, in use at different times of the year.

Table A1a.4.1: Spawning periods of selected species and groups

Group	J	F	M	A	M	J	J	A	S	O	N	D
Dab, long rough dab & solenette												
Plaice												
Sole												
Lemon sole												
Turbot & megrim												
Gurnards												
Scorpionfish												
Dragonets												
Bass												
Haddock												
Whiting												
Cod & saithe												
Norway pout & poor cod												
Dogfish & tope												
Monkfish												
Sandeels												
Argentines												
Blue whiting												
Northern North Sea herring												
Southern North Sea & English Channel herring												
Western stock herring												
Sprat												
North Sea mackerel												
Western stock mackerel												

Source: Information taken from Coull *et al.* (1998) and Cefas (2007)

Juvenile fish are vulnerable to predators and harsh conditions in the open water. Therefore it is typical for juvenile fish to stay in sheltered nursery grounds which provide an abundance of food. Selected nursery grounds are shown in Figures A1a.4.6 – A1a.4.9, with the information taken from Coull *et al.* (1998). These figures should be referred to where nursery sites are mentioned in this baseline. At nursery and spawning grounds, fish aggregate in large numbers and so are particularly vulnerable to disturbance.

Figure A1a.4.2: UK spawning grounds




Figure A1a.4.3: UK spawning grounds




Figure A1a.4.4: UK spawning grounds




Figure A1a.4.5: UK spawning grounds




Figure A1a.4.6: UK nursery grounds




Figure A1a.4.7: UK nursery grounds




Figure A1a.4.8: UK nursery grounds





Figure A1a.4.9: UK nursery grounds



A1a.4.3 Features of Regional Sea 1

A1a.4.3.1 Fish community

The northern North Sea is an ecosystem characterised by oceanic inflows from the Norwegian Sea and Atlantic Ocean and by seasonal stratification in the water column. Species diversity within the fish community is not as great in the central and northern North Sea as in the southern North Sea, with the exception of some areas of the Scottish coast (Callaway *et al.* 2002). The highest richness of northerly species is found in waters off the north-east of Scotland, including Orkney and Shetland (Daan 2006). Statistical analyses of North Sea fish assemblages using data collected in groundfish surveys (Callaway *et al.* 2002, Reiss *et al.* 2010) found that the 50m depth contour that separates the southern North Sea from the central and northern regions acts as a conspicuous boundary between different fish assemblages. Within the northern North Sea, a number of assemblages were identified, with divisions typically occurring at the 100m depth contour. At depths between 50-100m, the benthic community was dominated by dab and long rough dab. Of species more loosely associated with the seabed, haddock, whiting, herring and plaice were dominant. At depths of between 100-200m, the community was characterised by long rough dab, hagfish (*Myxine glutinosa*) and Norway pout. In the deepest areas of the region, below 200m, assemblages were notable for the low abundances of demersal fish and the prevalence of silvery pout (*Gadiculus argenteus*). The proportion of large fish (body length >40cm) by weight in the region was estimated at 22% in 2014 (Defra 2015).

A1a.4.3.2 Pelagic species

In the North Sea, there are thought to be three sub-populations of herring which spawn at different times, although the complex migratory behaviour of herring makes the population structure difficult to define. These groups are mixed for the majority of the year, but migrate to different spawning grounds during the breeding season (Daan *et al.* 1990), although fidelity to particular spawning sites may not be absolute (Slotte & Fiksen 2000). One of these populations is of importance in the northern North Sea - Buchan / Shetland herring, which spawns off the north-east coasts of Scotland and Shetland in August and September (Coull *et al.* 1998, Ellis *et al.* 2012). Juveniles migrate into nursery grounds in coastal areas such as the Moray Firth and the Firth of Forth. North Sea mackerel overwinter in deep water to the east and north of Shetland, before migrating south to spawn between May and August (Coull *et al.* 1998, Ellis & Heessen 2015a). Spawning may take place as far south as the Dogger Bank and following spawning, North Sea mackerel will mix with immigrant western stock mackerel in northern North Sea feeding grounds, before returning to over-wintering sites (Lockwood 1988). Juvenile fish remain in nursery areas in the shallow waters of the Southern Bight (Coull *et al.* 1998). Other pelagic species abundant in the region include sprat, which spawns over a wide area of the region within the summer months and the argentines, typically found in deeper waters.

A1a.4.3.3 Demersal species

Many of the larger, more commercially valuable gadoid species are abundant in the deeper waters of the central and northern North Sea. Whiting is one of the most numerous and widespread species. Recaptures of tagged whiting and the use of parasitic markers, show that the populations to the north and south of the Dogger Bank form separate populations (Hislop & MacKenzie 1976, Tobin *et al.* 2010). Whiting is one of the major predators of commercially important fish stocks in the North Sea (ICES 1991). In the region, major spawning areas are found north of the Dogger Bank and east of Scotland and the spawning season extends from February to June (Coull *et al.* 1998, Ellis *et al.* 2012). Much of the North Sea acts as a nursery ground for <1 year old whiting within the pelagic phase, particularly around the east coasts of Scotland and England (Coull *et al.* 1998, Ellis *et al.* 2012). Analysis of size-class indicate that larger whiting (>20cm length) are more abundant in waters to the east of Shetland and Orkney

than smaller fish, which tend to congregate in inshore areas such as the Moray Firth and northeast England (Hislop *et al.* 2015). Haddock is found throughout the northern North Sea, where spawning takes place during spring between Shetland and the Norwegian Deep (Coull *et al.* 1998), before adults disperse west and into the central North Sea to feed (Albert 1994). During the summer months, the density of juveniles is greatest off the north-east coast of Scotland, although once haddock have reached a size of 4-8cm, they adopt a demersal way of life and distributions of juveniles and adults overlap almost completely (Hislop *et al.* 2015).

Cod are more abundant in the northern North Sea than in the southern North Sea (a trend that has become increasingly distinct over recent decades (Hislop *et al.* 2015)) and there are a number of spawning sites in the region (Coull *et al.* 1998, Ellis *et al.* 2012), with significant larval abundance off the north coast of Scotland and in the Moray Firth (Edwards *et al.* 2011). Nursery areas are most extensive in the eastern North Sea, but the Firth of Forth and other coastal, shallow areas in the north-western North Sea act as important nurseries in the region (Coull *et al.* 1998, Gibb *et al.* 2007, Hislop *et al.* 2015). Tagging experiments have shown that there is a limited exchange of individuals between the northern North Sea and waters to the west of Scotland, although migrations are generally limited to 200 miles (Rogers & Stocks 2001). Analysis suggests that genetic differentiation of North Sea cod is greatest between those living at depths >100m and those in shallower areas (Heath *et al.* 2014). Analysis of size-class indicates that larger cod (>25cm) are abundant both along the coasts of Scotland and north-east England and in deeper, offshore waters, east of Shetland, while smaller cod are more restricted to coastal areas, both in the east and west of the North Sea (Hislop *et al.* 2015). Young saithe migrate into coastal waters of the north of Scotland, Orkney and Shetland, where they remain for 3-4 years, before recruiting to stocks in the northern North Sea (Newton 1984). Abundances are greatest in the deeper waters north of the 100m contour (Hislop *et al.* 2015). There are extensive spawning and nursery grounds for the species in the region. Other large gadoids that are abundant in the region include pollack and ling (*Molva molva*).

Small gadoid species include Norway pout and silvery pout. Norway pout is a small, commercially valuable species of particular importance to the industrial fleet. It is found throughout the central and northern North Sea, typically at depths of between 80-200m. Fish mature at 1-2 years and spawning takes place between January and April on the continental shelf and slightly later in deeper waters. No specific spawning sites have been identified and it is believed that the activity is widely dispersed. Silvery pout is a shelf edge species, commonly found at between 200-500m. They are small, short-lived, benthopelagic fish which spawn from mid-winter to spring. A number of flatfish are present in the region, including plaice (although juveniles are largely restricted to sheltered coastal waters (Goldsmith *et al.* 2015)), lemon sole, dab and witch (*Glyptocephalus cynoglossus*). Monkfish are widely distributed across the northern and central North Sea. Spawning is thought to take place in deeper waters and the presence of nursery sites in shallow waters around Shetland has been proposed (Laurenson *et al.* 2008), although juveniles are widely distributed around the North Sea (Ellis *et al.* 2012).

Hagfish are a very primitive group of fish, eel-like and lacking true jaws. They are remarkable for their ability to produce large quantities of mucus. The species is widely distributed throughout the northern North Sea usually in depths greater than 50m (and more abundant in deeper waters), where they favour muddy sediments in which they can bury themselves (Kloppmann 2015b). They are scavengers of dead and dying fish that have sunk to the seabed, but will also feed on benthic invertebrates (Shelton 1978). Very little is known about their life history, although it is thought that there may be 3 annual spawning events (Grant 2006), during which females produce a small number of large eggs.

Grey gurnards are a dominant demersal species in the North Sea, having undergone a substantial population increase in recent years (Sell & Heessen 2015). They are most

abundant in the central North Sea, to the north of the Dogger Bank during winter, their distribution spreading eastwards over summer (Sell & Heessen 2015). In the North Sea, they tend to live at depths of between 70-100m although they may move in surface or pelagic waters both during spawning and at night (Sell & Heessen 2015). Predation on juvenile gadoids by large gurnards may be a contributing factor slowing the population recovery of cod and whiting in the North Sea (Floeter *et al.* 2005).

A1a.4.3.4 Elasmobranchs

The most abundant sharks found in the northern North Sea are the lesser and greater spotted dogfish, spurdog and the tope. The starry ray is the most abundant skate, mainly found in the northern and central North Sea, between 50m and 100m depth (Ellis *et al.* 2015). The cuckoo ray has an abundance hotspot off the Aberdeenshire coast (Ellis *et al.* 2015). Dogfish show a wide but patchy distribution in the North Sea. Sightings of other species, such as the common skate, basking shark and porbeagle are rare in the northern North Sea (Rogers & Stocks 2001).

A1a.4.3.5 Diadromous species

The sea lamprey and river lamprey have both been recorded in the region. Although once an important fishery, lampreys are now rarely caught in large numbers. Several rivers in the region have a high concentration of salmon. In particular, the Rivers Spey, Tweed and Dee support large salmon populations. Sea trout are also present, but are more abundant on the west coast. The rivers and estuaries of the region provide an important habitat for the European eel (Robson 1996).

A1a.4.3.6 Shellfish

The soft, sandy sediments of the Fladen Ground provide an important habitat for *Nephrops* and *P. borealis*. These species are also found over soft sediments around the Farne Deep, while important populations of *Nephrops* also exist in the Firth of Forth and the Moray Firth (Lee & Ramster 1981). Scallop grounds may be found along the northern coast of the Moray Firth (Mason 1983). Cockles are particularly abundant in the Firth of Forth and at Findhorn Bay and the Culbin Bars in the Moray Firth (Chapman 2004), while mussels are harvested from the Dornoch Firth. Brown crabs, green and velvet crabs are widespread in coastal waters of the region, as are whelks and periwinkles.

A1a.4.4 Features of Regional Sea 2

A1a.4.4.1 Fish community

The southern North Sea is a dynamic ecosystem characterised by a sandy, flat, shallow seabed and considerable tidal mixing. Species diversity within the fish community is greater in the southern North Sea than in the central or northern North Sea (Callaway *et al.* 2002) and within the southern North Sea, fish diversity is greatest in the west (Rogers *et al.* 1998). Callaway *et al.* (2002) found that the southern North Sea is characterised by a high abundance of small demersal species (in 2014, fish in the region >40cm made up 6% of total weight (Defra 2015)), typically found closely associated with the seabed, including solenette (*Buglossidium luteum*), dab and common dragonet. Of species more loosely associated with the seabed, three distinct assemblages were identified in the region. The two most extensive of these could be characterised by whiting, grey gurnard, horse mackerel and dab. The third assemblage, located in the far south of the region, was characterised by high numbers of horse mackerel and mackerel. Reiss *et al.* (2010) identified whiting, dab, plaice, grey gurnard, and lesser weever (*Echiichthys vipera*) as characteristic demersal species in the southern North Sea.

Meanwhile, Corten & van de Kamp (1996) identified twelve, "southern," species within the North Sea either as those with a greater abundance in the southern North Sea during summer than

during winter, or as those with a distribution restricted to the southern North Sea. These species were: poor cod, bib, red mullet (*Mullus surmuletus*), sardine (*Sardina pilchardus*), lesser weever, anchovy (*Engraulis encrasicolus*), tub gurnard (*Chelidonichthys lucerna*), John Dory (*Zeus faber*), bass, black sea bream (*Spondyliosoma cantharus*), horse mackerel and mackerel. Other common species in the southern North Sea include pogge or hooknose (*Agonus cataphractus*), flounder (*Platichthys flesus*) and sand gobies (*Pomatoschistus minutus*) (Cefas 2007).

A1a.4.4.2 Pelagic species

There are two main herring populations of importance in the southern North Sea: Dogger Bank, or Banks, herring, which spawns from August to September or October; and Southern Bight or Downs herring which spawns from November to January in the Southern Bight of the North Sea and the Eastern English Channel (Coull *et al.* 1998, Ellis *et al.* 2012), after which pelagic larvae will drift towards important nursery grounds such as the Humber Estuary, Thames Estuary and the Wash. There is also a small spring-spawning stock of herring that spawns within the Thames estuary (Ellis *et al.* 2012). North Sea mackerel overwinter in deep water to the east and north of Shetland, before migrating south to spawn between May and August (Coull *et al.* 1998, Ellis *et al.* 2012). Catch rates of mackerel in the southern North Sea are much greater in summer and autumn than in winter and spring (Ellis & Heessen 2015a). Spawning may take place in waters as far south as the Dogger Bank and juvenile fish remain in nursery areas in the shallow waters of the Southern Bight (Coull *et al.* 1998), although Ellis & Heessen (2015a) report catch rates of small (<22cm) mackerel are much lower in the southern North Sea than larger mackerel, suggesting most juveniles move away from the area.

Sprat are most abundant in the shallow waters of the southern North Sea. Spawning takes place along the east coast of England from May to August with a peak during May and June (Coull *et al.* 1998). Nursery grounds are found around the Southern Bight and Dogger Bank and sprat mature to spawn for the first time at the age of two years. Other clupeids found in the southern North Sea include anchovies (spawn from April to August) and sardines (spawn from June to August). Horse mackerel around the UK are split into two stocks; a western stock and a North Sea stock. The North Sea stock spends most of the year in the central North Sea, Skagerrak and Kattegat, but migrates to the southern North Sea in the summer to spawn (Ellis 2015a). After spawning in the Southern Bight, horse mackerel will generally disperse to the north-east, although later in the year, they will tend to migrate west to over-winter in the warmer waters of the Channel and Celtic Sea (Ellis 2015a).

A1a.4.4.3 Demersal species

Small gadoid species, including poor cod and bib, are typical of the southern North Sea community. In general, poor cod is found in coastal waters at depths of between 25-300m, where they feed on a range of small crustaceans and fish. The species spawns in winter at depths of between 50-100m. Bib tends to congregate in large schools around reefs and wrecks. They spawn in shallow waters, between March and April.

The Southern Bight is a major spawning area for whiting. Much of the southern North Sea acts as a nursery ground for <1 year old whiting while in the pelagic phase, particularly around the mouth of the Thames and on the Dogger Bank (Coull *et al.* 1998). Cod are more abundant in the northern North Sea, but the southern North Sea remains an important area for spawning, with the main spawning season extending from January to April (Coull *et al.* 1998). The south-east North Sea acts as an important nursery ground for 1 and 2 year old cod. Righton *et al.* (2007) showed that there is a significant level of migration of cod from the southern North Sea northwards in spring and summer, although no migration was apparent between the southern North Sea and the English Channel. They also noticed behavioural differences (the use of tidal

stream transport in migrations by southern North Sea cod was not commonly observed in English Channel cod) between cod stocks in the region, suggesting different selection pressures may exist in the two areas.

Flatfish of particular importance in the region include solenette and dab, plaice, sole and lemon sole. Plaice is found in greatest abundance in the southern North Sea. Plaice spawn throughout the shallower parts of the southern North Sea, including the Dogger Bank and the Southern Bight, with spawning taking place between December and March (Goldsmith *et al.* 2015). Sandy, shallow bays on the coasts of England and continental Europe act as important nursery grounds for plaice, with juveniles moving further offshore as they mature. While large (>15cm) plaice are widespread and abundant over the southern North Sea, smaller, juvenile plaice are largely restricted to the south and east of the Dogger Bank, with the French and Dutch coasts having the greatest densities (Goldsmith *et al.* 2015).

The sole is close to the northern limits of its distribution in the southern North Sea and is confined to areas where temperatures do not fall below 5°C for prolonged periods (Horwood 1993). Consequently, during cold winters, dense aggregations of sole are known to accumulate in deeper, warmer waters, and will congregate in seabed pits and holes such as Silver Pit, where they become an easy target for fishermen (Rijnsdorp *et al.* 2015). Sole spawn in shallow, inshore areas and sandbanks between March and May. Major spawning grounds include the Thames Estuary and the Norfolk Banks (Coull *et al.* 1998, Ellis *et al.* 2012). Larvae are pelagic for approximately one month, after which they metamorphose into the demersal phase. As a result, local abundances of <1 year old sole are likely to reflect the spawning success of local spawning aggregations (van Beek *et al.* 1989). Nursery grounds are situated in coastal waters shallower than 20m (Rijnsdorp *et al.* 2015).

A1a.4.4.4 Elasmobranchs

The most abundant sharks found in the southern North Sea are the lesser and greater spotted dogfish and tope (a summer visitor). The outer Thames Estuary and the Wash are important areas for a number of ray species, including thornback rays, adults of which migrate into the Thames Estuary to breed in summer (Walker *et al.* 1997). Sightings of other species, such as the common skate, basking shark and porbeagle are rare in the southern North Sea (Rogers & Stocks 2001).

A1a.4.4.5 Diadromous species

The rivers and estuaries of south-east England provide an important habitat for the European eel. The up-river migration of elvers in the Thames occurs between April and October, with peak movement of eels in May and June (Naismith & Knights 1988). Eels are mainly found in the shallower waters of the North Sea, suggesting they are resident yellow eels (Walker & Ellis 2015). The sea lamprey and river lamprey are rare in the region, although they once comprised an important fishery in the rivers of south-east England. The allis and twait shad are present in the Thames Estuary and catches at sea are most commonly reported along the eastern English coast (Dickey-Collas *et al.* 2015). The Atlantic salmon and sea trout are most abundant around the northern and western coasts of the UK, but are also present in the southern North Sea. Large numbers of sea trout are found off the north coast of Norfolk, feeding on sprat and sandeels, prior to returning to their home rivers in north-east England (Aprahamian & Robson 1998).

A1a.4.4.6 Shellfish

The main site for *Nephrops* is present to the north and west of the shallow Dogger Bank, while pink and brown shrimp are abundant in the Wash and the Thames and Humber Estuaries. The east coast of England is a site of particularly intense spawning by brown crab (Rogers & Stocks

2001) but they are found throughout the region. Large populations of cockles are found in the Wash and the Thames Estuary. Mussels are abundant in the Wash, as are wild and cultivated oysters along the Essex and Kent coast (Rogers & Stocks 2001). Whelks and periwinkles are widespread in the region. Razor clams, including the introduced species *Ensis directus* as well as native species, are abundant in the Wash and locally elsewhere.

A1a.4.5 Features of Regional Sea 3

A1a.4.5.1 Fish community

The English Channel acts as a biogeographical boundary between the western waters and the North Sea for many species, but is also an important migration route for others (Defra 2005). The region supports a similar fish community to the southern North Sea and many species, such as plaice, migrate between the regions (Arnold & Metcalfe 1996). While much of the region is shallow with fine sediment on the seabed, some central areas such as the Dover Straits are deeper with coarser sediment and significant tidal streams (Kaiser *et al.* 1999). These habitats support a high diversity of invertebrates which in turn support small gadoids such as bib and poor cod, along with demersal species such as thickback sole (*Microchirus variegatus*) and red gurnard (Defra 2005). Vaz *et al.* (2007) found the region to be strongly spatially structured. The lower temperatures and salinities in coastal areas favour benthodemersal species (which particularly dominate when the autumn is cool and wet), while further offshore, or in warmer winters, pelagic species (horse mackerel) and demersal species (red mullet) tend to dominate the assemblage.

The seahorses *H. hippocampus* and *H. guttulatus* are most abundant in the English Channel (Defra 2005). Both species are on the IUCN red list (classified as data deficient).

A1a.4.5.2 Pelagic species

Mackerel is a seasonal visitor to the region, as the western spawning stock migrates eastward through the English Channel into nursery and feeding grounds in the southern and central North Sea over the summer months (Ellis & Heessen 2015a). The Channel represents the southern limit of the herring population and they are most abundant in the eastern Channel and along the French coast (Dickey-Collas *et al.* 2015), where they occupy summer and spring feeding grounds, although there is no major nursery or spawning ground in the region. Horse mackerel are abundant further offshore. Sprat are common in winter particularly in Lyme Bay and between Portland Bill and the Isle of Wight (Pawson & Robson 1998).

A1a.4.5.3 Demersal species

Of the gadoids, whiting and cod are common, with important spawning regions off east Sussex. Haddock and saithe are not common in the region, while other gadoids such as ling and pollack can be found around reefs and rocky outcrops (Pawson & Robson 1998).

Plaice and dab are the most abundant flatfish in the region, with important winter spawning taking place in the centre of the Channel (Goldsmith *et al.* 2015). Some of the North Sea population spawns in the English Channel (Houghton & Harding 1976). Dab spawn between January and June and they, along with sole, migrate to coastal nursery waters before moving into deeper waters as they grow. Lemon sole, flounder, turbot (*Psetta maxima*) and brill (*Scophthalmus rhombus*) are also found in the English Channel. Flounder nursery sites may be found in riverine and estuarine areas. Solenette, common dragonet and lesser weever fish are also abundant in the region (Kaiser *et al.* 1999).

In UK waters, black sea-bream are mainly found in the Channel, where they overwinter in the offshore western Channel before moving eastwards in spring to spawn and spend summer at

coastal feeding grounds (Heessen 2015b). Striped red mullet are also most abundant in the Channel, where they spawn in summer, and along the European coastline (Heessen 2015c). Bass are most commonly reported along the Channel and the southern North Sea. Juvenile fish congregate in inshore waters, but disperse into a range of inshore and offshore habitats as they mature (Jennings & Ellis 2015).

Seahorses are found along the seagrasses of the southern English coast often in sheltered areas such as Poole Harbour or the Solent. Studland Bay is an important site for the two species typically found in UK waters. The population at Studland Bay is subject of a long-running tagging and monitoring project run by the Seahorse Trust, as part of the British Seahorse Survey. This work has confirmed that seahorse populations along the south coast are resident, rather than migrating visitors from across the Channel (The Seahorse Trust 2012).

A1a.4.5.4 Elasmobranchs

The thornback ray, starry smooth-hound (*Mustelus asterias*) (Farrell *et al.* 2015) and lesser and greater spotted dogfish are present in the eastern English Channel. The distinctive undulate ray is abundant along the southern coast and around the Isle of Wight (Ellis *et al.* 2015). The basking shark is not common in this region, although is occasionally sighted in waters around the Isle of Wight (Swaby & Potts 1998) or the Hurd Deep, near the Channel Islands (Solandt & Chassin 2014).

A1a.4.5.5 Diadromous species

There are no major salmon rivers in the region, although the Rivers Ouse and Rother are important for sea trout. Eels are relatively abundant in the rivers of the south east of England, although sea lampreys are not abundant in the region (Swaby & Potts 1998).

A1a.4.5.6 Shellfish

Substantial scallop grounds are found along the coast of this region (Mason 1983). Cockles, mussels, periwinkles and whelks are all present at shorelines. Crabs and lobsters are abundant on rocky ground in the region, with brown crabs typically found further offshore than lobsters. The Solent supports a healthy population of native oysters (*Ostrea edulis*).

A1a.4.6 Features of Regional Seas 4 & 5

A1a.4.6.1 Fish community

The Western Approaches may be split into three main regions: the western English Channel, the Celtic Sea and the Bristol Channel. The fish communities of these regions are influenced by the Atlantic Ocean and warm water pelagic species are occasionally reported in the area (Stebbing *et al.* 2002).

The western English Channel is deeper than the eastern Channel, with a steep shelving seabed, leading to the open sea. Consequently, the fish community is significantly different from that found further east. Monkfish and cuckoo rays, which are virtually absent from the eastern Channel, are relatively abundant in the western Channel (Ellis & Velasco 2015, Ellis *et al.* 2015). The high number of wrecks and reefs in the region provide habitats for pollack and conger eels (*Conger conger*). In the Celtic Sea, offshore species become increasingly abundant with depth, and species such as hake (*Merluccius merluccius*), megrim (*Lepidorhombus whiffiagonis*), long rough dab, blue whiting and boarfish (*Capros aper*) are common (Warnes & Jones 1995). The Bristol Channel contains a number of important spawning and nursery grounds, such as Carmarthen Bay where a number of juvenile rays and flatfish mature (Ellis *et al.* 2012). Large numbers of the smalleyed ray are present in the Bristol Channel (Ellis *et al.* 2015).

A1a.4.6.2 Pelagic species

Mackerel is abundant at the shelf edge. Over winter they migrate into coastal waters off Cornwall, with the area acting as an important feeding ground (Ellis & Heessen 2015a). Herring are locally abundant, particularly in the Bristol Channel (Dickey-Collas *et al.* 2015) with a limited amount of spring spawning taking place to the south of Cornwall and Pembrokeshire (Coull *et al.* 1998). The western English Channel is notable for the populations of sardine and horse mackerel, which are abundant in the region. Horse mackerel is very abundant throughout the year in the waters along the southern coast of Cornwall, spawning along the western shelf edge (Ellis 2015a), while sardines spawn to the south of the Scilly Isles and Land's End in spring and early summer (Pawson 1995). Summer spawning sprat migrate inshore over the winter months. Argentines are also present in the region, particularly in the Bristol Channel (Heessen 2015a).

The enormous and peculiar ocean sunfish (*Mola mola*) is often observed basking at the surface off the coasts of Cornwall and Pembrokeshire, where they feed on jellyfish, fish larvae and other plankton.

A1a.4.6.3 Demersal species

Cod are numerous in the region, thought to move inshore over winter after spending summer feeding around deep water reefs and wrecks (Pawson 1995). Aggregations of spawning cod form in spring off Trevoze Head and to the south of Cornwall. Whiting are also abundant in inshore areas and spawn in similar locations to cod, while pollack and saithe are locally abundant, particularly around reefs and rocky outcrops. Haddock are relatively rare in the region, although more common in the northern part of the Celtic Sea, while hake may be found in the deeper waters of the region. The area around the Cornish peninsula acts as an important nursery ground for ling (Pawson 1995, Ellis *et al.* 2012). Poor cod and bib are also present.

Plaice and dab are the most commercially important flatfish in the region, although sole and lemon sole are also common. All these species spawn in the area, although the spawning periods are staggered throughout the year, perhaps avoiding competition (plaice from December to March, sole from March to May, lemon sole from April to September). These flatfish tend to move into deeper waters as they mature, with relatively little movement along the coast taking place. Sole is most abundant in the Bristol Channel and tends to be confined to areas with high sea bottom temperatures (Rijnsdorp *et al.* 2015).

Bass are abundant in inshore areas from spring as they migrate from warmer waters offshore. Bass spawn in the region from March to May and estuaries in the region, such as the Severn, act as important nurseries for juveniles. It is thought that the strongest recruitment takes place when the sea temperature is warmest at this time of year (Pawson *et al.* 2007). Sandeels are present throughout the region, and other species commonly found include sea bream, John Dory, red mullet, gurnards, conger eel and various wrasse. Rocklings, boarfish, greater pipefish (*Syngnathus acus*), pogge, butterfish (*Pholis gunnellus*), blennies and dragonets have been recorded in trawl surveys in this region.

A1a.4.6.4 Elasmobranchs

A number of ray species are present in the region, including the thornback ray and the cuckoo ray as well as the smalleyed ray and spotted ray (which are both locally abundant in the Bristol Channel) (Ellis *et al.* 2015). The thornback ray spawns in shallow bays around the region. Spurdogs, starry smooth-hounds and lesser and greater spotted dogfish may also be found. High catches of juvenile porbeagles in the Bristol Channel suggest this may be a nursery for the sharks (Bendall *et al.* 2013). Oceanic sharks such as blue sharks (*Prionace glauca*), thresher shark (*Alopias vulpinus*) and mako (*Isurus oxyrinchus*) sharks make occasional, seasonal visits

to the region. The Cornish coast is an area where basking sharks are particularly common, with numerous sightings reported annually in the summer months (Solandt & Chassin 2014).

A1a.4.6.5 Diadromous species

Salmon and sea trout are present in many rivers in the region, particularly along the south coast of Wales and are regularly caught in trawls in the Bristol Channel (Heessen & Daan 2015). The Bristol Channel is a particularly important region for shads and lampreys. The Rivers Severn, Usk, Wye and Tywi are the last four rivers in the UK known to contain spawning populations of twaite shad, and the allis shad is regularly recorded in the Severn Estuary, with the Tamar Estuary home to one of only two known UK spawning populations (DECC 2010, Maitland & Hatton-Ellis 2003). Eels are present in rivers throughout the region. Eels captured at sea in the region are typically found in shallow waters, and are likely to be resident yellow eels. Those caught in deeper waters are more likely to represent silver eels on their migration (Walker & Ellis 2015).

A1a.4.6.6 Shellfish

Offshore *Nephrops* grounds exist in the Celtic Sea, to the south of Ireland. Scallops (and to a lesser extent, queen scallops) are abundant on sandy sediments along the coasts of Dorset, Devon and Cornwall and also south Wales (Mason 1983). Brown crabs, green crabs, velvet crabs and lobsters are all abundant in the region, while the spider crab, a species with a south-westerly distribution in UK waters (Clark 1986), is particularly abundant in the region. The Severn Estuary, and particularly the Burry Inlet and Carmarthen Bay, provide important grounds for cockles, whelks and razor clams. Crawfish are more abundant off the extreme south-west of England and Wales than many other parts of the UK.

A1a.4.7 Features of Regional Sea 6

A1a.4.7.1 Fish community

Beam trawl surveys of the fish community of the Irish Sea identified three distinct categories of demersal fish (Defra 2005). Sandy, inshore areas are dominated by flatfish species, including plaice, solenette, sole and dab and other benthic-demersal fish such as the tub gurnard, lesser weever fish, dragonets and gobies. The offshore assemblage is characterised by species such as thickback sole, lemon sole and red gurnard as well as by elasmobranchs such as the greater spotted dogfish and the cuckoo and spotted rays (*Raja montagui*). The third grouping is found over muddy sediments found to the west of the Isle of Man and is characterised by the presence of witch and long rough dab. Otter trawl surveys reveal a distinction between western and eastern inshore fish assemblages in the region, with haddock, Norway pout and various clupeid species all more abundant in the west.

A1a.4.7.2 Pelagic species

Mackerel is present in the region, but there is little spawning or nursery activity in the region with the greatest abundances following the edges of the continental slope along the west coast of Ireland (Ellis *et al.* 2012). Herring is the most important pelagic stock in the Irish Sea, with particularly large abundance around the Isle of Man and Pembrokeshire. Spawning grounds are present to the east and to the north of the Isle of Man (late summer/autumn spawning) and off the Pembrokeshire coast near Milford Haven (spring spawning), with juvenile fish entering nursery grounds along the English and Irish coasts (Coull *et al.* 1998, Ellis *et al.* 2012). The Clyde Estuary also supports a spring spawning population. Sprat are widely distributed throughout the region, spawning over a wide area between May and August (Dickey-Collas *et al.* 2015).

A1a.4.7.3 Demersal species

Cod are widely distributed throughout the region in the summer, particularly in the deep water off the coast of Cumbria. Spawning grounds are situated in this area and to the west of the Isle of Man, with an important nursery ground located between the Isle of Man and the coast of Ireland. Whiting is abundant and widely distributed, particularly around coastal areas, while haddock, ling, pollack and saithe are found around wrecks and rocky reefs. Haddock are found in greatest numbers in the deeper waters to the west of the region. Important spawning grounds for whiting exist off the west of England and in Cardigan Bay, between February and June (Parker-Humphreys 2004, Ellis *et al*, 2012).

Important flatfish in the Irish Sea include plaice, sole and dab, which are found over sandy areas of the seabed. Spawning areas for these species exist in coastal areas in the east of the region, such as within Cardigan Bay. Lemon sole can be found in deeper offshore areas. Bass and grey mullet are seasonally abundant inshore, with abundance decreasing further north. Sandeels are widely distributed throughout the region and juvenile and some adult monkfish may also be found in coastal waters.

Other demersal species common in the Irish Sea include the poor cod, abundant in the deeper central waters, dragonets and lesser weevers which are particularly abundant in the shallow waters of Liverpool Bay, as well as coastal species such as grey and red gurnards, the thickback sole and the solenette (Parker-Humphreys 2004).

A1a.4.7.4 Elasmobranchs

There are a number of sharks and rays present in the region. Lesser and greater spotted dogfish, tope, cuckoo rays, thornback rays and spotted rays are all present, particularly around the north Wales coast and in deeper offshore regions. Anglesey and the Lleyn peninsula is an area of particular abundance for the greater spotted dogfish, with large numbers of egg cases washed up on local beaches (Defra 2005, Ellis 2015c). Cardigan Bay is a probable nursery site for thornback rays (Ellis *et al*. 2015). The basking shark is often observed in the region, particularly to the south and west of the Isle of Man, but also off the Pembrokeshire coastline. Satellite tracking reveals the Irish Sea to be an important migration route for basking sharks between the west of Scotland, the Isle of Man and south-west England (Witt *et al*. 2016).

A1a.4.7.5 Diadromous species

Sea trout is abundant in Welsh and other rivers, while sea lamprey, allis and twaite shads and smelt (*Osmerus eperlanus*) have all been recorded in the Dee estuary. The Solway Firth contains one of only two known spawning populations of allis shad in the UK (DECC 2010). Atlantic salmon is also present in various parts of the region (Potts & Swaby 1995). Four principal salmon rivers drain into Liverpool Bay (Clwyd, Conwy, Dee and Ribble), while the Mersey also supports a small population of salmon (Dong Energy 2013).

A1a.4.7.6 Shellfish

Nephrops is found in the soft sediments to the east and west of the Isle of Man as well as in the Firth of Clyde. The Irish Sea is an important region for scallops and queen scallops, with large grounds in Cardigan Bay, around the Isle of Man, the Solway Firth, Morecambe Bay and around islands in the Firth of Clyde. Cockles and oysters are also abundant throughout the region, particularly in the Solway Firth. Lobsters and brown crabs are abundant, particularly on the rocky shores of north Wales and the Lleyn Peninsula, while spider crabs are also present in the region (Clarke 1986). Whelks are also abundant in places, including around the Isle of Man and off the north Wales coast.

A1a.4.8 Features of Regional Sea 7

A1a.4.8.1 Fish community

The Minch and the firths along the west coast provide important sheltered, inshore nursery grounds for many species, with maerl beds an important habitat for large numbers of juvenile gadoids (Bailey *et al.* 2011). The fish assemblage in this region is comprised mainly of the gadoids haddock, hake and whiting and Norway pout as well as small pelagic species such as sprat (Gordon 1981). There is a high diversity of small benthic-demersal species including flatfish, gobies and blennies in the shallow, warm waters and sheltered sea lochs (Potts & Swaby 1997).

A1a.4.8.2 Pelagic species

A small number of mackerel migrate into the Minch after spawning in the autumn, although a change in the normal migration route in the 1980s has led to the western stock increasingly moving into the northern North Sea and the Norwegian Sea instead (Ellis & Heessen 2015a). Spent herring migrate into the Minch in winter following spawning. A number of spawning cohorts exist to the west of Scotland, with spawning taking place to the south-east of the Outer Hebrides, to the west of the Hebrides and to the north of Skye from August to September and again in the latter two sites in March and April. Larvae from the west coast spawning populations may be carried into the North Sea, mixing the stocks (Dickey-Collas *et al.* 2015). Sprat is very common in winter in the Minch. Sprat spawn over the summer and juveniles use the sea lochs and firths of the coastline as nursery areas (De Silva 1973).

A1a.4.8.3 Demersal species

Cod spawn to the north of the region in the early months of the year with peak activity in February. Whiting and Norway pout also spawn in the region. The Minch is an important nursery ground for juvenile gadoids, including cod, whiting, saithe, haddock and Norway pout. Haddock are widespread in the area throughout the year.

Plaice, lemon sole and sandeels use the region as spawning and nursery grounds and are present throughout the year. Other demersal species abundant in the region include bib, cuckoo wrasse (*Labrus mixtus*), common dragonet, butterfish, bullrout, pogge, topknot (*Zeugopterus punctatus*) and red and grey gurnards. A number of gobies and blennies, at the northern edges of their distributions, inhabit the lochs of the region, including the tompot blenny (*Parablennius gattorugine*), black goby (*Gobius niger*) and butterfly blenny (*Blennius ocellaris*) which are more typically found in waters to the south-west of the UK (Ellis & Rogers 2015, Ellis 2015d). The leopard spotted goby (*Thorogobius ephippiatus*) and Fries' goby (*Leseurigobius friesii*), reportedly rare in most UK waters, are frequently recorded in the sheltered sea lochs of the region. The former is a shy species that is rarely caught by remote methods, and is almost certainly not as rare as once thought. It is regularly observed by divers by its preferred habitat of small crevices or fissures (MacDougall 2002).

A1a.4.8.4 Elasmobranchs

The lesser spotted dogfish (but rarely the greater spotted dogfish), spurdog, common skate, cuckoo ray, nursehound (*Scyliorhinus stellaris*) and thornback ray have all been recorded in the Minch. The coastal waters of the west of Scotland are notable for the high abundance of basking sharks, particularly in the summer (Robson 1997). The areas around Hyskier, Coll and Tyree were identified as, "hotspots," by Witt *et al.* (2012), with satellite tracking revealing that individual sharks stayed close to the islands over the summers of 2012, 2013 and 2014 (Witt *et al.* 2016).

A1a.4.8.5 Diadromous species

Sea trout and salmon are abundant in the rivers and lochs of this region. Important lochs for these species include Loch Ewe, Loch Morar, Loch Shiel and Loch Lochy. Eels are likely to be present in the region. The twaite shad and allis shad have also been recorded in the region (Robson 1997).

A1a.4.8.6 Shellfish

The waters the Minch provide important habitats for *Nephrops*. The seabed in the region is soft and muddy and the species can be found in greatest abundance from the west of Skye to the Stanton Bank, between Islay and Jura and along the Kintyre Peninsula (Chapman 2006). *Nephrops* may also be found in the sea lochs around the coast of the mainland. Two species of shrimp are abundant in the region, *P. montagui* and the similar *Dichelopandalus bonnieri* (which typically has a more western distribution (Lee & Ramster 1981)). Scallops and queen scallops are abundant in sandy areas around the islands. Lobsters are present in greatest numbers around rocky areas of the coasts of the Hebridean Islands, while brown crabs are common throughout the region (Chapman 2006). Cockles are present, although individual cockle beds in the region are not large (Chapman 2006). Razor clams, whelks, periwinkles and velvet crabs are also abundant in the region and crawfish are present.

A1a.4.9 Features of Regional Sea 8

A1a.4.9.1 Fish community

The Scottish continental shelf is an ecosystem characterised by oceanic inflows from the Atlantic Ocean, such as the warm continental shelf current. The water is generally deeper than areas closer to the coast and the shelf edge is located at a depth of 300-450m. To the west of the Hebrides and around the north coast of Scotland, haddock, whiting, Norway pout, poor cod and grey gurnards dominate the assemblage. In deeper water along the shelf edge silvery pout, bluemouth (*Helicolenus dactylopterus*) and the hollowsnout rattail (*Caelorinchus caelorhincus*) are key members of the community (Pinnegar *et al.* 2010).

A1a.4.9.2 Pelagic species

Large numbers of mackerel move northwards along the continental shelf edge towards the northern North Sea in spring, having overwintered to the south-west of the UK. Consequently, this region is an area of mixing between the western and North Sea mackerel stocks with mackerel originating from both regions (Ellis & Heessen 2015a). Limited spawning takes place around Shetland and Orkney during the summer feeding migration, but the majority of the stock goes past the islands to feed in the deep waters at the shelf edge to the north-west of Shetland (Gordon 2006). However, during the return trip in autumn, most of the stock passes around Shetland. The western mackerel stock spawns from March to July in waters to the west of the Hebrides (Coull *et al.* 1998). Herring is locally abundant in the region, particularly when feeding in summer and autumn, and also around Orkney and to the south-west of Shetland during the late summer spawning period (Gordon 2006). To the west of the Hebrides, spawning takes place from March to April and again in late summer (Coull *et al.* 1998). Sprat is abundant with spawning occurring around the west and north coasts of the mainland. In some years, sprat is present around the coast of Shetland during migration, but their presence is irregular and unpredictable (Lee & Ramster 1981). Blue whiting is very abundant in deep waters to the north of Orkney in February, and spawning takes place between February and April along the continental slope to the west of Scotland at depths of 300-600m (Gordon 2006). After spawning, fish disperse to the North and Norwegian Seas to feed (Gordon 2006). Juvenile blue whiting remain on the nursery grounds for 2-4 years before returning to spawn (Gordon 2006). Argentines are abundant along the edge of the coastal shelf (Heessen 2015a).

A1a.4.9.3 Demersal species

Cod, haddock and whiting are all abundant in the region, with cod widely distributed off Orkney in the summer and whiting abundant in inshore waters. Spawning areas for these species exist to the west and north of the Hebrides and in the case of whiting, to the west of Shetland. Ling, pollack and saithe are also present, with saithe particularly abundant around Shetland. Norway pout is abundant in offshore regions and spawns around the shelf from January to April and further offshore a few months later (Defra 2005).

Plaice, dab and long rough dab are abundant in areas of sandy seabed, while megrim and witch are more abundant in the deeper waters of this region than elsewhere. Megrim is particularly abundant to the north-west of Orkney. The main area for witch in the North Sea is to the east of Orkney. Halibut (*Hippoglossus hippoglossus*), which are rare elsewhere, are frequently found off the coast of Shetland. Flounders spawn offshore in late winter. Monkfish and gurnards are also abundant in the area, with juvenile monkfish and non-spawning adults found in inshore regions of Orkney and Shetland. Sandeels are common in sandy areas across the region. A number of cold water species such as Vahl's eelpout (*Lycodes vahlii*), Esmark's eelpout (*Lycodes esmarkii*) and the wolf-fish (*Anarhichas lupus*) are present in the northern parts of the region (Pinnegar *et al.* 2010).

A1a.4.9.4 Elasmobranchs

The spurdog, lesser spotted dogfish, thornback ray and cuckoo ray are all abundant in the region. Common skate are locally abundant within certain sea lochs (Pinnegar *et al.* 2010). Porbeagle sharks are found along the edge of the continental shelf (Bendall *et al.* 2013), and genetic studies suggest they migrate south in the winter. They have been recorded, on occasion, as far out to sea as the oceanic ridge (Pade *et al.* 2009), and have been reported to gather south of Sumburgh Head in Shetland to give birth (Swaby & Potts 1997). The basking shark is frequently observed throughout this region, particularly west of the Hebrides. The common skate, listed as, "critically endangered," on the IUCN Red List, is relatively abundant in northern parts of the Scottish continental shelf, particularly around Shetland. In 2015, a rarely encountered false catshark (*Pseudotriakis microdon*) (colloquially, the sofa shark) was found off the Hebrides during a demersal survey, a reminder that open ocean and deep-sea fish may occasionally venture into waters near the coast.

A1a.4.9.5 Diadromous species

There are few rivers in the region to support populations of salmon. Lochs such as the Loch of Spiggie in Shetland, contain sea trout, while eels can be found in the small rivers of Orkney and the Hebrides.

A1a.4.9.6 Shellfish

The most important shellfish species on the Scottish continental shelf and slope is *Nephrops*, which is found on suitable sediments across the region down to depths of 600m (Chapman 2006). The shrimps *P. montagui* and *D. bonnierii* are also abundant. Lobsters and brown and velvet crabs are present in coastal regions, including the east of the Hebrides, the north of Scotland, Orkney and Shetland. Cockles, mussels and whelks are abundant around the coasts of Orkney and Shetland (Chapman 2006).

A1a.4.10 Features of Regional Seas 9, 10 & 11

A1a.4.10.1 Fish community

These Regional Seas may largely be characterised as offshore, deep water regions and consequently the fish assemblages found here are quite different from those of other regions. Many of the key pelagic and demersal species from Regional Sea 8 are still found in these regions, but there are fewer small, coastal species and a community of deep water species. This section will therefore concentrate on the deep water communities. The regions within this area can broadly be characterised as: Faroe-Shetland Channel (Regional Sea 9); Rockall Trough; and Rockall Bank and Plateau (Regional Sea 10). Regional Sea 11 comprises the Atlantic north-west approaches, where the water depth exceeds 1,000m. Deep-water fish tend to be slow-growing and long-lived. They often have a high age of maturation and the sparsity of life in the deep sea requires adult fish to aggregate at certain sites, usually seamounts, to spawn.

The Faroe-Shetland Channel is separated from the deep shelf edge waters to the west of Scotland by the Wyville Thomson Ridge, which rises to a depth of approximately 500m. This separation means that the fish communities on either side of it are quite distinct, particularly below 500m (Gordon 2001). At this depth, the water of the Faroe-Shetland Channel is cold and comprised mainly of Norwegian water, while to the west, water is of warmer Atlantic origin. The continental slope margin of the Faroe-Shetland Channel may be divided into three zones: an upper slope, the transition zone and the deep Norwegian Sea zone (Bullough *et al.* 1998).

A1a.4.10.2 Pelagic species

Blue whiting and mackerel may pass through the Faroe-Shetland Channel when migrating south to reach spawning grounds (Pinnegar *et al.* 2010). Herring are unlikely to be found in these deep seas, which are out of reach of their spawning grounds. They are also notably absent from the Rockall Bank (Dickey-Collas *et al.* 2015). Argentines, in contrast, aggregate at the Rockall Bank (Heessen 2015a).

Little is known about mesopelagic species. Some of these species can be very abundant and it is thought that the dominant fish species to the west of the UK are the light-emitting lantern-fish (*Notoscopelus kroyeri*) and the pearlside (*Maurolicus muelleri*) (Pinnegar *et al.* 2010, Kloppmann & Ellis 2015). The snipe-eel (*Nemichthys scolopaceus*) and the dragonfish (*Melanostomias bathyphilus*) are sometimes reported in mid-water surveys, while large ocean wanderers such as the dealfish (*Trachipterus arcticus*) and the oarfish (*Regalecus glesne*), the longest bony fish in the world, are occasionally washed up on UK coasts (Pinnegar *et al.* 2010).

A1a.4.10.3 Demersal species

Within the Faroe-Shetland Channel, the upper slope zone lies approximately between 200-500m depth and is characterised by Atlantic water with temperatures similar to those of the west coast. The dominant species found in this region are rabbitfish (*Chimaera monstrosa*), Norway redfish (*Sebastes viviparus*), bluemouth and blue whiting (Gordon & Swan 1997, cited by Gordon 2006). A similar community can be found at similar depths in Regional Seas 10 and 11, although redfish (*Sebastes* spp.) are less abundant to the west of the Hebrides (Gordon 2003).

The transition zone was identified by Bullough *et al.* (1998) and defined as the area of the slope where the bottom temperature changes rapidly with depth at the transition between the warm Atlantic water and the cold Norwegian Sea water. The main species identified by Bullough *et al.* (1998) were the Greenland halibut (*Reinhardtius hippoglossoides*) and roughhead grenadier (*Macrourus berglax*) (which are associated with cold water overflows (Gordon 2003)) along with blue ling (*Molva dypterygia*), tusk (*Brosme brosme*), two species of redfish (*S. marinus* and *S.*

mentella) and the Arctic skate (*Amblyraja hyperborea*). This assemblage is quite diverse and similar to that of the Norwegian continental shelf margin. A number of these species, such as the Greenland halibut and the roughhead grenadier, are rare to the west of the Wyville Thomson Ridge (Gordon 2003). Analysis of the results of MAFF (Ministry of Agriculture, Fisheries and Food) surveys carried out in the mid 1970s (Gordon and Swan 1997, cited by Gordon 2006) suggests that the transition zone in the Faroe-Shetland Channel occurs approximately between 625-785m depth.

The deep Norwegian Sea Water lies below the transition zone and is an area of low biomass and diversity. In surveys carried out by the Scottish Association for Marine Science (SAMS) in 1996 at depths of between 1,060m and 1,520m, a number of species of eelpout were identified as well as the Greenland halibut and Arctic skate.

Over 120 demersal fish species have been identified on the continental slopes to the west of Scotland below 400m depth (Gordon *et al.* 1994). There is no evidence of zonation in the fish assemblage in deep waters to the west of Scotland, such as is seen in the Faroe-Shetland Channel. The slope is characterised by a gradual decrease in temperature with depth and as each species occupies its own, highly variable, depth range, there is also a gradual change in the fish fauna with depth (Gordon 2003). Key members of the deepwater community in Regional Seas 10 and 11 include ling, blue ling (which spawn in these deep waters (Ellis *et al.* 2012)), tusk, roundnose grenadier (*Coryphaenoides rupestris*), argentines, black scabbardfish (*Aphanopus carbo*), orange roughy (*Hoplostethus atlanticus*) and greater forkbeard (*Phycis blennoides*).

Within the Rockall Trough, the dominant species at 250m depth are silvery pout, bluemouth and blue whiting, while in deeper waters a variety of morid cod (*Lepidion guentheri*), grenadiers, arrowtooth (or cut-throat) eels and deep water sharks dominate (Gordon & Bergstad 1992).

The Rockall Bank maintains a demersal fish community not dissimilar from that of the continental shelf, with the assemblage dominated by blue whiting, poor cod, Norway redfish, haddock and grey gurnard (Pinnegar *et al.* 2010). Other large gadoids on the Rockall Bank include cod, saithe, whiting and ling (Hislop *et al.* 2015). Flatfish include megrim (Velasco *et al.* 2015), witch, long-rough dab and lemon sole (Goldsmith *et al.* 2015). Monkfish (Ellis & Velasco 2015) and snake pipefish (Daan 2015b) are also common at the Rockall Bank.

A1a.4.10.4 Elasmobranchs

The Arctic skate is a key feature of the assemblage in the Faroe-Shetland Channel. There is a greater abundance of elasmobranchs to the west of Scotland than in the Faroe Shetland Channel with deepwater sharks such as the leafscale gulper shark (*Centrophorus squamosus*) and the blackmouth dogfish (*Galeus melastomus*) present (Gordon *et al.* 1994). The velvet belly (*Etmopterus spinax*), a lantern shark that uses light producing cells in its stomach as camouflage against the ocean surface, is sometimes caught in demersal hauls in the deep waters of the Rockall Plateau (Ellis & Heessen 2015b). On the Rockall Bank, thornback rays, common skate and the black-mouth dogfish are present (Ellis *et al.* 2015, Ellis 2015c). The mid-Atlantic skate (*Rajella kukujevi*) and the little gulper shark (*Centrophorus uyato*) have been caught on longlines in the Rockall Trough in recent years (Pinnegar *et al.* 2010).

A1a.4.10.5 Diadromous species

The area is distant from the coast and so diadromous species will only occur on migration and are unlikely to be encountered.

A1a.4.10.6 Shellfish

Although most of the species considered in this baseline tend to have a largely coastal, shallow water distribution, the Rockall Bank supports a range of crustacean and mollusc species and *Nephrops* may be found. The deep-sea red crab (*Chaecon affinis*) which inhabits seamounts throughout the east Atlantic is present in greatest numbers at depths between 600-900m and is targeted in fisheries in the region.

A1a.4.11 Evolution of the baseline

Recent research has suggested that there have been substantial changes in the fish communities in the north-east Atlantic over several decades. These communities consist of species that have complex interactions with one another and the natural environment. Fish species will undergo natural variation in population size, largely as a result of year to year variation in recruitment success. These population trends will be influenced by human exploitation and broad-scale climatic and hydrological variations.

A1a.4.11.1 Climate

As well as coming under severe pressure from anthropogenic factors, fish communities are likely to be affected by future climate change. Climate change may influence fish distribution and abundance through affecting growth rates, recruitment rates, behaviour, survival and responses to changes at other trophic levels, although exact responses are very difficult to predict. This could have a major effect on the community structure of the region. Sea surface temperatures (SST) of UK waters are predicted to rise by between 1.5-2.5°C this century (Lowe *et al.* 2009). Sea surface temperatures are driven by the North Atlantic Oscillation (NAO). Increased mortality in Atlantic salmon has been shown to be linked strongly to a positive phase in the NAO and the associated increase in sea surface temperature (Peyronnet *et al.* 2008).

An analysis of 50 species around the UK demonstrated that 70% changed distribution and abundance in response to warming between 1980 and 2008, with three-quarters of these species increasing in abundance (Simpson *et al.* 2011). Alheit & Hagen (1997) analysed data on herring and sardine landings at ports around the English Channel and southern North Sea dating back to the 10th Century. They found that large landings of herring were correlated with cold winter weather, while warm winters lead to large catches of sardine. Perry *et al.* (2005) showed marked changes in the distributions of North Sea fish over the past 25 years with the distributions of two-thirds of species having shifted in mean latitude. They found a northwards shift in the southern or northern population boundaries of a number of species had occurred. The northern boundary of bib, a southern North Sea species, was shown to have moved northwards by 342km between 1978 and 2001. Based on the projected SST increases of Hulme *et al.* (2002), Perry *et al.* (2005) predicted that the bib population may extend over the entire North Sea by 2080, while blue whiting may have retracted from the region by 2050. Drinkwater (2005) predicts that cod stocks in the Celtic and Irish Seas will have disappeared by 2100 as a result of temperature and hydrodynamic changes. Studies have shown that anchovy, which were absent from the North Sea until the mid 1990s, have extended their distribution as far north as the west coast of Norway (Brander *et al.* 2003, Alheit *et al.* 2012).

Habitat requirements are likely to play a significant role in vulnerability to climate change, with species such as plaice and herring likely to be vulnerable at different stages in their life-cycles (nursery grounds and spawning grounds, respectively (Petitgas *et al.* 2013)). Increasing freshwater temperatures over the past four decades have implications for the survival rates of juvenile diadromous fish, although not necessarily negative ones (Simpson *et al.* 2013). Although salmon populations in Scottish rivers have been declining since the 1990s, partly due to poor growth of juveniles at sea, twaite shad (Arahamian *et al.* 2010) and sea lamprey (Rodríguez-Muñoz *et al.* 2001) demonstrate increased larval survival at warmer temperatures.

Shellfish populations are often strongly tied to particular sediment types and so these too will be vulnerable to climate-induced impacts on their preferred habitat. The settlement of many bivalve species (e.g. cockles, mussels) is dependent on a range of environmental factors (Chapman 2004). As a result, significant changes in the environment may cause low stock recruitment and potentially the disappearance of a species from a ground. Changing environmental conditions may also affect the distributions of migratory crustaceans, such as lobsters and crabs.

Hedger *et al.* (2004) found that North Sea aggregations were present in deeper water in the 1990s than the 1980s, while a study of distribution data from 1980-2004 conducted by Dulvy *et al.* (2008) found that the North Sea demersal fish assemblage had deepened by 3.6m per decade over this time. However, modelling carried out by Rutterford *et al.* (2015) suggests that demersal fish are constrained by the availability of habitat at a suitable depth. Consequently, the capacity of fish to adapt to increasing water temperature by relocating to deeper water may have already been exhausted and little or no further movement to deeper water is possible. Unable to move into deeper, cooler water, fish will be exposed to higher temperatures, with potential physiological and ecological consequences. Evidence of the physiological stresses experienced by cod has been demonstrated by studies of the otoliths (ear bones), which reveal slower growth rates during warmer years (Pilling *et al.* 2007, Millner *et al.* 2011). Sandeels are also unlikely to be able to migrate easily into deeper waters, as they are reliant on a coarse, sandy seabed habitat (Holland *et al.* 2005). Declining recruitment in sandeel populations around northern UK waters is inversely correlated with increasing temperature (Heath *et al.* 2012). Species such as sandeel, a key link between trophic levels in the food web, that are unable to adapt to changing temperatures as a result of specific habitat requirements are likely to be most vulnerable to the effects of climate change. Declining recruitment in the sandeel population of the northern UK is inversely correlated with sea temperature (Heath *et al.* 2012).

Species richness in the North Sea has increased, with 8 times more species increasing their distributions than have decreased their distributions (Hiddink & ter Hofstede 2008 (see also Simpson *et al.* 2011)). Species richness shows a positive correlation to average bottom temperature (Henderson 2007), and it is predicted that a 2°C rise in sea bottom temperature would result in an increase in species richness in the Bristol Channel of 10% (Henderson 2007, Henderson & Bird 2010). The increase in species richness, along with an increase in small, southerly species is possibly influenced by the release of predation pressure from large, exploited commercial fish, acting in combination with the changing climate (Hiddink & ter Hofstede 2008). Changes in species composition may mean a change in the community structure. Twelve fish species which were classified as typical southern North Sea species by Corten & van de Kamp (1996) were observed by Beare *et al.* (2004) to have increased in abundance dramatically in the northern North Sea since the mid 1990s. Vaz *et al.* (2007) detected an increase in species diversity in the English Channel community between 1997 and 2002, and sightings of a number of warm water fish such as blue-fin tuna (*Thunnus thynnus*), grey triggerfish (*Balistes caprisacus*) and large pelagic sharks have increased (Stebbing *et al.* 2002). The distribution range of the short-snouted seahorse seems to be moving northwards from the English Channel and is increasingly recorded in the southern North Sea (Pinnegar *et al.* 2008).

The Celtic and Irish Seas, have seen increases in the distribution and abundance of warm water species such as the John Dory, and species richness increased between 1999 and 2008. In contrast, waters off the west of Scotland saw a reduced number of species over this period (ter Hofstede *et al.* 2010). There is, however, significant spatial variation in these general trends; increases in richness can be observed at smaller spatial scales within the region, for example off the north coast of Ireland (Heath & Speirs 2012). Analysis of trawl survey results from 1986-

2008 reveal the Rockall Bank to have been colonised by whiting and the oceanic, southern species Ray's sea bream (*Brama brama*) in recent years (Neat & Campbell 2011).

Dippner (1997) found a high correlation between SST and variability in the recruitment of cod, whiting and mackerel in the North Sea. A reduction in the survival of cod larvae (and consequently, recruitment) in the North Sea since the mid 1980s has been linked to a change in the copepod community and in particular to the decline of *C. finmarchicus*, an important prey item for larval cod (Beaugrand *et al.* 2003). Similarly, shifts in the *Calanus* species composition linked to climate (see Section A1a.1) has led to a mismatch in timing of the occurrence of the early larval stages of sandeel and food availability, a factor behind recent recruitment failure (van Deurs *et al.* 2009). The large-scale distribution of basking sharks in recent years has been positively correlated with sea temperature, a pattern that may be linked to the distribution of the warm water copepod *C. helgolandicus*, the abundance of which is indicative of shark distribution over small spatial scales (Cotton *et al.* 2005).

A changing climate may affect migration routes of some species. There has been a northerly shift in the mackerel spawning grounds and a change in the timing of adult migration into these grounds with the result that fewer mackerel now pass through the Minch during their migration (Walsh *et al.* 1995). Flounder has been observed to undertake spawning migrations up to two months earlier in cooler conditions (Sims *et al.* 2004), while the navigation of salmon back to home rivers may be severely affected as it relies on a range of environmental cues, potentially affecting recruitment success (Pinnegar *et al.* 2008). The timing of spawning may also be affected, with Greve *et al.* (2001) noting the earlier appearance of fish larvae in the southern North Sea in recent years. Kjesbu *et al.* (2010) report that temperature increases may act to bring forward the onset of spawning in bass. In addition, female bass have been found not to reach maturity unless they remain in water above 10°C during maturation (Pawson *et al.* 2000), which may, in part, explain the increased abundance in sea bass in the southern North Sea and Channel. Fincham *et al.* (2013) report that four out of seven stocks of sole showed significant long-term trends towards earlier spawning (at a rate of 1.5 weeks per decade), with winter SST significantly affecting the date of peak spawning. In addition, any effects climate change may have on larval dispersion are currently a major unknown factor in predicting the effects of climate change (Petitgas *et al.* 2013). As described in Section A1a.4.2.1, Fässler *et al.* (2011) have demonstrated how the timing of spawning in herring may influence larval dispersal and development.

Lenoir *et al.* (2011) attempted a forecast of the distributions of eight fish species in the north-east Atlantic, using ecological niche modelling with explanatory variables including SST, salinity and bathymetry under various greenhouse gas emission scenarios. The forecast concluded that by 2090 horse mackerel and anchovy would show increased occurrence in northern UK waters, while pollack, sole, haddock and saithe would show a decrease in southern UK waters; turbot and sprat would show no significant change. However, climate change effects on marine fish populations are difficult to predict and the evidence is not easy to interpret. As an example, from 2003, the snake pipefish, previously rare in the region, showed a sudden and dramatic increase in northern European waters, particularly to the west of Scotland and the northern North Sea (Harris *et al.* 2007) and could be found as far north as Svalbard (Fleischer *et al.* 2007). This sudden expansion had significant impacts on seabird populations as chicks, used to a diet of sandeels, struggled to eat the bony, awkward prey item (Harris *et al.* 2007). Numbers peaked in 2007 and this was considered by many to be indicative of a major marine ecosystem shift in response to climate change. Since 2007, however, catch rates of snake pipefish have dwindled back to their pre-2003 average, with the sudden outburst remaining unexplained (Heath *et al.* 2012, Daan 2015b). A similar situation arose when large catches of bluemouth, generally a rare species in the North Sea were reported near Shetland in 1991, spreading over the entire North Sea within a few years (Heessen & Blasdale 2015). By 1997,

however the species had almost completely disappeared again. It is thought that the fish all belonged to a single cohort, perhaps swept into the North Sea by an abnormally large inflow of Atlantic water that year (Mamie *et al.* 2007). These examples serve as a reminder that the factors governing fish distribution and abundance are complicated and interdependent.

A1a.4.11.2 Overfishing

Many fish are subject to considerable fishing pressure, which will act to reduce biomass of commercially valuable species and, potentially, non-target species. Fishing may affect the abundance, diversity, size, composition and life-history of fish communities, through the pressure exerted and the selective removal of larger, mature individuals from populations. The impacts of long-term exploitation of a fish stock are typically a decrease in body size, age of maturation and productivity. The benefits of sustainable management of fish stocks can be seen in the herring population. Following a collapse in the early 1970s, the spawning stock biomass of this species has risen to healthy levels, with a short moratorium (1977-1981) and subsequent technical measures ensuring careful management and sustainable harvesting (ICES 2008).

Data indicate that the biomass of fish from high trophic levels declined by two thirds in the North Atlantic in the second half of the 20th Century (Christensen *et al.* 2003). Thurstan *et al.* (2010) showed that over 118 years from 1889-2007, the landings per unit of fishing power in the bottom trawl sector fell by 94%, a demonstration of the scale of the decline in demersal fish stocks since the advent of industrial fishing. Anderson *et al.* (2008) and Perry *et al.* (2010) argue that the de-stabilisation of fish stocks caused by heavy fishing leads to increased fluctuations in abundance and therefore increased vulnerability to natural events. This was also illustrated by Lindegren *et al.* (2010) who demonstrated the increased resilience to environmental change of the Atlantic cod in the Sound, the narrow strait separating the North and Baltic Seas, where a trawling ban has been in place since 1932, compared to the neighbouring waters.

Consequently, fishing pressure is likely to enhance the impacts of climate change. Off the south-west of Britain, the pattern generally observed is for warm-water species to have increased in abundance in recent decades. However, Genner *et al.* (2004, 2010) showed that warm-water species of commercial importance, such as skates, rays and brill have declined over the same period. The conclusion drawn from this was that species under fishing pressure may be unable to respond to the more favourable environmental conditions. Modelling carried out by Beggs *et al.* (2013) suggests that cod is more sensitive to climate variability during periods of low spawning stock biomass.

Bailey *et al.* (2009a) showed that fish abundance recorded from depths of 800-2,500m in the Porcupine Seabight and Abyssal Plain (south-west of Ireland), has fallen significantly since 1977, possibly as a result of impacts on the species in shallower waters resulting in declines in the deeper parts of the range. These depths are considerably deeper than the maximum depth of commercial fishing (approximately 1,600m), indicating that the effects of fishing extend into deep, un-fished waters, where there is little or no routine monitoring or management. The tendency of deep-sea fish to form spawning aggregations makes them an easy target for trawlers, while their slow growth rates and long life expectancies make them highly vulnerable to over-fishing.

The latest Charting Progress report (Defra 2010) states that the majority of UK stocks are still fished well above the levels expected to provide the highest long-term yield, although of 20 indicator stocks, the proportion being harvested sustainably rose from 10% in the early 1990s to about 40% in 2007. Fishing mortality estimates have declined significantly in recent years in 67% of assessed stocks in UK waters, and recent ICES advice suggests a greater degree of

optimism about the future of these stocks than has been present for a number of years (leading to an increase in the quotas for several species in 2016). The OSPAR Quality Status Report 2010 (OSPAR 2010) concluded that too many fish stocks are still outside safe biological limits, although there have been significant improvements in a number of stocks. Recent reductions in fishing effort have been partially offset by a more technologically advanced and efficient fleet (OSPAR 2010).

Recent assessments of indicator stocks suggest that 4 of 13 are at full reproductive capacity and being harvested sustainably (Defra 2015), an improvement on the average between 1990 and 2009. Of these four stocks, three (North Sea plaice, north-east Atlantic mackerel and blue whiting) were also being fished at or below the rate providing maximum sustainable yield (MSY), indicating that harvesting of these stocks is both sustainable and delivering the largest possible catches. Fernandes & Cook (2013) meanwhile report a substantial reduction in exploitation rate since the turn of the century, with 22 out of 36 assessed stocks within EU waters fished below MSY, compared to 10 in 2002.

A1a.4.12 Environmental issues

A1a.4.12.1 Trophic interactions

Fish and shellfish are important components of marine ecosystems, operating at a number of trophic levels. They utilise a variety of feeding strategies, including filter feeding for plankton and detritus suspended in the water column, scavenging for detritus on the seabed, and both pelagic and demersal predation of plankton, small fish, cephalopods, crustaceans and other benthic organisms. Pelagic fish such as mackerel and herring primarily feed on planktonic crustaceans, other zooplankton and small fish. Demersal fish such as gadoids and flatfish often consume a wide range of benthic invertebrates including crustaceans, polychaetes, molluscs and echinoderms, along with cephalopods and fish. Sandeels main prey is calanoid copepods, but other planktonic prey including fish larvae are also taken; larger individuals may also take benthic prey such as polychaetes. Most benthic crustaceans are scavengers to some extent, feeding on detritus, although many species are also active predators of a variety of benthic organisms. Many molluscs such as scallops, cockles and mussels are filter feeders of material suspended in the water column.

Fish and shellfish play a pivotal role in the transfer of energy from some of the lowest to the highest trophic levels within the ecosystem, and also through the recycling of nutrients from higher levels through the consumption of detritus. Consequently, their populations will be determined by both top-down factors, such as predation, and bottom-up factors, such as ocean climate and plankton abundance. Fish and shellfish are important prey items for top marine predators including elasmobranchs, seabirds, cetaceans and humans, and small planktivorous species such as sandeels and herring act as important links between zooplankton and top predators (Frederiksen *et al.* 2006). The influence that fish abundance can have on predators is best illustrated by sandeels, which have declined in abundance dramatically in recent years, with sandeel landings in the North Sea decreasing by over 50% since 2003 (Frederiksen *et al.* 2006). This decline has coincided with a series of breeding failures amongst sandeel-dependent seabirds such as puffins (*Fratercula arctica*) and kittiwakes (*Rissa tridactyla*). Sandeel recruitment and their interactions with higher predators are complex and poorly understood. Further information on the ecological importance of fish and shellfish species to seabirds and marine mammals is provided in Appendices A1a6 and A1a7 respectively.

A1a.4.12.2 Habitat impacts

Many fish and shellfish species are dependent on specific habitats either throughout, or at particular stages of their lives. Herring spawn on coarse sand or gravel, while sandeels live on sandbanks and *Nephrops* require muddy sediments in which they can build burrows.

Seahorses and many other cryptic species find refuge in seagrass beds of the south coast, while maerl beds provide sheltered nursery grounds for juvenile gadoids off the west of Scotland. Diadromous fish are dependent on passage through shallow coastal waters, into and up rivers to spawn (or in the opposite direction for eels). Damage through contact with anchors, fishing gear and offshore or nearshore construction can all impact on fish communities.

Stelzenmüller *et al.* (2010) present maps of the sensitivity of 11 fish and shellfish species to aggregate extraction around the southern and western coasts of England and Wales. The maps indicate that the highest sensitivity is typically in coastal regions where spawning and nursery areas are found. The most sensitive species to extraction were scallop, lobster and queen scallop with whiting and plaice the least sensitive. It is estimated that it would take between 2-3 years after a cessation of aggregate extraction for signs of community recovery to show in sandy gravel habitats with moderate wave exposure and tidal currents (Defra 2010).

A1a.4.12.3 Contamination

A significant source of contamination is that of riverine discharge, which may result in toxic algal blooms which can result in fish and shellfish kills and shellfishery closures. Hormonal disruption of flounder through oestrogenic contamination has been observed around UK estuaries (Allen *et al.* 1999). Hormonal disruption through contamination also occurs in gastropod and bivalve shellfish. Female whelks and dog whelks (*Nucella lapillus*) exposed to tributyltin (TBT), an antifoulant used on the hulls of boats until 1987, have been reported to develop male sex organs (Polockzanska & Ansell 1999). This condition, called imposex, can have significant impacts on the reproductive success of a population. Since the ban on the use of TBT, monitoring of dog whelks at sites around Scotland have shown the incidence and severity of cases of imposex to be much reduced (Scottish Government website⁸)

The effects of contamination from oil on demersal fish have been assessed in a number of species both in the field and in the laboratory. A number of indicators of hydrocarbon contamination, such as the activity of certain enzymes in the liver, may be used to investigate this. Within two weeks of the *Sea Empress* spill off the coast of Pembrokeshire, Kirby *et al.* (1999) collected dab and plaice samples from near to the site. They found elevated hepatic enzyme activity, providing evidence of significant levels of hydrocarbon contamination. Aas & Klungsøyr (1998) concluded that cod and haddock were not significantly affected by locally elevated poly-aromatic hydrocarbon (PAH) concentrations surrounding platforms in Norwegian waters.

Micro-plastics in the marine environment may result from the break-down of plastic waste over time, or from specifically manufactured micro-beads or fibres typically found in products such as toothpaste or exfoliants. Analyses of micro-plastics recorded from Continuous Plankton Recorder (CPR) samples indicate that the quantity is increasing over time in the north-east Atlantic, with greatest concentrations found at the coasts (Edwards *et al.* 2014). It has been demonstrated that micro-plastics may be ingested by zooplankton and will be passed to higher trophic levels by predation on affected zooplankton. Desforges *et al.* (2015) estimated that adult salmon in the north-east Pacific could be ingesting up to 90 particles each day.

⁸ <http://www.gov.scot/Publications/2011/03/16182005/33>

A1a.4.12.4 Noise disturbance

Fish exhibit large variation in their ability to emit and detect sound, dependent on diversity in anatomical features, hearing physiology and behaviour. For example, drumming noises produced by the swimbladder have been found to have an important role in mating of cod, haddock and probably other gadoids (Rowe & Hutchings 2006). Staaterman *et al.* (2014) made the first acoustic recordings of sounds produced by fish larvae (those of the grey snapper (*Lutjanus griseus*)), while the larvae of some coral reef fish and shellfish have been shown to navigate to suitable settling locations on the reef using sound signals (Stanley *et al.* 2012). This suggests that even at this young, under-developed stage, fish are sensitive to acoustic signals in the marine environment. A study conducted by Buscaino *et al.* (2010) exposed European sea bass and gilthead sea bream (*Sparus aurata*) to acoustic stimuli of a frequency typical of vessel traffic, with the noise resulting in increased motility and intense muscle activity in both species.

Research on the effects of anthropogenic noise on marine fish and invertebrates continues to improve our understanding of this potential issue. Nevertheless, significant data gaps remain and the difficulty of designing and conducting suitable experiments makes it hard to draw firm conclusions on the impacts that activities such as seismic surveys and pile-driving might have on fish and invertebrate species (Hawkins *et al.* 2015). Mueller-Blenkle *et al.* (2010) played recordings of pile-driving noise to captive cod and sole, and observed the responses of the two species. Both species displayed significant, but different, behavioural responses at relatively low sound pressure levels. Cod showed a freezing response at the onset and cessation of the noise stimulus, while sole showed an increase in swimming speed for the duration of the stimulus. Both species made movements away from the source of the noise. Pile-driving stimuli were shown to result in a significant increase in barotrauma injuries to Chinook salmon (*Oncorhynchus tshawytscha*) (Halvorsen *et al.* 2012). The repetitive, intense pulses of sound characteristic of seismic survey air guns were found to cause long-term damage to the epithelial cells of the pink snapper (*Pagrus auratus*), with the injuries sustained resulting in the fish being unable to respond to a similar stimulus in a repeat of the experiment 58 days later (McCauley *et al.* 2003). However, a similar experiment conducted in several species of riverine fish by Popper *et al.* (2005) found that the limited hearing loss incurred had recovered with 24 hours, with no apparent damage to hair cells.

Acoustic disturbance on early life stages is also an area of significant interest. Laboratory experiments carried out by Simpson *et al.* (2015) found that predator avoidance by juvenile European eels migrating to the continent from the Sargasso Sea was significantly impaired by the noise of vessel traffic in harbours. The migration route of older eels back to spawning grounds does not appear to be affected by noise generated by offshore wind farms in the southern Baltic Sea (Andersson *et al.* 2012).

A1a.4.12.5 Electromagnetic Field disturbance

Many fish species are able to detect electric and magnetic fields (EMF) and it is a matter of concern that anthropogenic sources of EMF, in particular subsea cables, might be detected by and negatively influence sensitive species. Elasmobranchs possess an electrosensory system called the Ampullae of Lorenzini, an array of receptors that allow them to detect the weak electric fields produced by prey items. Consequently, they are most frequently linked with potential EMF effects. Anthropogenic EMF may affect elasmobranchs during seasonal migrations, feeding and reproductive behaviour or normal habitat use. An additional, untested area of concern is the potential effects of subsea cables on juvenile sharks and rays in coastal nursery areas.

Available data assessing elasmobranch responses to subsea cable EMFs is limited. The first and only documented example of a marine fish responding to an emission from a subsea cable comes from evidence of shark bites on optical telecommunication cables (Marra 1989). Gill *et al.* (2009) demonstrate that the behaviour of a number of species of elasmobranchs are affected by EMFs from undersea cables. The authors indicated that the effect was only likely to be observed within a very small distance around the cable and it was unclear whether the effect was positive, negative or neutral.

Some teleost fish have been demonstrated to be EMF sensitive. Relevant species in UK waters include plaice and the diadromous species European eel, Atlantic salmon, sea trout and river and sea lamprey (Gill & Bartlett 2010). Migratory eels and salmonids are able to use the earth's magnetic field for orientation during migrations (Durif *et al.* 2013) and magnetic field detection has been demonstrated in a range of fish species, notably by Formicki *et al.* (2004), who found a number of species to be more attracted to fyke nets on which a magnet was mounted than to nets with no magnet. While there is a range of evidence to support the EMF detection abilities of these fish, and little doubt that highly mobile, migratory species will encounter anthropogenic EMF sources, evidence is limited as to whether such EMFs are likely to provoke a behavioural response (Ohman *et al.* 2007, Gill & Bartlett 2010).

A1a.4.12.6 Conservation frameworks

In addition to fisheries management measures, a number of conservation frameworks apply to fish species within UK waters (see Table A1a.4.2). The *Wildlife and Countryside Act 1981 (as amended)* applies to territorial waters and lists (Schedule 5) several species of marine and estuarine fish which receive protection under the Act. The Act makes it an offence (subject to exceptions) to kill, injure, or otherwise disturb any wild animal listed on the schedule and prohibits interference with places used for shelter or protection by these species. The Act does not apply in Northern Ireland, where the equivalent is the Wildlife (Northern Ireland) Order 1985 (although no marine fish are covered by this Act). European Protected Species (Annex IV species) receive protection via the various UK Habitats Regulations. The 2007 amendments to these Regulations resulted in consequential amendments to the Wildlife and Countryside Act with respect to whole or partial removal from Schedule of the Act 5 of European Protected Species to ensure consistency of regulation. There are six diadromous species that require the designation of SACs (Special Area of Conservation) in UK waters under Annex II of the EC Habitats Directive, although the salmon is only protected in freshwater habitats. Other diadromous species are protected under Annex IV or V, which offer protection against deliberate disturbance, capture or killing.

As well as this, a number of species have been listed under Annex V of the OSPAR list of threatened and/or declining species and habitats and by CITES (Convention on International Trade in Endangered Species), an international agreement between governments to regulate international trade in wild animals and plants. A number of species are also listed on the IUCN (International Union for the Conservation of Nature) Red List of Threatened Species. A number of fish species are also the subject of UK Biodiversity Action Plans (BAPs), as priority species. In addition to, individual lists exist for Scottish, Welsh and Northern Irish species and there is some variation between these lists.

A European Council Regulation (EC No 1100/2007) establishing measures for the recovery of European eel stocks entered into force in September 2007. The regulation requires the development of Eel Management Plans (there are 15 such Management Plans in the UK) which include measures to allow the escapement to sea of at least 40% of the silver eel biomass. Under the regulation, from 2013 EU countries with a glass eel fishery will be required to reserve 60% of the eel catches with a body length less than 12 cm for restocking European eel river

basins. European eel was included in CITES Appendix II in 2007 and in 2008 was listed as “Critically Endangered” on the IUCN Red List.

Table A1a.4.2: Marine fish afforded protection under national legislation and international conventions

Species	Wildlife and Countryside Act ¹	EC Habitats Directive	OSPAR	CITES	IUCN	Priority Species ²
Pelagic fish						
Mackerel (<i>Scomber scombrus</i>)					Least concern	England, Scotland (PMF), Wales, NI
Herring (<i>Clupea harengus</i>)					Least concern	England, Wales, Scotland (PMF), NI
Horse mackerel (<i>Trachurus trachurus</i>)					Vulnerable	England, Scotland (PMF), Wales, NI
Blue whiting (<i>Micromesistius poutassou</i>)						England, Scotland (PMF)
Northern bluefin tuna (<i>Thunnus thynnus</i>)			Y		Endangered	England
Demersal species						
Atlantic cod (<i>Gadus morhua</i>)			Y		Vulnerable	England, Wales, Scotland (PMF), NI
Whiting (<i>Merlangius merlangus</i>)					Least concern	England, Wales, Scotland (PMF), NI
Saithe (<i>Pollachius virens</i>)						Scotland (PMF)
Hake (<i>Merluccius merluccius</i>)						England, Wales, Scotland, NI
Ling (<i>Molva molva</i>)						England, Wales, Scotland (PMF), NI
Blue ling (<i>Molva dypterygia</i>)						England, Scotland (PMF)
Tusk (<i>Brosme brosme</i>)						Scotland
Norway pout (<i>Trisopterus esmarkii</i>)					Least concern	Scotland (PMF)
Monkfish (<i>Lophius piscatorius</i>)					Least concern	England, Scotland (PMF), Wales, NI

Species	Wildlife and Countryside Act ¹	EC Habitats Directive	OSPAR	CITES	IUCN	Priority Species ²
Black scabbardfish (<i>Aphanopus carbo</i>)						England, Scotland (PMF)
Roundnose grenadier (<i>Coryphaenoides rupestris</i>)					Critically endangered	England, Scotland (PMF)
Norway redfish (<i>Sebastes viviparus</i>)						Scotland
Plaice (<i>Pleuronectes platessa</i>)					Least concern	England, Wales + Scotland, NI
Sole (<i>Solea solea</i>)					Data deficient	England, Wales, NI
Deepwater sole (<i>Bathysolea profundicola</i>)					Least concern	Scotland
Atlantic halibut (<i>Hippoglossus hippoglossus</i>)					Endangered	England, Scotland (PMF)
Greenland halibut (<i>Reinhardtius hippoglossoides</i>)						England, Scotland (PMF)
Giant goby (<i>Gobius cobitis</i>)	Schedule 5					
Couch's goby (<i>Gobius couchii</i>)	Schedule 5				Least concern	
Sandeel (<i>Ammodytes marinus</i>)						England, Wales, Scotland (PMF)
Sandeel (<i>Ammodytes tobianus</i>)					Data deficient	Scotland (PMF)
Short snouted seahorse (<i>Hippocampus hippocampus</i>)	Schedule 5 (England only)		Y	Appendix II	Data deficient	England
Long snouted seahorse (<i>Hippocampus guttulatus</i>)	Schedule 5 (England only)		Y	Appendix II	Data deficient	England, Wales
Orange roughy (<i>Hoplostethus atlanticus</i>)			Y			England, Scotland (PMF)
Elasmobranch species						
Basking shark (<i>Cetorhinus maximus</i>)	Schedule 5		Y	Appendix II	Vulnerable	England, Wales, Scotland (PMF), NI
Tope (<i>Galeorhinus galeus</i>)					Vulnerable	England, Wales, NI

Species	Wildlife and Countryside Act ¹	EC Habitats Directive	OSPAR	CITES	IUCN	Priority Species ²
Spurdog (<i>Squalus acanthias</i>)			Y		Vulnerable	England, Scotland (PMF), Wales, NI
Porbeagle (<i>Lamna nasus</i>)			Y	Appendix II	Vulnerable	England, Scotland (PMF), Wales, NI
Blue shark (<i>Prionace glauca</i>)					Near threatened	England, Wales
Shortfin mako (<i>Isurus oxyrinchus</i>)					Vulnerable	England
Angel shark (<i>Squatina squatina</i>)	Schedule 5 (England only – not between 6-12nm of coast)		Y		Critically endangered	Wales, NI
Gulper shark (<i>Centrophorus granulosus</i>)			Y			England
Leafscale gulper shark (<i>Centrophorus squamosus</i>)			Y		Vulnerable	England, Scotland (PMF)
Portuguese dogfish (<i>Centroscymnus coelolepis</i>)			Y		Near threatened	England, Scotland (PMF)
Kitefin shark (<i>Dalatias licha</i>)					Near threatened	England
Common skate (<i>Dipturus batis</i>)			Y		Critically endangered	England, Wales, Scotland (PMF), NI
Spotted ray (<i>Raja montagui</i>)			Y		Least concern	
Sandy ray (<i>Leucoraja circularis</i>)					Endangered	Scotland (PMF)
Undulate ray (<i>Raja undulata</i>)					Endangered	England, Wales, NI
White skate (<i>Rostroraja alba</i>)			Y		Endangered	England, Wales
Thornback ray (<i>Raja clavata</i>)			Y		Near threatened	Wales, Scotland
Diadromous species						
European sturgeon (<i>Acipenser sturio</i>)	Schedule 5 (not Scotland)	Annex II & IV	Y	Appendix I	Critically endangered	England, Wales, Scotland
Allis shad (<i>Alosa alosa</i>)	Schedule 5	Annex II & V	Y		Least concern	England, Wales, Scotland, NI

Species	Wildlife and Countryside Act ¹	EC Habitats Directive	OSPAR	CITES	IUCN	Priority Species ²
Twaite shad (<i>Alosa fallax</i>)	Schedule 5	Annex II & V			Least concern	England, Wales, Scotland, NI
River lamprey (<i>Lampetra fluviatilis</i>)		Annex II & V			Least concern	England, Wales, Scotland (PMF), NI
Sea lamprey (<i>Petromyzon marinus</i>)		Annex II	Y		Least concern	England, Wales, Scotland (PMF)
Smelt (<i>Osmerus eperlanus</i>)					Least concern	England, Wales, Scotland (PMF), NI
Whitefish (<i>Coregonus lavaretus</i>)	Schedule 5	Annex V	Y		Vulnerable	England, Wales, Scotland
European eel (<i>Anguilla anguilla</i>)			Y	Appendix II	Critically endangered	England, Wales, Scotland (PMF), NI
Atlantic salmon (<i>Salmo salar</i>)		Annex II (freshwater only)	Y		Lower risk / least concern	England, Wales, Scotland (PMF), NI
Sea trout (<i>Salmo trutta</i>)					Least concern	England, Scotland (PMF), Wales, NI
Commercial Shellfish species						
Crawfish (<i>Palinurus elephas</i>)					Vulnerable	England, Scotland (PMF), Wales, NI
Native oyster (<i>Ostrea edulis</i>)			Y			England, Wales, Scotland (PMF), NI
Horse mussel (<i>Modiolus modiolus</i>)						NI

Note: ¹Does not apply to Northern Ireland and may vary between England, Wales and Scotland. ²Priority species include those listed under Section 41 (England) and Section 42 (Wales) of the Natural Environment and Rural Communities (NERC) Act 2006, Section 2(4) of the Nature Conservation (Scotland) Act 2004, Section 3(1) of the Wildlife and Natural Environment Act (Northern Ireland) 2011 and Schedule 5 of the Wildlife (Northern Ireland) Order 1985, also includes species on Scotland's list of Priority Marine Features (PMF).

Sources: Information sourced January 2016.

References

- Aas E & Klungsøyr J (1998). PAH metabolites in bile and EROD activity in North Sea fish. *Marine Environmental Research* **46**: 229-232.
- Albert OT (1994). Ecology of haddock in the Norwegian Sea. *ICES Journal of Marine Science* **51**: 31-44.
- Alheit J & Hagen E (1997). Long-term climate forcing of European herring and sardine populations. *Fisheries Oceanography* **6**: 130-139.
- Alheit J, Pohlmann T, Casini M, Greve W, Hinrichs R, Mathis M, O'Driscoll K, Vorberg R & Wagner C (2012). Climate variability drives anchovies and sardines into the North and Baltic Seas. *Progress in Oceanography* **96**: 128-139.
- Allen Y, Matthieson P, Scott AP, Haworth S, Feist S & Thain JE (1999). The extent of oestrogenic contamination in the UK estuarine and marine environments – further surveys of flounder. *The Science of the Total Environment* **233**: 5-20.
- Amara R, Poulard JC, Lagardère F & Desaunay Y (1998). Comparison between the life-cycles of two Soleidae, the common sole, *Solea solea*, and the thickback sole, *Microchirus variegatus*, in the Bay of Biscay (France). *Environmental Biology of Fishes* **53**: 193-209.
- Anderson CNK, Hsieh C, Sandin SA, Hewitt R, Hollowed A, Beddington J, May RM & Sugihara G (2008). Why fishing magnifies fluctuations in fish abundance. *Nature* **452**: 835-839.
- Andersson MH, Lagenfelt I & Sigray P (2012). Do ocean based windfarms alter the migration pattern in the endangered European silver eel (*Anguilla Anguilla*) due to noise disturbance? In: Popper AN & Hawkins A (2012). *The effects of noise on aquatic life. Advances in Experimental Medicine and Biology* **730**: 393-396.
- Aprahamian M & Robson CF (1998). Chapter 5.8. Fish: salmon, sea trout and eels. In: JH Bame, CF Robson, SS Kaznowska, JP Doody, NC Davidson & AL Buck Eds. *Coasts and seas of the United Kingdom. Region 7 South-east England: Lowestoft to Dungeness (Coastal Directory Series)*. Joint Nature Conservation Committee, Peterborough, pp. 126-127.
- Aprahamian MW, Aprahamian CD & Knights AM (2010). Climate change and the green energy paradox: the consequences for twaite shad *Alosa fallax* from the River Severn, UK. *Journal of Fish Biology* **77**: 1912-1930.
- Arnold GP & Metcalfe JD (1996). Seasonal migrations of plaice (*Pleuronectes platessa*) through the Dover Strait. *Marine Biology* **127**: 151-160.
- Bailey DM, Collins MA, Gordon JDM, Zuur AF & Priede IG (2009). Long-term changes in deep water fish populations of the Northeast Atlantic: a deeper reaching effect of fisheries. *Proceedings of the Royal Society B* **276**: 1965-1969.
- Bailey M, Bailey DM, Bellini LC, Fernandes PG, Fox C, Heymans S, Holmes S, Howe J, Hughes S, Magill S, McIntyre F, McKee D, Ryan MR, Smith IP, Tyldsely G, Watret R & Turrell WR (2011). The west of Scotland marine ecosystem: a review of scientific knowledge. Marine Scotland Science Report No. 0911, 292pp.
- Bark A, Williams B & Knights B (2007). Current status and temporal trends in stocks of European eel in England and Wales. *ICES Journal of Marine Science* **64**: 1368-1378.
- Bauer G (1992). Variation in the life span and size of the freshwater pearl mussel. *Journal of Animal Ecology* **61**: 425-436.
- Beare DJ, Burns F, Greig A, Jones EG, Peach K, Kienle M, McKenzie E & Reid DG (2004). Long-term increases in prevalence of North Sea fishes having southern biogeographic affinities. *Marine Ecology Progress Series* **284**: 269-278.
- Beaugrand G, Brander KM, Lindley JA, Souissi S & Reid PC (2003). Plankton effect on cod recruitment in the North Sea. *Nature* **426**: 661-664.
- Beggs SE, Cardinale M, Gowen RJ & Bartolino V (2013). Linking cod (*Gadus morhua*) and climate: investigating variability in Irish Sea cod recruitment. *Fisheries Oceanography* **23**: 54-64.
- Béguer-Pon M, Benchetrit J, Castonguay M, Aarestrup K, Campana SE, Stokesbury MJW & Dodson JJ (2012). Shark predation on migrating adult American eels (*Anguilla rostrata*) in the Gulf of St Lawrence. *PLoS ONE* **7**: e46830.
- Bendall V, Ellis JR, Hetherington SJ, McCully S, Righton D & Silva JF (2013). Preliminary observations on the biology and movements of porbeagle *Lamna nasus* around the British Isles. *Collective Volume of Scientific Papers ICCAT* **69**: 1702-1722.
- Blaxter JHS & Hunter JR (1982). The biology of the clupeoid fishes. *Advances in Marine Biology* **20**: 1-223.
- Boulcott P & Wright PJ (2011). Variation in fecundity in the lesser sandeel: implications for regional management. *Journal of the Marine Biological Association of the United Kingdom* **91**: 1273-1280.
- Brander KM, Blom G, Borges MF, Erzini K, Henderson G, MacKenzie BR, Mendes H, Ribiero J, Santos AMP & Tørensen R (2003). Changes in fish distribution in the eastern North Atlantic: are we seeing a coherent response to changing temperature? *ICES Journal of Marine Science* **219**: 261-270.

- Bullough LW, Turrell WR, Buchan P & Priede IG (1998). Commercial deep water trawling at sub-zero temperatures – observations from the Faroe Shetland Channel. *Fisheries Research* **39**: 33-41.
- Buscaino G, Filiciotto F, Buffa G, Bellante A, di Stefano V, Assenza A, Fazio F, Caola G & Mazzola S (2010). Impact of an acoustic stimulus on the motility and blood parameters of European sea bass (*Dicentrarchus labrax* L.) and gilthead sea bream (*Sparus aurata* L.). *Marine Environmental Research* **69**: 136-142.
- Callaway R, Alsvag J, de Boois I, Cotter J, Ford A, Hinz H, Jennings S, Kroncke I, Lancaster J, Piet G, Prince P & Ehrich S (2002). Diversity and community structure of epibenthic invertebrates and fish in the North Sea. *ICES Journal of Marine Science* **59**: 1199-1214.
- CEFAS (2007). Fish and fish assemblages of the British Isles. Report no. C2983. Report to the Department of Trade and Industry, 109pp.
- Chapman CJ (2004). Northern North Sea Shellfish and Fisheries. SEA 5 technical report to the Department of Trade and Industry, 69pp.
- Chapman CJ (2006). Coastal shellfish resources and fisheries in SEA 7. SEA 7 technical report to the Department of Trade and Industry, 76pp.
- Christensen V, Guenette S, Heymans JJ, Walters CJ, Watson R, Zeller D & Pauly D (2003). Hundred year decline of North Atlantic predatory fishes. *Fish and Fisheries* **4**: 1-24.
- Clark PF (1986). North-east Atlantic crabs: an atlas of distribution. The Marine Conservation Society, Ross-on-Wye, 252 pp.
- Corten A & Van de Kamp G (1996). Variation in the abundance of southern fish species in the southern North Sea in relation to hydrography and wind. *ICES Journal of Marine Science* **53**: 1113-1119.
- Cotton PA, Sims DW, Fanshawe S & Chadwick M (2005). The effect of climate variability on zooplankton and basking shark (*Cetorhinus maximus*) relative abundance off southwest Britain. *Fisheries Oceanography* **14**: 151-155.
- Coull KA, Johnstone R & Rogers SI (1998). Fisheries Sensitivity Maps in British Waters. Report to United Kingdom Offshore Operators Association, Aberdeen, 58pp.
- Daan N (2006). Spatial and temporal trends in species richness and abundance for the southerly and northerly components of the North Sea fish community separately, based on IBTS data 1977-2005. *ICES CM 2006 D:02*: 1-10.
- Daan N (2015a). Dragonets (Callionymidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp. 390-395.
- Daan N (2015b). Pipefish (Syngnathidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp. 267-276.
- Daan N, Bromley PJ, Hislop JRG & Nielsen NA (1990). Ecology of North Sea fish. *Netherlands Journal of Sea Research* **26**: 343-386.
- De Silva SS (1973). Abundance, structure, growth and origin of inshore clupeid populations of the west coast of Scotland. *Journal of Experimental Marine Biology and Ecology* **12**: 119-144.
- DECC (2010). Severn tidal power SEA topic paper: migratory and estuarine fish, 280pp.
- Defra (2005). Charting Progress: 3. Marine habitats and species – An integrated assessment of the state of UK seas. A joint report published by the Department for Environment, Food and Rural Affairs, London, 168pp.
- Defra (2010). Charting Progress 2 - Feeder report: Healthy and biologically diverse seas. Draft published by the Department for Environment Food and Rural Affairs on behalf of the UK Marine Monitoring and Assessment Strategy community, London, 744pp
- Defra (2015). Biodiversity 2020: a strategy for England's wildlife and ecosystem services. Report by Defra, 210pp.
- Desforges J-PW, Galbraith M & Ross PS (2015). Ingestion of microplastics by zooplankton in the Northeast Pacific Ocean. *Archives of Environmental Contamination and Toxicology* **69**: 320-330.
- Dickey-Collas M, Heessen H & Ellis J (2015). Shads, herring, pilchard, sprat (Clupeidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp. 139-151.
- Dippner JW (1997). Recruitment success of different fish stocks in the North Sea in relation to climate variability. *Ocean Dynamics* **49**: 277-293.
- Dong Energy (2013). Burbo Bank extension OWF. Environmental Statement Annex 13: Fish and shellfish ecology. Document ref: 5.1.5.13, 135pp.
- Dulvy NK, Rogers SI, Jennings S, Stelzenmüller V, Dye SR & Skjoldal HR (2008). Climate change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. *Journal of Applied Ecology* **45**: 1029-1039.
- Durif CMF, Browman HI, Phillips JB, Skiftesvik AB, Vøllestad LA & Stockhausen HH (2013). Magnetic compass orientation in the European eel. *PLoS ONE* **8**: e59212.

- Edwards E (1979). *The edible crab and its fishery in British waters*. Fishing News Books Ltd, Farnham, 142 pp.
- Edwards M, Helaouet P, Halliday N, Beaugrand G, Fox C, Johns DG, Licandro P, Lynam C, Pitois S, Stevens D, Coombs S & Fonseca L (2011). Fish larvae atlas of the NE Atlantic: results from CPR survey 1948-2005. Sir Alister Hardy Foundation for Ocean Science Report, 22pp.
- Edwards M, Helaouet P, Johns DG, Batten S, Beaugrand G, Chiba S, Hall J, Head E, Hosie G, Kitchener J, Koubbi P, Kreiner A, Melrose C, Pinkerton M, Richardson AJ, Robinson K, Takahashi K, Verheye HM, Ward P & Wootton M (2014). Global Marine Ecological Status Report: results from the global CPR survey 2012/2013. *SAHFOS Technical Report 10*: 1-37.
- Ellis J & Velasco F (2015). Anglerfish (Lophiidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 240-245.
- Ellis J (2015a). Jacks (Carangidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 329-332.
- Ellis J (2015b). Scorpionfish (Scorpaenidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 288.
- Ellis J (2015c). Catsharks (Scyliorhinidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 67-75.
- Ellis J (2015d). Blennies (Blenniidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 387-389.
- Ellis J & Heessen H (2015a). Mackerels and tunnies (Scombridae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 413-419.
- Ellis J & Heessen H (2015b). Lantern sharks (Etmopteridae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 87-88.
- Ellis J & Rogers S (2015). Gobies (Gobiidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 396-411.
- Ellis J & Velasco F (2015). Anglerfish (Lophiidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 240-245.
- Ellis JR, Pawson MG & Shackley SE (1996). The comparative feeding ecology of six species of shark and four species of ray (Elasmobranchii) in the North-East Atlantic. *Journal of the Marine Biological Association of the United Kingdom* **76**: 89-106.
- Ellis JR, Milligan SP, Readdy L, Taylor N & Brown MJ (2012). Spawning and nursery grounds of selected fish species in UK waters. Science Series Technical Report, Cefas, Lowestoft, 147: 56pp.
- Ellis J, Heessen H & McCully Phillips S (2015). Skates (Rajidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 96-124.
- FAO website [accessed February 2016]. <http://www.fao.org/fishery/species/search/en>
- Fariña AC, Azevedo M, Landa J, Duarte R, Sampedro P (2008). *Lophius* in the world; a synthesis on the common features and life strategies. *ICES Journal of Marine Science* **65**: 1272-1280.
- Farrell E, McCully S & Ellis J (2015). Hound sharks (Triakidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.76-80.
- Fässler SMM, Payne MR, Brunel T & Dickey-Collas M (2011). Does larval mortality really influence population productivity? An analysis of North Sea herring time series. *Fisheries Oceanography* **20**: 530-543.
- Fernandes PG & Cook RM (2013). Reversal of fish stock decline in the north-east Atlantic. *Current Biology* **23**: 1432-1437.
- Fincham JI, Rijnsdorp AD & Engelhard GH (2013). Shifts in the timing of spawning in sole linked to warming in sea temperatures. *Journal of Sea Research* **75**: 69-76.
- FishBase website [accessed January 2016]. <http://www.fishbase.org/summary/1365>
- Fleischer D, Schaber M & Pipenburg D (2007). Atlantic snake pipefish (*Entelurus aequoreus*) extends its northwards distribution range to Svalbard (Arctic Ocean). *Polar Biology* **30**: 1359-1362.
- Floeter J, Kempf A, Vinther M, Schrum C & Temming A (2005). Grey gumard (*Eutrigla gurnardus*) in the North Sea: an emerging key predator? *Canadian Journal of Fisheries and Aquatic Science* **62**: 1853-1864.
- Formicki K, Sadowski M, Tanski A, Korzelecka-Orkisz A & Winnicki A (2004). Behaviour of trout (*Salmo trutta* L.) larvae and fry in a constant magnetic field. *Journal of Applied Ichthyology* **20**: 290-294.
- Frederiksen M, Edwards M, Richardson AJ, Halliday NC & Wanless S (2006). From plankton to top predators: bottom-up control of a marine food web across four trophic levels. *Journal of Animal Ecology* **75**: 1259-1268.

- Gauld JA & Hutcheon JR (1990). Spawning and fecundity in the lesser sandeel, *Ammodytes marinus* Raitt in the north-western North Sea. *Journal of Fish Biology* **36**: 611-613.
- Genner MJ, Sims DW, Wearmouth VJ, Southall EJ, Southward AJ, Henderson PA & Hawkins SJ (2004). Regional climate warming drives long-term community changes of British marine fish. *Proceedings of the Royal Society B* **271**: 655-661.
- Genner MJ, Sims DW, Southward AJ, Budd GC, Masterson P, McHugh M, Rendle P, Southall EJ, Wearmouth VJ & Hawkins SJ (2010). Body size dependent responses of a marine fish assemblage to climate change and fishing over a century long scale. *Global Change Biology* **16**: 517-527.
- Gill AB & Bartlett M (2010). Literature review on the potential effects of electromagnetic fields and subsea noise from marine renewable energy developments on Atlantic salmon, sea trout and European eel. Report No. 401, commissioned by Scottish Natural Heritage, 27pp.
- Gibb FM, Gibb IM & Wright PJ (2007). Isolation of Atlantic cod (*Gadus morhua*) nursery areas. *Marine Biology* **151**: 1185-1194.
- Gill AB, Huang Y, Gloyne-Philips I, Metcalfe J, Quayle V, Spencer J & Wearmouth V (2009). COWRIE 2.0 Electromagnetic Fields (EMF) Phase 2: EMF-sensitive fish response to EM emissions from sub-sea electricity cables of the type used by the offshore renewable energy industry. Commissioned by COWRIE Ltd (project reference COWRIE-EMF-1-06), 128pp inc. appendices.
- Gillibrand P, Hay D, Penston M & Murray A (2005). Sea lice and sea trout: are salmon farms causing increased parasitism on wild salmonids in Scotland? *The Challenger Society for Marine Science* **14**: 24-31.
- Goldsmith D, Rijnsdorp A, Vitale F & Heessen H (2015). Right-eyed flounders (Pleuronectidae). In: *HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 452-471.
- González-Irusta JM & Wright PJ (2015). Spawning grounds of Atlantic cod (*Gadus morhua*) in the North Sea. *ICES Journal of Marine Science* **73**: 304-315.
- Gordon JDM (1981). The fish populations of the west of Scotland shelf, part 2. *Oceanography and Marine Biology Annual Review* **19**: 405-441.
- Gordon JDM (2001). Deep-water fisheries at the Atlantic frontier. *Continental Shelf Research* **21**: 987-1003.
- Gordon JDM (2003). The Rockall Trough, Northeast Atlantic: the cradle of deep sea biological oceanography that is now being subjected to unsustainable fishing activity. *Journal of Northwest Atlantic Fishery Science* **31**: 57-83.
- Gordon JDM (2006). Fish and fisheries in the SEA 7 area. Report to the Department of Trade and Industry by Scottish Association for Marine Science (SAMS), Dunstaffnage Marine Laboratory, Oban, 122pp.
- Gordon JDM & Bergstad (1992). Species composition of demersal fish in the Rockall trough, northeastern Atlantic, as determined by different trawls. *Journal of the Marine Biological Association of the United Kingdom* **72**: 213-230.
- Gordon JDM, Harrison EM & Swan SC (1994). A guide to the deep-water fish of the Northeastern Atlantic. Scottish Association for Marine Science (SAMS), Dunstaffnage, 24pp.
- Gore MA, Rowat D, Hall J, Gell FR & Ormond RF (2008). Transatlantic migration and deep mid-ocean diving by basking shark. *Biology Letters* **4**: 395-398.
- Grant SM (2006). An exploratory research survey and biological resource assessment of Atlantic hagfish occurring on the SW slope of Newfoundland Grand Bank. *Journal of North-western Atlantic Fishery Science* **36**: 91-110.
- Greve W, Lange U, Rieners F & Nast J (2001). Predicting the seasonality of North Sea zooplankton. *Senckenbergiana maritima* **31**: 263-268.
- Guerin AJ, Jackson AC, Bowyer PA & Youngson AF (2014). Hydrodynamic models to understand salmon migration in Scotland. Marine Research Report for The Crown Estate, 116pp.
- Halvorsen MB, Casper BM, Woodley CM, Carlson TJ & Popper AN (2012). Threshold for onset of injury in Chinook salmon from exposure to impulsive pile-driving sounds. *PLoS ONE* **7**: e38968.
- Hansen LP, Jonsson N & Jonsson B (1993). Oceanic migration and homing in Atlantic salmon. *Animal Behaviour* **45**: 927-941.
- Harris MP, Beare D, Toresen R, Nøttestad L, Kloppmann M, Dörner H, Peach K, Rushton DRA, Foster-Smith J & Wanless S (2007). A major increase in snake pipefish (*Entelurus aequoreus*) in northern European seas since 2003: potential implications for seabird breeding success. *Marine Biology* **151**: 973-983.
- Haugland M, Holst JC, Holm M & Hansen LP (2006). Feeding of Atlantic salmon (*Salmo salar* L.) post-smolts in the north-east Atlantic. *ICES Journal of Marine Science* **63**: 1488-1500.
- Hawkings AD, Pembroke AE & Popper AN (2015). Information gaps in understanding the effects of noise on fishes and invertebrates. *Reviews in Fish Biology and Fisheries* **25**: 39-64.
- Hay SJ, Hislop JRG & Shanks AM (1990). North Sea Scyphomedusae; summer distribution, estimated biomass and significance, particularly for O-group gadoid fish. *Netherlands Journal of Sea Research* **25**: 113-130.

- Heath M & Gallego A (1997). From the biology of the individual to the dynamics of the population: bridging the gaps in fish early life studies. *Journal of Fish Biology* **51**: 1-29.
- Heath MR & Spiers DC (2012). Changes in species diversity and size composition in the Firth of Clyde demersal fish community, 1927-2009. *Proceedings of the Royal Society B* **279**: 543-552.
- Heath MR, Neat FC, Pinnegar JK, Reid DG, Sims DW & Wright PJ (2012). Review of climate change impacts on marine fish and shellfish around the UK and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems* **22**: 337-367.
- Heath MR, Culling MA, Crozier WW, Fox CJ, Gurney WSC, Hutchinson WF, Nielsen EE, O'Sullivan M, Preedy KF, Righton DA, Spiers DC, Taylor MI, Wright PJ & Carvalho GR (2014). *ICES Journal of Marine Science* doi.: 10.1093/icesjms/fst185.
- Hedger R, MacKenzie E, Heath M, Wright P, Scott B, Gallego A & Bridson J (2004). Analysis of the spatial distributions of mature cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) abundance in the North Sea (1980-1999) using Generalised Additive Models. *Fisheries Research* **70**: 17-25.
- Heessen H & Blasdale T (2015). Redfish (Sebastidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.279-287.
- Heessen H & Daan N (2015). Salmon (Salmonidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.160-166.
- Heessen H (2015a). Argentines (Argentinidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.152-154.
- Heessen H (2015b). Sea-breems and porgies (Sparidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.336-343.
- Heessen HJL, Daan H & Ellis JR (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, 572pp.
- Henderson PA (2007). Discrete and continuous change in the fish community of the Bristol Channel in response to climate change. *Journal of the Marine Biological Association of the UK* **87**: 589-598.
- Henderson PA & Bird DJ (2010). Fish and macro-crustacean communities and their dynamics in the Severn Estuary. *Marine Pollution Bulletin* **61**: 100-114.
- Hiddink JG & ter Hofstede R (2008). Climate induced increases in species richness of marine fishes. *Global Change Biology* **14**: 453-460.
- Hislop JRG & McKenzie K (1976). Population studies of the whiting *Merlangius merlangus* (L.) of the northern North Sea. *Journal du Conseil International pour l'Exploration de la Mer* **37**: 98-111.
- Hislop JRG, Gallego A, Heath MR, Kennedy FM, Reeves SA (2001). A synthesis of early life history of the anglerfish, *Lophius piscatorius* (Linnaeus 1758) in northern British waters. *ICES Journal of Marine Science* **58**: 70-86.
- Hislop J, Bergsat OA, Jakobsen T, Sparholt H, Blasdale T, Wright P, Kloppmann M, Hillgruber N & Heessen H (2015). Codfishes (Gadidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.186-236.
- Holland GJ, Greenstreet SPR, Gibb IM, Fraser HM & Robertson MR (2005). Identifying sandeel *Ammodytes marinus* sediment habitat preferences in the marine environment. *Marine Ecology Progress Series* **303**: 269-282.
- Horwood J (1993). The Bristol Channel sole (*Solea solea* (L.)): a fisheries case study. *Advances in Marine Biology* **29**: 215-367.
- Houghton RG & Harding D (1976). The plaice of the English Channel: spawning and migration. *ICES Journal of Marine Science* **36**: 229-239.
- Hulme M, Jenkins GJ, Lu X, Turnpenny JR, Mitchell TD, Jones RG, Lowe J, Murphy JM, Hassel D, Boorman P, McDonald R & Hill S (2002). Climate change scenarios for the United Kingdom: The UKCIP02 Scientific Report. Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich, 120pp.
- ICES (1991). Report of the multi-species working group. ICES Document CM 1991/G:7.
- ICES (2008). Report of the Study Group for Bycatch of Protected Species (SGBYC), January 29-31 2008, Copenhagen Denmark. ICES CM 2008/ACOM: 48. 80 pp.
- ICES website [accessed January 2016].
<http://www.ices.dk/news-and-events/news-archive/news/Pages/Latest-ICES-advice-on-European-Eel---stocks-remain-critical.aspx>
- Iversen SA, Skogen MD & Svendsen E (2002). Availability of horse mackerel (*Trachurus trachurus*) in the north-eastern North Sea, predicted by the transport of Atlantic water. *Fisheries Oceanography* **11**: 245-250.
- Jansen T & Gjalson H (2013). Population structure of Atlantic mackerel. *PLoS ONE* **8**: e64744.

- Jennings S & Ellis J (2015). Bass (Moronidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp.324-326.
- JNCC website [accessed January 2016].
<http://jncc.defra.gov.uk/protectedsites/sacselection/species.asp?FeatureIntCode=S1095>
- Kaiser MJ, Rogers SI & Ellis JR (1999). Importance of benthic habitat complexity for demersal fish assemblages. *American Fisheries Society Symposium* **22**: 212-223.
- King PA, Fives JM & McGrath D (1994). Reproduction, growth and feeding of the dragonet, *Callionymus lyra* (Teleostei: Callionymidae), in Galway Bay, Ireland. *Journal of the Marine Biological Association of the United Kingdom* **74**: 513-526.
- Kirby MF, Neall P & Tylor T (1999). EROD activity measured in flatfish from the area of the Sea Empress oil spill. *Chemosphere* **38**: 2929-2949.
- Kjesbu OS, Righton D, Kruger-Johnson M, Thorsen A, Michalsen K, Fonn M, Witthames PW (2010). Thermal dynamics of ovarian maturation in Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Science* **67**: 605-625.
- Kloppmann M (2015a). Lampreys (Petromyzontidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp.55-59.
- Kloppmann M (2015b). Hagfish (Myxiniidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp.52-54.
- Kloppmann M & Ellis J (2015). Hatchetfish and pearlsides (Sternoptychidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp. 167-170.
- Laurenson CH, Dobby H, McLay HAS & Leslie B (2008). Biological features of the *Lophius piscatorius* catch in Scottish waters. *ICES Journal of Marine Science* **65**: 1281-1290.
- Lee AJ & Ramster JW (1981). *Atlas of the seas around the British Isles*. Ministry of Agriculture, Fisheries and Food, Lowestoft, 102 pp.
- Lenoir S, Beaugrand G & Lecuyer E (2011). Modelled spatial distribution of marine fish and projected modifications in the North Atlantic Ocean. *Global Change Biology* **17**: 115-129.
- Lindegren M, Diekmann R & Möllmann C (2010). Regime shifts, resilience and recovery of a cod stock. *Marine Ecology Progress Series* **402**: 239-253.
- Lockwood SJ (1988). *The mackerel: its biology, assessment and the management of a fishery*. Fishing News Books, Farnham, Surrey, UK, 181pp.
- Lowe JA, Howard TP, Pardaens A, Tinker J, Holt J, Wakelin S, Milne G, Leake J, Wolf J, Horsburgh K, Reeder T, Jenkins G, Ridley J, Dye S, Bradley S (2009). UK Climate Projections science report: Marine and coastal projections. Met Office Hadley Centre, Exeter, UK, 95pp.
- MacDougall CA (2002). *Thorogobius ephippiatus*. Leopard-spotted goby. Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme. Plymouth: Marine Biological Association of the United Kingdom. Available from: <http://www.marlin.ac.uk/species/Thorogobiusephippiatus.htm>
- Maitland PS & Hatton-Ellis TW (2003). *Ecology of the Allis and Twaite Shad: Conserving Natura 2000 Rivers Ecology Series No. 3*. English Nature, Peterborough, 28pp.
- Makhrov AA & Bolotov IN (2011). Does freshwater pearl mussel (*Margaritifera margaritifera*) change the lifecycle of Atlantic salmon (*Salmo salar*)? *Advances in Gerontology* **1**: 186-194.
- Malcolm IA, Godfrey J & Youngson AF (2010). Review of migratory routes and behaviour of Atlantic salmon, sea trout and European eel in Scotland's coastal environment: implications for the development of marine renewables. Scottish Marine and Freshwater Science, Report No 1:14, 77pp.
- Malcolm IA, Miller CP & Millidine KJ (2015). Spatio-temporal variability in Scottish smolt emigration times and sizes. Scottish Marine and Freshwater Science, Report No 6:2, 19pp.
- Mamie JCJ, Beare DJ, Jones EG, Kienzle M, Dobby H, Heath MR & Reid DG (2007). Aspects of the distribution and growth of bluemouth (*Helicolenus dactylopterus*, Delaroche 1809) since its invasion of the northern North Sea in 1991. *Fisheries Oceanography* **16**: 85-94.
- Marine Scotland website [accessed February 2016].
<http://www.gov.scot/Topics/marine/marine-environment/species/fish/sandeels>
- Marine Scotland website [accessed December 2015].
<http://www.gov.scot/Topics/marine/marine-environment/species/fish/shellfish/nephrops>
- Marine Scotland website [accessed January 2016].
<http://www.gov.scot/Topics/marine/marine-environment/species/fish/shellfish/scallop>
- Marra LJ (1989). Sharkbite on the SL submarine lightwave cable system: history, causes and resolution. *IEEE Journal of Oceanic Engineering* **14**: 230-237.

- Mason J (1983). *Scallop and queen fisheries in the British Isles*. Fishing News Books Ltd, Farnham, 144 pp.
- McCleave JD & Arnold GP (1999). Movements of yellow- and silver-phase European eels (*Anguilla anguilla* L.) tracked in the western North Sea. *ICES Journal of Marine Science* **56**: 510-536.
- McCauley RD, Fewtrell J & Popper AN (2003). High intensity anthropogenic sound damages fish ears. *Journal of the Acoustical Society of America* **113**: 638-642.
- Millner RS, Pillner GM, McCully SR & Høie H (2011). Changes in the timing of otolith zone formation in North Sea cod from otolith records: an early indicator of climate-induced temperature stress? *Marine Biology* **158**: 21-30.
- Mueller-Blenkle C, McGregor PK, Gill AB, Andersson MH, Metcalfe J, Bendall V, Sigray P, Wood DT & Thomsen F (2010). Effects of pile-driving noise on the behaviour of marine fish. Technical report produced for COWRIE Ref: Fish 06-08, 57pp inc. appendices.
- Naismith IA & Knights B (1988). Migration of elvers and juvenile European eels, *Anguilla anguilla* L. in the River Thames. *Journal of Fish Biology* **33**: 161-175.
- Navarro MR, Vilamor B, Myklevoll S, Gil J, Abaunza P & Canoura J (2012). Maximum size of Atlantic mackerel (*Scomber scombrus*) and Atlantic chub mackerel (*Scomber colias*) in the north-east Atlantic. *Cybius* **36**: 406-408.
- Neat F & Campbell N (2011). Demersal fish diversity of the isolated Rockall Plateau compared with the adjacent west coast of Scotland. *Biological Journal of the Linnean Society* **104**: 138-147.
- Nelson JS (2006). *Fishes of the World*. John Wiley & Sons, New Jersey, USA, 601pp.
- Newton AW (1984). Scottish tagging experiments in the North Sea and in division VIa. *ICES CM 1984 G:67*: 4.
- Öhman MC, Sigray P & Westerberg H (2007). Offshore windmills and the effects of electromagnetic fields on fish. *Ambio* **36**: 630-633.
- OSPAR (2010a). Quality Status Report 2010. OSPAR Commission, London, 176pp.
- Pade NG, Quieroz N, Humphries NE, Witt MJ, Jones CS, Noble LR & Sims DW (2009). First results from satellite linked archival tagging of porbeagle shark, *Lamna nasus*: Area, fidelity, wider-scale movements and plasticity in diel depth changes. *Journal of Experimental Marine Biology and Ecology* **370**: 64-74.
- Parker-Humphreys (2004). Distribution and relative abundance of demersal fishes from beam trawl surveys in the Bristol Channel (ICES Division VIIF), 1993-2001. Report by Cefas: Science Series Technical Report **123**: 69pp.
- Pawson MG & Robson CF (1998). Chapter 5.7. Fish: exploited sea-fish. In: JH Barne, CF Robson, SS Kaznowska, JP Doody, NC Davidson & AL Buck Eds. Coasts and seas of the United Kingdom. Region 8 Sussex: Rye Bay to Chichester Harbour (Coastal Directory Series). Joint Nature Conservation Committee, Peterborough pp. 95-98.
- Pawson MG (1995). Biogeographical identification of English Channel fish and shellfish stocks. Report No. 99. Fisheries Research Technical Report by the Centre for Environment, Fisheries and Aquaculture Science, Lowestoft, 72pp.
- Pawson M, Pickett GD & Witthames PR (2000). The influence of temperature on the onset of first maturity in sea bass. *Journal of Fish Biology* **56**: 319-327.
- Pawson MG, Kupschus S & Pickett GD (2007). The status of sea bass (*Dicentrarchus labrax*) stocks around England and Wales, derived using a separable catch-at-age model, and implications for fisheries management. *ICES Journal of Marine Science* **64**: 346-356.
- Peres MB & Vooren CM (1991). Sexual development, reproductive cycle, and fecundity of the school shark *Galeorhinus galeus* off southern Brazil. *Fishery Bulletin* **89**: 655-667.
- Perry AL, Low PJ, Ellis JR & Reynolds JD (2005). Climate change and distribution shifts in marine fishes. *Science* **308**: 1912-1915.
- Petitgas P, Rijnsdorp AD, Dickey-Collas M, Engelhard GH, Peck MA, Pinnegar JK, Drinkwater K, Huret M & Nash RDM (2013). Impacts of climate change on the complex life-cycles of fish. *Fisheries Oceanography* **22**: 121-139.
- Peyronnet A, Friedland KD & Ó Maoileidigh N (2008). Different ocean and climate factors control the marine survival of wild and hatchery Atlantic salmon in the north-east Atlantic Ocean. *Journal of Fish Biology* **73**: 945-962.
- Pilling GM, Millner RS, Easy MW & Tidd AN (2007). Phenology and North Sea cod *Gadus morhua* L.: has climate change affected otolith annulus formation and growth? *Journal of Fish Biology* **70**: 584-599.
- Pinnegar JK, Sims DW & Heath M (2008). Marine Climate Change Impacts Partnership (MCCIP) Annual Report Card 2007-2008 Scientific Review – Fish and Fisheries.
- Pinnegar J, Blasdale T, Campbell N, Coates S, Colclough S, Fraser H, Greathead C, Greenstreet S, Neat F, Sharp R, Simms D, Stevens H & Waugh A (2010). Charting Progress 2: Healthy and biologically diverse seas. Feeder Report, Section 3.4: Fish. Published by Defra, 128pp.
- Polockzanska ES & Ansell AD (1999). Imposex in the whelks *Buccinum undatum* and *Neptunea antiqua* from the west coast of Scotland. *Marine Environmental Research* **47**: 203-212.

- Popper AN, Smith ME, Cott PA, Hanna BW, MacGillivray AO, Austin ME & Mann DA (2005). Effects of exposure to seismic airgun use on hearing of three fish species. *Journal of the Acoustic Society of America* **117**: 3958-3971.
- Pörtner HO & Peck MA (2010). Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. *Journal of Fish Biology* **77**: 1745-1779.
- Potts GW & Swaby SE (1997). Chapter 5.9. Fish: other species. In: JH Barne, CF Robson, SS Kaznowska, JP Doody & NC Davidson Eds. *Coasts and seas of the United Kingdom. Regions 15 & 16 North-west Scotland: The Western Isles and West Highlands (Coastal Directory Series)*. Joint Nature Conservation Committee, Peterborough, UK, pp. 138-140.
- Power M & Attrill MJ (2002). Factors affecting long-term trends in the estuarine abundance of pogge (*Agonus cataphractus*). *Estuarine, Coastal and Shelf Science* **54**: 941-949.
- Reiss H, Degraer S, Duineveld GCA, Kröncke I, Aldridge J, Craeymeersch JA, Eggleton JD, Hillewaert H, Lavaleye MSS, Moll A, Pohmann T, Rachor E, Robertson M, Vanden Berghe E, van Hoey G & Rees HL (2010). Spatial patterns of infauna, epifauna and demersal fish communities in the North Sea. *ICES Journal of Marine Science* **67**: 278-293.
- Righton D, Quayle VA, Hetherington S & Burt G (2007). Movements and distribution of cod (*Gadus morhua*) in the southern North Sea and English Channel: results from conventional and electronic tagging experiments. *Journal of the Marine Biological Association of the United Kingdom* **87**: 599-613.
- Rijnsdorp A, Goldsmith D, Heessen H & van Hal R (2015). Soles (Soleidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp. 472-482.
- Robson CF (1996). Chapter 5.8. Fish: salmon, sea trout and eels. In: JH Barne, CF Robson, SS Kaznowska, JP Doody, NC Davidson & AL Buck Eds. *Coasts and seas of the United Kingdom. Region 3 North-east Scotland: Cape Wrath to St. Cyrus (Coastal Directory Series)*. Joint Nature Conservation Committee, Peterborough, UK, pp. 107-108.
- Robson CF (1997). Chapter 5.7. Fish: exploited sea-fish. In: JH Barne, CF Robson, SS Kaznowska, JP Doody & NC Davidson Eds. *Coasts and seas of the United Kingdom. Regions 15 & 16 North-west Scotland: The Western Isles and West Highlands (Coastal Directory Series)*. Joint Nature Conservation Committee, Peterborough, UK, pp. 131-135.
- Rodríguez-Muñoz R, Nicieza AG & Braña F (2001). Effects of temperature on developmental performance, survival and growth of sea lamprey embryos. *Journal of Fish Biology* **58**: 475-486.
- Rogers S & Stocks R (2001). North Sea Fish and Fisheries. Technical report produced by the Centre for Environment, Fisheries and Aquaculture Science for Strategic Environment Assessment - SEA 2, 72pp.
- Rogers SI, Rijnsdorp AD, Damm U & Vanhee W (1998). Demersal fish populations in the coastal waters of the UK and continental NW Europe from beam trawl survey data collected from 1990-1995. *Journal of Sea Research* **39**: 79-102.
- Rowe S, Hutchings JA (2004). The function of sound production by Atlantic cod as inferred from patterns of variation in drumming muscle mass. *Canadian Journal of Zoology* **82**: 1391-1398.
- Russell FS (1976). *The eggs and planktonic stages of British marine fishes*. Academic Press, London, 524 pp.
- Rutterford LA, Simpson SD, Jennings S, Johnson MP, Blanchard JL, Schön P-J, Sims DW, Tinker J & Genner MJ (2015). Future fish distributions constrained by depth in warming seas. *Nature Climate Change* doi: 10.1038/nclimate2607.
- Sell A & Heessen H (2015). Gurnards (Triglidae). In: HJL Heessen, H Daan & JR Ellis (2015). *Fish Atlas of the Celtic Sea, North Sea and Baltic Sea*. Wageningen Academic Publishers, the Netherlands, pp.289-301.
- Shelton RGJ (1978). On the feeding of the hagfish *Myxine glutinosa* in the North Sea. *Journal of the Marine Biological Association of the United Kingdom* **58**: 81-86.
- Simpson SD, Jennings S, Johnson MP, Blanchard JL, Schön P-J, Sims DW & Genner MJ (2011). Continental shelf-wide response of a fish assemblage to rapid warming of the sea. *Current Biology* **21**: 1565-1570.
- Simpson S, Blanchard J & Genner M (2013). Impacts of climate change on fish. *MCCIP Science Review 2013*, pp. 113-124.
- Simpson SD, Purser J & Radford AN (2015). Anthropogenic noise compromises anti-predator behaviour in European eels. *Global Change Biology* **21**: 586-593.
- Sims DW, Southall EJ, Richardson AJ, Reid PC & Metcalfe JD (2003). Seasonal movements and behaviour of basking sharks from archival tagging; no evidence of winter hibernation. *Marine Ecology Progress Series* **248**: 187-196.
- Sims DW, Wearmouth VJ, Genner MJ, Southward AJ & Hawkins J (2004). Low-temperature driven early migration of a temperate marine fish. *Journal of Animal Ecology* **73**: 333-341.
- Sims DW, Southall EJ, Metcalfe JD & Pawson MG (2005a). Basking shark population assessment. Final report for Global wildlife Division of Defra Tender CR0247.

- Sims DW, Southhall EJ, Tarling GA & Metcalfe J (2005b). Habitat-specific normal and reverse diel vertical migration in the plankton-feeding basking shark. *Journal of Animal Ecology* **74**: 755-761.
- Slotte A & Fiksen Ø (2000). State-dependent spawning migration in Norwegian spring-spawning herring (*Clupea harengus* L.). *Journal of Fish Biology* **56**: 138-162.
- Soldandt J-L & Chassin E (2014). Basking shark watch – overview of data from 2009-2013. Report by the Marine Conservation Society, 6pp.
- Sparholt H (2015). Sandeels (Ammodytidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp.377-381.
- Staaterman E, Paris BB & Kough AS (2014). First evidence of fish larvae producing sounds. *Biology Letters* **10**: 20140643.
- Stanley JA, Radford CA & Jeffs AG (2012). Effects of underwater noise on larval settlement. In: Popper AN & Hawkins A (2012). The effects of noise on aquatic life. *Advances in Experimental Medicine and Biology* **730**: 371-374.
- Stebbing ARD, Turk SMT, Wheeler A & Clarke KR (2002). Immigration of southern fish species to south-west England linked to warming of the North Atlantic (1960-2001). *Journal of the Marine Biological Association of the United Kingdom* **82**: 177-180.
- Stelzenmüller V, Ellis JR & Rogers SI (2010). Towards a spatially explicit risk assessment for marine management: assessing the vulnerability of fish to aggregate extraction. *Biological Conservation* **143**: 230-238.
- Swaby SE & Potts GW (1998). Chapter 5.9. Fish: other species. In: JH Barne, CF Robson, SS Kaznowska, JP Doody, NC Davidson & AL Buck Eds. Coasts and seas of the United Kingdom. Region 8 Sussex: Rye Bay to Chichester Harbour (Coastal Directory Series). Joint Nature Conservation Committee, Peterborough pp. 101-102.
- Ter Hofstede R, Hiddinck JG & Rijnsdorp AD (2010). Regional warming changes fish species richness in the eastern North Atlantic Ocean. *Marine Ecology progress Series* **414**: 1-9.
- The Seahorse Trust (2012). British Seahorse Survey 2011, 50pp.
- Thurstan RH, Brockington S & Roberts CM (2010). The effects of 118 years of industrial fishing on UK bottom trawl fisheries. *Nature Communications* **1**: 1-6.
- Tobin D, Wright PJ, Gibb FM & Gibb IM (2010). The importance of life stage to population connectivity in whiting (*Merlangius merlangus*) from the northern European shelf. *Marine Biology* **157**: 1063-1073.
- Van Beek FA, Rijnsdorp AD & de Clerck R (1989). Monitoring juvenile stocks of flatfish in the Wadden Sea and the coastal areas of the southeastern North Sea. *Helgolander Marine Research* **43**: 461-477.
- Van Damme CJG & Thorsen A (2014). Mackerel historic fecundity and atresia data from the mackerel egg survey. Working document for the mackerel benchmark at the WKPELA, 6pp.
- Van Deurs M, van Hal R, Tomczak MT, Jónasdóttir SH & Dolmer P (2009). Recruitment of lesser sandeel *Ammodytes marinus* in relation to density dependence and zooplankton composition. *Marine Ecology Progress Series* **381**: 249-258.
- Vaz F, Carpentier A & Coppin F (2007). Eastern English Channel fish assemblages: measuring the structuring effect of habitats on distinct sub-communities. *ICES Journal of Marine Science* **64**: 271-287.
- Velasco F, Heessen H, Rijnsdorp A & de Boois I (2015). Turbots (Scophthalmidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp. 429-446.
- Walker A & Ellis J (2015). Freshwater eels (Anguillidae). In: HJL Heessen, H Daan & JR Ellis (2015). Fish Atlas of the Celtic Sea, North Sea and Baltic Sea. Wageningen Academic Publishers, the Netherlands, pp.127-129.
- Walker P, Howlett G & Millner R (1997). Distribution, movement and stock structure of three ray species in the North Sea and eastern English Channel. *ICES Journal of Marine Science* **54**: 797-808.
- Walsh M, Reid DG & Turrell WR (1995). Understanding mackerel migration off Scotland: tracking with echosounders and commercial data and including environmental correlates and behaviour. *ICES Journal of Marine Science* **52**: 925-939.
- Wames S & Jones BW (1995). Species distributions from English Celtic Sea groundfish surveys, 1984 to 1991. Report No. 98. Fisheries Research Technical Report by the Centre for Environment, Fisheries and Aquaculture Science, Lowestoft, 42pp.
- Wheeler A (1978). *Key to the Fishes of Northern Europe*. Warne, London, 380pp.
- Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J & Tortonese E (1986). *Fishes of the North-eastern Atlantic and the Mediterranean*. UNESCO, Paris.
- Winslade P (1974). Behavioural studies on the lesser sandeel *Ammodytes marinus* (Raitt) I. The effect of food availability on activity and the role of olfaction in food detection. *Journal of Fish Biology* **6**: 565-576.

Witt MJ, Hardy T, Johnson L, McClellan CM, Pikesley SK, Ranger S, Richardson PB, Solandt J-L, Speedie C, Williams R & Godley BJ (2012). Basking sharks in the north-east Atlantic: spatio-temporal trends from sightings in UK waters. *Marine Ecology Progress Series* **459**: 121-134.

Witt MJ, Doherty PD, Godley BJ, Graham RT, Hawkes LA & Henderson SM (2016). Basking shark satellite tagging project: insights into basking shark (*Cetorhinus maximus*) movement, distribution and behaviour using satellite telemetry. Scottish Natural Heritage Commissioned Report, No. 908, 80pp.