



Local Tobacco Control Profiles

Indicators at a glance February 2016

Notes:

- Value cells are shaded **red**, **amber** or **green** where the England value can be benchmarked against a goal.
- In the change columns, prev refers to the change in England value compared to the previous data point; base refers to the change in England value compared to the baseline value.
- Changes are only shown if they are statistically significant. An upwards arrow (↑, ↑ or ↑) represents a significant increase in the indicator value, a downwards arrow (↓, ↓ or ↓) represents a significant decrease. A sideways arrow (↔) is displayed if there has been no significant change.
- The arrows are coloured **green** and **red** for those indicators where a change can be described as improving or worsening respectively.

Domain: smoking prevalence in adults				
Indicator	Period	Value	Unit	Change from previous
Smoking Prevalence in adults - current smokers (IHS)	2014	18.0	%	↓
Smoking Prevalence in adults - ex-smokers (IHS)	2014	33.9	%	
Smoking Prevalence in adults - never smokers (IHS)	2014	48.1	%	
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	28.0	%	↓
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014	30.8	%	
Smoking prevalence in adults in routine and manual occupations - never smokers (IHS)	2014	41.2	%	
Smoking prevalence in adults - current smokers (QOF)	2014/15	18.6	%	↓
Smoking status at time of delivery	2014/15	11.4	%	↓

Domain: smoking prevalence in young people				
Indicator	Period	Value	Unit	Change from previous
Smoking prevalence age 15 years - regular smokers (SDD survey)	2014	8	%	↔
Smoking prevalence age 15 years - occasional smokers (SDD survey)	2014	5	%	↓
Smoking prevalence at age 15 - current smokers (WAY survey)	2014/15	8.2	%	
Smoking prevalence at age 15 - regular smokers (WAY survey)	2014/15	5.5	%	
Smoking prevalence at age 15 - occasional smokers (WAY survey)	2014/15	2.7	%	
Use of e-cigarettes at age 15 years (WAY survey)	2014/15	18.4	%	
Use of other tobacco products at age 15 years (WAY survey)	2014/15	15.2	%	
Smoking prevalence at ages 11-15 - regular smokers (modelled estimates)	2009 - 12	3.1	%	
Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates)	2009 - 12	1.4	%	
Smoking prevalence at age 15 - regular smokers (modelled estimates)	2009 - 12	8.7	%	
Smoking prevalence at age 15 - occasional smokers (modelled estimates)	2009 - 12	3.9	%	
Smoking prevalence at ages 16-17 - regular smokers (modelled estimates)	2009 - 12	14.7	%	
Smoking prevalence at ages 16-17 - occasional smokers (modelled estimates)	2009 - 12	5.8	%	

Domain: smoking related mortality

Indicator	Period	Value	Unit	Change from previous
Smoking attributable mortality	2012 - 14	274.8	DSR per 100,000	↓
Deaths from lung cancer	2012 - 14	59.5	DSR per 100,000	↔
Deaths from chronic obstructive pulmonary disease	2012 - 14	51.7	DSR per 100,000	↔
Smoking attributable deaths from heart disease	2012 - 14	29.7	DSR per 100,000	↓
Smoking attributable deaths from stroke	2012 - 14	9.3	DSR per 100,000	↔

Domain: smoking related ill health

Indicator	Period	Value	Unit	Change from previous
Premature births (<37 weeks gestation)	2010 - 12	75.8	Crude rate per 1,000	
Low birth weight of term babies	2014	2.9	%	↔
Hospital admissions for asthma (under 19 years)	2013/14	197.1	DSR per 100,000	
Smoking attributable hospital admissions	2014/15	1671.2	DSR per 100,000	↑
Cost per capita of smoking attributable hospital admissions	2011/12	38.0	£ per capita	
Lung cancer registrations	2010 - 12	76.0	DSR per 100,000	↔
Oral cancer registrations	2010 - 12	13.2	DSR per 100,000	↑

Domain: impact of smoking

Indicator	Period	Value	Unit	Change from previous
Indicative tobacco sales figures (£ millions)	2013	15446.1	£	↑
Illicit tobacco: market share (£ millions)	2014/15	1200	£	↔
Illicit tobacco: share of cigarette sales (%)	2014/15	10	%	↔
Fatalities from accidental fires ignited by smoking materials	2013/14	72	Count	↓
Accidental fires ignited by smoking related materials	2013/14	3300	Count	↑
Attitudes to smoking in 15 year olds - 'smoking causes harm to others' (WAY survey)	2014/15	91	%	

Domain: smoking quitters

No. setting a quit date	2014/15	5549.0	Crude rate per 100,000	↓
Successful quitters at 4 weeks	2014/15	2829.0	Crude rate per 100,000	↓
Successful quitters (CO validated) at 4 weeks	2014/15	1954.0	Crude rate per 100,000	↓
Completeness of NS-SEC recording by Stop Smoking Services	2014/15	88.3	%	↓
Cost per quitter	2014/15	420.0	£ per capita	