



Local Tobacco Control Profiles

Indicators at a glance (May 2015)

| Tobacco summary of indicators - May 2015 | | | | |
|---|-----------|---------|------------------------|----------------------|
| Indicator | Period | Value | Unit | Change from previous |
| Smoking Prevalence (IHS) | 2013 | 18.4 | % | ↓ |
| Smoking prevalence - routine & manual | 2013 | 28.6 | % | ↓ |
| Successful quitters at 4 weeks | 2013/14 | 3524 | Crude rate per 100,000 | |
| Successful quitters (CO validated) at 4 weeks | 2013/14 | 2472 | Crude rate per 100,000 | |
| Completeness of NS-SEC recording by Stop Smoking Services | 2013/14 | 86.2 | % | |
| Smoking status at time of delivery | 2013/14 | 12.0* | % | ↓ |
| Low birth weight of term babies | 2012 | 2.8 | % | ↔ |
| Smoking prevalence age 15 years - regular smokers | 2013 | 8 | % | ↔ |
| Smoking prevalence age 15 years - occasional smokers | 2013 | 10 | % | ↔ |
| Lung cancer registrations | 2010 - 12 | 76.0 | DSR per 100,000 | ↔ |
| Oral cancer registrations | 2010 - 12 | 13.2 | DSR per 100,000 | ↑ |
| Deaths from lung cancer | 2011 - 13 | 60.2 | DSR per 100,000 | ↔ |
| Deaths from chronic obstructive pulmonary disease | 2011 - 13 | 51.5 | DSR per 100,000 | ↑ |
| Smoking attributable mortality | 2011 - 13 | 288.7 | DSR per 100,000 | ↓ |
| Smoking attributable deaths from heart disease | 2011 - 13 | 32.7 | DSR per 100,000 | ↓ |
| Smoking attributable deaths from stroke | 2011 - 13 | 11 | DSR per 100,000 | ↓ |
| Smoking attributable hospital admissions | 2012/13 | 1688 | DSR per 100,000 | ↔ |
| Cost per capita of smoking attributable hospital admissions | 2011/12 | 38.0 | £ per capita | |
| Accidental fires ignited by smoking related materials | 2012/13 | 3143 | Count | ↓ |
| Fatalities from accidental fires ignited by smoking materials | 2012/13 | 73 | Count | ↓ |
| Smoking prevalence modelled estimates – occasional smokers aged 11-15 years | 2009-12 | 1.4 | % | |
| Smoking prevalence modelled estimates – occasional smokers aged 15 years | 2009-12 | 3.9 | % | |
| Smoking prevalence modelled estimates – occasional smokers aged 16-17 years | 2009-12 | 5.8 | % | |
| Smoking prevalence modelled estimates – regular smokers aged 11-15 years | 2009-12 | 3.1 | % | |
| Smoking prevalence modelled estimates – regular smokers aged 15 years | 2009-12 | 8.7 | % | |
| Smoking prevalence modelled estimates – regular smokers aged 16-17 years | 2009-12 | 14.7 | % | |
| Indicative tobacco sales figures (£ millions) | 2013 | 15446.1 | Count | ↑ |

Notes:

- Changes are only shown if they are statistically significant. An upwards arrow represents a significant increase in the indicator value, a downwards arrow represents a significant decrease. A sideways arrow is displayed if there has been no significant change.
- The arrows are coloured green and red for those indicators where a change can be described as improving or worsening respectively