

Ad Hoc Statistical Bulletin

Deliberate Self Harm (DSH) in the UK Armed Forces

1 April 2010 – 31 March 2015 (revised)

Date 22 September 2016

Overview

This is an ad hoc statistical bulletin providing information on the number and rate of UK Armed Forces personnel who had at least one episode of Deliberate Self Harm (DSH) recorded between 2010/11 and 2014/15 on MOD held systems.

In July 2014 High Court judges ordered a new inquest into the death of Private Cheryl James at Deepcut Barracks in 1995. This bulletin has been developed following the provision of data to the Coroner's Inquest and to ensure the public has equal access to the information and supports the MOD's commitment to release information where possible.

The current measure of DSH in the UK Armed Forces is based on the initial notification of casualty system (NOTICAS), an administrative system used to inform chain of command of casualties and; primary care data, as captured on the Defence Medical Information Capability Programme (DMICP). The measure counts the total number of UK Armed Forces personnel who had at least one DSH event recorded after 2009 either on NOTICAS and/or DMICP.

Issued by: Head of Defence Statistics (Health) | MoD | Abbey Wood | Oak 0 West | Bristol | BS34 8JH

Telephone: 030679 84423 Email: DefStrat-Stat-Health-Hd@mod.uk

Press office: 020721 83253

Link to stats: https://www.gov.uk/government/statistics/mod-national-and-official-statistics-by-topic

Results

Table 1 below presents the number and rate of UK Armed Forces personnel who had at least one DSH event recorded over the whole time period by demographics and year.

Table 1: UK Armed Forces personnel DSH¹, by demographics and year, numbers, rates per 1,000 personnel at risk per annum and 95% Confidence Interval (CI) 2010/11 - 2014/15

	2010/11			2011/12			2012/13			2013/14			2014/15			Percentage change over	
	n	rate	95% CI	n	rate	95% CI	r	rate	95% CI	n	rate	95% CI	n	rate	95% CI	whole pe	riod
All ²	339	1.7	(1.5 - 1.9)	287	1.5	(1.3 - 1.6)	301	1.6	(1.4 - 1.8)	365	2.1	(1.9 - 2.3)	383	2.3	(2.1 - 2.5)	36%	1
Service																	
Naval Service	63	1.6	(1.2 - 2.0)	43	1.2	(0.8 - 1.5)	34	1.0	(0.6 - 1.3)	61	1.8	(1.3 - 2.2)	55	1.6	(1.2 - 2.1)	2%	
Army	249	2.1	(1.9 - 2.4)	213	1.8	(1.6 - 2.1)	236	2.1	(1.8 - 2.4)	268	2.5	(2.2 - 2.9)	290	3.0	(2.6 - 3.3)	40%	1
RAF	27	0.6	(0.4 - 0.9)	31	0.7	(0.5 - 1.0)	31	8.0	(0.5 - 1.1)	36	1.0	(0.7 - 1.3)	38	1.1	(0.7 - 1.4)	75%	
Gender																	
Male	275	1.5	(1.3 - 1.7)	231	1.3	(1.1 - 1.5)	251	1.5	(1.3 - 1.7)	306	1.9	(1.7 - 2.1)	319	2.1	(1.9 - 2.4)	40%	4
Female	64	3.4	(2.6 - 4.3)	54	3.0	(2.2 - 3.8)	50	2.8	(2.0 - 3.6)	59	3.5	(2.6 - 4.4)	64	4.0	(3.0 - 4.9)	16%	
Rank																	
Officer	8	0.2	(0.1 - 0.5)	7	0.2	(0.1 - 0.4)	5	0.1	(0.0 - 0.3)	13	0.4	(0.2 - 0.7)	14	0.5	(0.3 - 0.8)	102%	
Rank	331	2.0	(1.8 - 2.2)	277	1.7	(1.5 - 1.9)	296	1.9	(1.7 - 2.2)	346	2.4	(2.1 - 2.6)	369	2.7	(2.4 - 3.0)	36%	1
Training Status																	
Trained	294	1.8	(1.6 - 2.0)	265	1.6	(1.4 - 1.8)	272	1.7	(1.5 - 2.0)	327	2.2	(2.0 - 2.5)	355	2.6	(2.3 - 2.9)	45%	1
Untrained	45	2.8	(2.0 - 3.6)	20	1.5	(0.9 - 2.3)	29	2.1	(1.4 - 3.0)	38	2.9	(2.0 - 3.9)	28	2.4	(1.6 - 3.4)	-16%	
Age																	
< 20	65	4.7	(3.6 - 5.9)	28	2.7	(1.8 - 3.9)	42	4.5	(3.2 - 5.9)	44	5.2	(3.7 - 6.7)	43	5.5	(3.9 - 7.2)	17%	
20-24	133	2.9	(2.4 - 3.4)	121	2.8	(2.3 - 3.3)	130	3.2	(2.6 - 3.7)	148	4.0	(3.3 - 4.6)	146	4.3	(3.6 - 5.0)	47%	1
25-29	81	1.9	(1.5 - 2.3)	68	1.6	(1.2 - 1.9)	59	1.4	(1.0 - 1.7)	88	2.2	(1.7 - 2.6)	76	1.9	(1.5 - 2.4)	4%	
30-34	27	0.9	(0.6 - 1.3)	28	0.9	(0.6 - 1.2)	31	0.9	(0.6 - 1.3)	49	1.5	(1.1 - 2.0)	49	1.6	(1.2 - 2.1)	85%	
35-39	19	0.6	(0.4 - 1.0)	20	8.0	(0.5 - 1.2)	21	0.9	(0.5 - 1.3)	26	1.2	(0.8 - 1.7)	35	1.6	(1.1 - 2.1)	148%	1
40-44	6	0.3	(0.1 - 0.6)	~	0.7	(0.4 - 1.2)	11	0.6	(0.3 - 1.0)	~	0.3	(0.1 - 0.7)	25	1.6	(1.0 - 2.3)	434%	1
45+	8	0.5	(0.2 - 0.9)	~	0.2	(0.1 - 0.6)	6	0.3	(0.1 - 0.8)	~	0.2	(0.1 - 0.6)	9	0.5	(0.2 - 1.0)	12%	

Source: Initial NOTICAS and DMICP

Reported rates of DSH among UK Armed Forces personnel were low at **0.2%** in 2014/15. However, there was an increasing trend in the reported rates of DSH among UK Armed Forces personnel in recent years with a significant increase of 36% between 2010/11 and 2014/15. It is difficult to quantify how much of this rise was accounted for by improved methods of capturing DSH data and what may be a true rise in DSH among UK Armed Forces personnel.

The UK Armed Forces population at highest risk of DSH between 2010/11 and 2014/15 were:

- Army personnel
- Females
- Other ranks
- Personnel aged under 24

The risk groups for DSH in the Armed Forces were similar to those seen in the general population, where females and younger age groups were found to be at greatest risk of presentation at a hospital with a self-harm episode^a. This finding was also in line with literature available on the general population where females were more likely to present with mental health

^{1.} Only those with at least one DSH event since 2009, captured.

^{2.} Sub-group totals do not always sum to the overall total as a number of records had missing demographic data at the time of the incident (see data source JPA notes)

^{3.} Naval Service includes Royal Navy and Royal Marines

^{4. ~} Numbers fewer than five have been suppressed to protect patient confidentiality (see methodology notes)

a Skegg, K. (2005) Self Harm, Lancet, 366, 1471-83.

problems compared to males. It is suggested this was because females were likely to have more interactions with health professionals than males^b.

Comparisons between UK Armed Forces DSH rates and the UK population are difficult. However, from the evidence available the military population appears healthier with a lower lifetime prevalence of attempted suicide and self-harm, within the range of general population estimates^c.

Limitations

There are known difficulties in accurately capturing DSH episodes common to the UK general population and military populations. The reporting of DSH is dependent on when the DSH is identified and brought to the attention of the appropriate parties either by the individual themselves seeking help or if discovered by a third party. Potential barriers to seeking care include:

- The associated stigma relating to DSH
- The mechanism they use to self-harm some of which may not be visible
- It may be possible for the individual to treat themselves at home (for example cuts)^d.

Information on numbers and rates of DSH in the whole UK population is not available, thus any comparisons between the UK military and civilians has been based on small location based studies.

Data limitations:

From the available data it is not possible to differentiate between new episodes and the on-going treatment of a DSH episode, thus this bulletin counts personnel who had a DSH event recorded from 2009 onwards only once over the whole time period presented.

DMICP data was sourced using read codes from the data warehouse where the read code indicated Deliberate Self Harm. Information entered using free text has not been included as the information is not held in the central data warehouse; therefore the figures provided are a minimum.

The NOTICAS data relied on either a DSH tick box on the form being completed or via a free text search of the comments section, thus the figures provided are a minimum.

Background notes

This ad hoc statistical bulletin has been released in response to a request from the Coroner of the Deepcut inquiry, requesting information on DSH in the UK Armed Forces, in particular in Army personnel in comparison with the UK general population.

This statistical bulletin ensures MOD is open and transparent about the methodology and quality of any statistics and that equal access is given to all, as required by the Code of Practice for Official Statistics.

b Office for National Statistics (2003) Better or Worse: A follow up study of the mental health of adults in Great Britain. London: National Statistics.

c Pinder et al., (2011) Self-harm and attempted suicide among UK Armed Forces personnel: results of a cross sectional survey.

d McAllister, M. (2003). Multiple meanings of self-harm: A critical review. International Journal of Mental Health Nursing, 12, 177-185.

Care management

Assessment and care-management within the Armed Forces for personnel experiencing mental health problems is available at three levels:

- Primary Health Care (PHC), by the patient's own Medical Officer (MO).
- Through specialists in military Departments of Community Mental Health (DCMH).
- In hospitals, either the NHS or the contracted In-Patient Service Provider (ISP).

The level of care a patient may require is determined by a number of factors, including the severity of symptoms and the degree of risk posed by the patient's current condition.

Data used in this response covers all aspects of the care management pathway where entered onto NOTICAS and/or DMICP.

Data sources

Initial Notification of Casualty (NOTICAS)

Notification of Casualty (or "NOTICAS") is the name for the formalised system of reporting casualties within the UK Armed Forces. The NOTICAS reports raised for casualties contain information on how seriously medical staff judge their condition to be. They are not strictly medical categories but are designed to give an indication of the severity of the injury or illness to inform what the individual's next of kin are told.

Initial NOTICAS casualty reporting system data was used in this bulletin as it covers incidents where personnel have been admitted to an NHS Emergency Department and where the next of kin has been informed.

Information supplied by the Joint Personnel Administration Business Information Cell (JPA-BIC) highlighted incidents where the DSH box was ticked. In addition to these, NOTICAS incidents where a free text search of the comments section highlighted it was a DSH related incident was also included. Free text search included the words 'Deliberate', 'Self Harm', Deliberate Self Harm' and 'DSH'.

Defence Medical Information Capability Program (DMICP)

DMICP is the MOD electronic integrated primary health care record for UK Armed Forces personnel. DMICP was rolled out in 2007 and legacy medical data for currently serving personnel was migrated across during rollout.

UK Armed Forces personnel have access to specialist mental health services via referrals made by their GP, provided through MOD Department of Community Mental Health (DCMH) or the MOD in-patient provider. DCMH are specialised psychiatric services based on community mental health teams closely located with primary care services at MOD sites in the UK and abroad. DCMH staff record the initial mental health assessment during a patient's first appointment, based on presenting complaints. The information is based on a clinician's diagnosis at initial assessment and final diagnoses may differ as some patients do not present the full range of symptoms, signs or clinical history during their first appointment. Data presents those assessed with a mental disorder at initial appointment.

All UK based and aero-medically evacuated Service personnel based overseas requiring inpatient admission are treated by one of eight NHS trusts in the UK which are part of a consortium headed by the South Staffordshire and Shropshire NHS Foundation trust (SSSFT). UK based Service personnel from British Forces Germany (BFG) were treated at Guys and St Thomas Hospital in the UK up until April 2013 and from this date, at Gilead IV hospital, Bielefeld, under a contract with Soldiers, Sailors and Airmen Family Association (SSAFA) through the Limited Liability Partnership. Data is received monthly direct from SSSFT and SSAFA. See Background Quality Report for the UK Armed Forces Mental Health Report for further information on data sources found at www.gov.uk.

Primary care data, as captured on DMICP were used to compile the response where the following read codes were entered; DMS4691, DMS4692, DMS4693, DMS4698, DMS4707, DMS4708, DMS4710, DMS4711, DMS4713, DMS4714, DMS4716, DMS4717, DMS4719, DMS4720, DMS4722, DMS4723, DMS4725, DMS4726, DMS4729, EMISCSE4TK-1, TK-2, TK-4, TK-5, TK60, TK601, TRIQQIN7, U2, U200, U200-1, U200-2, U201U202-1, U204-3, U208, U208, U21, U2-1, U22, U2-3, U29, U290, U2B, U2C, U2D, U2E and U2v.

Please note if the DSH incident were recorded as free text only in the patient medical record the information does not transfer into the central data warehouse, thus was not available for analysis. It would require many hours of a clinicians time to review the patient records to code the information and thus make the information centrally available; in the timeframe required to provide the analysis, this was deemed to be disproportionate effort.

There has been no audit of the clinical accuracy of the DMICP mental health data entered in the patient record and no validation of the patient record with data held in the data warehouse.

Joint Personnel Administration System (JPA)

JPA is the system used by the Armed Forces to deal with matters of pay, leave and other personnel administrative tasks. JPA replaced a number of single-Service IT systems and was implemented in April 2006 for RAF, November 2006 for Naval Service and April 2007 for Army.

The patient data from each data source were cross referenced with the Joint Personnel Administration (JPA) system for UK Armed Forces personnel. JPA is the source for demographic information on UK Armed Forced personnel and is used to gather information on a person's service, rank, training status, gender and age.

Some demographic data i.e. gender, rank, training status and age was not available at the time of extraction for the years 2011/12 – 2013/14. This primarily affected individuals who had a DSH code entered onto DMICP in the initial days of joining the UK Armed Forces.

Methodology

UK Armed Forces population used within this bulletin included regulars (including Gurkhas and Military Provost Guard Staff), mobilised reservists, Full Time Reserve Service personnel and Non-regular Permanent Staff and trained and untrained personnel.

This bulletin presents the total number of UK Armed Forces personnel with at least one DSH event recorded after 2009 either in the MOD primary care system (DMICP) and/or the casualty notification system (NOTICAS). The data excludes all personnel recorded with DSH ideation and/or thoughts of DSH but for whom no act of actual DSH was coded in the primary care record.

All Service personnel who had a DSH event recorded from 2009 onwards were counted once over the whole time period presented. It is possible that personnel may have had subsequent new episodes of DSH however these have been excluded from the bulletin. Therefore the annual rates present a minimum count of DSH in the UK Armed Forces.

Rates

Rates enable comparisons between groups and over time, taking account of the number of personnel in a group (personnel at risk) at a particular point in time. The number of events (i.e. DSH) is divided by the number of personnel at risk and multiplied by 1,000 to calculate the rate.

In order to calculate the rates in this bulletin, an estimate of person at risk was required for the denominator. The estimate was calculated by using the average number of personnel serving in a 13 month period (e.g. the number of personnel serving at the first of every month between April 2013 and April 2014 divided by thirteen for FY 2013/2014). This methodology for calculating the estimate was in line with the method used for the UK Armed Forces Mental Health Annual report^e.

In order to understand if a difference in rates was statistically significant, 95% confidence intervals were used. Statistical significance indicates the likelihood that a finding was not due to chance. The 95% confidence interval for a rate provides the range of values within which we expect to find the real value of the indicator under study, with a probability of 95%.

If two confidence intervals do not overlap, a comparable statistical test would indicate a statistically significant difference. The rates and confidence intervals presented have been rounded to 1 decimal place and therefore when small numbers are presented the rate may lie towards one end of the confidence interval instead of more centrally between the lower and upper confidence interval.

The small number in some of the sub-group analysis may result in wide confidence intervals in the corresponding rate. The impact of this is that the range in which we expect the true value of that statistic to lie is much larger, making it harder to interpret the true underlying trend.

The information presented in this bulletin has been structured to release information into the public domain in a way that contributes to the MOD accountability to the public but which doesn't risk breaching individual's rights to medical confidentiality. In line with Joint Service Publication (JSP) 200 and in keeping with the Office for National Statistics Guidelines, all numbers less than five have been suppressed and presented as '~' to prevent the inadvertent disclosure of individual identities. Where there is only one cell in a row or column that is less than five, the next smallest number (or numbers where there are tied values) has also been suppressed so that numbers cannot simply be derived from totals.

 $^{\text{e}} \ \underline{\text{https://www.gov.uk/government/statistics/uk-armed-forces-mental-health-annual-statistics-financial-year-} \underline{201415}$

6

Glossary

Army - The British Army consists of the General Staff and the deployable Field Army and the Regional Forces that support them, as well as Joint elements that work with the Royal Navy and Royal Air Force. Its primary task is to help defend the interests of the UK.

Confidence Interval - The 95% confidence interval for a rate provides the range of values within which we expect to find the real value of the indicator under study, with a probability of 95%.

Deliberate Self Harm – includes incidents of self-injury (SI) and self-poisoning within this bulletin. It excludes personnel who had thoughts of deliberate self-harm or suicidal ideation.

Defence Medical Information Capability Programme (DMICP) - The DMICP programme commenced during 2007 and comprises an integrated primary Health Record (iHR) for clinical use and a pseudo-anonymised central data warehouse.

MOD Specialist Mental Health Services - encompass the delivery of care through MOD's Department for Community Mental Health (DCMH) for outpatient care, and all admissions to the MOD's in-patient care contractor. It does not cover mental health care for patients treated wholly in the primary care setting by GPs.

New Case of DSH – all UK Armed Forces personnel who had DSH event recorded from 2009 onwards were counted once over the whole time period presented.

NOTICAS – Notification of Casualty (or "NOTICAS") is the name for the formalised system of reporting casualties within the UK Armed Forces.

Officer - An officer is a member of the Armed Forces holding the Queen's Commission to lead and command elements of the forces. Officers form the middle and senior management of the Armed Forces. This includes ranks from Sub-Lt/2nd Lt/Pilot Officer up to Admiral of the Fleet/Field Marshal/Marshal of the Royal Air Force, but excludes Non-Commissioned Officers.

Other Ranks - Other ranks are members of the Royal Marines, Army and Royal Air Force who are not officers but Other Ranks include Non-Commissioned Officers.

Royal Air Force (RAF). The Royal Air Force (RAF) is the aerial defence force of the UK.

Royal Marines (RM) Royal Marines are sea-going soldiers who are part of the Naval Service. RM officer ranks were aligned with those of the Army on 1 July 1999.

Royal Navy (RN) The sea-going defence forces of the UK but excludes the Royal Marines and the Royal Fleet Auxiliary Service (RFA).

UK Armed Forces population - is defined as the number of serving UK Armed Forces personnel.

UK Armed Forces - are full time Service personnel, including Nursing Services and Gurkhas, Naval activated Reservists, mobilised Reservists, Military Provost Guarding Service (MPGS), Non Regular Permanent Service (NRPS) and reservist personnel. Unless otherwise stated, includes trained and untrained personnel.

Further information

Contact us

Defence Statistics welcome feedback on our statistical products. If you have any comments or questions about this publication or about our statistics in general, you can contact us as follows:

Defence Statistics (Health)

Telephone: 030679 84423

Email: DefStrat-Stat-Health-PQ-FOI@mod.uk

If you require information which is not available within this or other available publications, you may wish to submit a Request for Information under the Freedom of Information Act 2000 to the Ministry of Defence. For more information, see:

https://www.gov.uk/make-a-freedom-of-information-request/the-freedom-of-information-act

Other contact points within Defence Statistics are:

Defence Expenditure Analysis	030 679 34531	DefStrat-Econ-ESES-DEA-Hd@mod.uk					
Price Indices	030 679 32100	DefStrat-Econ-ESES-PI-Hd@mod.uk					
Naval Service Manpower	023 9254 7426	DefStrat-Stat-Navy-Hd@mod.uk					
Army Manpower	01264 886175	DefStrat-Stat-Army-Hd@mod.uk					
RAF Manpower	01494 496822	DefStrat-Stat-Air-Hd@mod.uk					
Tri-Service Manpower	020 7807 8896	DefStrat-Stat-Tri-Hd@mod.uk					
Civilian Manpower	020 7218 1359	DefStrat-Stat-Civ-Hd@mod.uk					
Health Information	030 6798 4423	DefStrat-Stat-Health-Hd@mod.uk					

If you wish to correspond by mail, our postal address is:

Defence Statistics (Health) Ministry of Defence Abbey Wood North Oak 0 West #6028 Bristol

BS34 8JH

For general MOD enquiries, please call: 020 7218 9000