Routine childhood immunisations from September 2012

| When to immunise | Diseases protected against | | | Vaccine given | Immunisation site** |
|---|--|------|-----|---|------------------------|
| Two months old | Diphtheria, tetanus, pertussis, polio and Haemophilus | | | DTaP/IPV/Hib (Pediacel) | Thigh |
| | Withdray | vn A | ٩ug | ust 2015 | |
| Three months old | polio and Hib | | | (Pediacel) | rriign |
| | Meningococcal group C disease (MenC) | | | MenC (Menjugate or Neisvac C) | Thigh |
| Four months old | Diphtheria, tetanus, pertussis, polio and Hib | | | DTaP/IPV/Hib (Pediacel) | Thigh |
| | MenC | | | MenC (Menjugate or Neisvac C) | Thigh |
| | Pneumococcal disease | | | PCV (Prevenar 13) | Thigh |
| Between 12 and 13 months old – within a month of the first birthday | Hib/MenC | | | Hib/MenC (Menitorix) | Upper arm/thigh |
| | Pneumococcal disease | | | PCV (Prevenar 13) | Upper arm/thigh |
| | Measles, mumps and rubella (German measles) | | | MMR (Priorix or MMR VaxPRO) | Upper arm/thigh |
| Three years four months old or soon after | Diphtheria, tetanus, pertussis and polio | | | dTaP/IPV (Repevax) or DTaP/IPV (Infanrix-IPV) | Upper arm |
| | Measles, mumps and rubella | | | MMR (Priorix or MMR VaxPRO) (check first dose has been given) | Upper arm |
| Girls aged 12 to 13 years old | Cervical cancer caused by human papillomavirus types 16 and 18 (and genital warts caused by types 6 and 11) | | | Gardasil | Upper arm |
| Thirteen to 18 years old | Tetanus, diphtheria and polio | | | Td/IPV (Revaxis), and check MMR status | Upper arm |

^{**} Where two or more injections are required at once, these should ideally be given in different limbs. Where this is not possible, injections in the same limb should be given 2.5cm apart. For more details see Chapters 4 and 11 in the Green Book.

Non-routine immunisations for at-risk babies

| At birth, 1 month old, 2 months old and 12 months old | Hepatitis B | Нер В | Thigh |
|---|--------------|-------|-------------------------|
| At birth | Tuberculosis | BCG | Upper arm (intradermal) |



*NB Where a vaccine is manufactured by more than one supplier, it may, on occasion be necessary to substitute an alternative brand. All vaccines for use in the routine childhood programme can be ordered free of charge at www.immform.dh.gov.uk





404119 © Crown copyright 2010 1p 2k Dec10 (MRP)