



Local Tobacco Control Profiles Indicators at a glance June 2017

Notes:

- Changes are only shown if they are statistically significant. An upwards arrow (↑, ↑ or ↑) represents a significant increase in the indicator value, a downwards arrow (↓, ↓ or ↓) represents a significant decrease. A sideways arrow (↔) is displayed if there has been no significant change.
- The arrows are coloured green and red for those indicators where a change can be described as improving or worsening respectively.
- A blue dash (-) signifies that there is only one data point and therefore no comparison can be made with previous data. A blue asterisk (*) is used where data should not be compared with previous time periods.

Domain: smoking prevalence in adults				
Indicator	Period	Value	Unit	Change from previous
Smoking Prevalence in adults - current smokers (APS)	2016	15.5	%	↓
Smoking Prevalence in adults - ex smokers (APS)	2016	26.2	%	*
Smoking Prevalence in adults - never smoked (APS)	2016	58.3	%	*
Smoking Prevalence in adults in routine and manual occupations - current smokers (APS)	2016	26.5	%	↓
Smoking prevalence in adults in routine and manual occupations - ex smokers (APS)	2016	21.9	%	*
Smoking prevalence in adults in routine and manual occupations - never smoked (APS)	2016	51.5	%	*
Smoking Prevalence in adults - current smokers (IHS)	2014	18.0	%	↓
Smoking Prevalence in adults - ex-smokers (IHS)	2014	33.9	%	↔
Smoking Prevalence in adults - never smoked (IHS)	2014	48.1	%	↔
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	28.0	%	↔
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014	30.8	%	↔
Smoking prevalence in adults in routine and manual occupations - never smoked (IHS)	2014	41.2	%	↔
GPPS smoking prevalence - current smokers	2015/16	16.4	%	↔
GPPS smoking prevalence - ex smoker	2015/16	27.5	%	↔
GPPS smoking prevalence - never smoked	2015/16	56.1	%	↔
Smoking prevalence in adults - current smokers (QOF)	2015/16	18.1	%	↓
Smoking prevalence in adults with serious mental illness (SMI)	2014/15	40.5	%	-
Smoking status at time of delivery	2015/16	10.6	%	↓

Domain: smoking prevalence in young people

Indicator	Period	Value	Unit	Change from previous
Smoking prevalence age 15 years - regular smokers (SDD survey)	2014	8.0	%	↔
Smoking prevalence age 15 years - occasional smokers (SDD survey)	2014	5.0	%	↓
Smoking prevalence at age 15 - current smokers (WAY survey)	2014/15	8.2	%	-
Smoking prevalence at age 15 - regular smokers (WAY survey)	2014/15	5.5	%	-
Smoking prevalence at age 15 - occasional smokers (WAY survey)	2014/15	2.7	%	-
Use of e-cigarettes at age 15 years (WAY survey)	2014/15	18.4	%	-
Use of other tobacco products at age 15 years (WAY survey)	2014/15	15.2	%	-
Smoking prevalence at ages 11-15 - regular smokers (modelled estimates)	2009 - 12	3.1	%	-
Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates)	2009 - 12	1.4	%	-
Smoking prevalence at age 15 - regular smokers (modelled estimates)	2009 - 12	8.7	%	-
Smoking prevalence at age 15 - occasional smokers (modelled estimates)	2009 - 12	3.9	%	-
Smoking prevalence at ages 16-17 - regular smokers (modelled estimates)	2009 - 12	14.7	%	-
Smoking prevalence at ages 16-17 - occasional smokers (modelled estimates)	2009 - 12	5.8	%	-

Domain: smoking related mortality

Indicator	Period	Value	Unit	Change from previous
Smoking attributable mortality	2013 - 15	283.5	DSR per 100,000	↑
Deaths from lung cancer	2013 - 15	58.7	DSR per 100,000	↓
Deaths from chronic obstructive pulmonary disease	2013 - 15	52.6	DSR per 100,000	↑
Smoking attributable deaths from heart disease	2013 - 15	28.9	DSR per 100,000	↓
Smoking attributable deaths from stroke	2013 - 15	9.4	DSR per 100,000	↔
Potential years of life lost due to smoking related illness	2013 - 15	1399.1	DSR per 100,000	↓
Stillbirth rate	2013 - 15	4.6	Crude rate per 1,000	↔
Neonatal mortality	2013 - 15	2.7	Crude rate per 1,000	↔

Domain: smoking related ill health

Indicator	Period	Value	Unit	Change from previous
Premature births (<37 weeks gestation)	2013 - 15	78.4	Crude rate per 1,000	↑
Low birth weight of term babies	2015	2.8	%	↓
Hospital admissions for asthma (under 19 years)	2015/16	202.4	DSR per 100,000	↓
Smoking attributable hospital admissions	2015/16	1726.4	DSR per 100,000	↑
Cost per capita of smoking attributable hospital admissions	2011/12	38.0	£ per capita	*
Emergency hospital admissions for COPD	2015/16	411.2	DSR per 100,000	↔
Lung cancer registrations	2013 - 15	78.9	DSR per 100,000	↔
Oral cancer registrations	2013 - 15	14.5	DSR per 100,000	↔

Domain: impact of smoking

Indicator	Period	Value	Unit	Change from previous
Indicative tobacco sales figures (£ millions)	2013	15446.1	£	*
Illicit tobacco: tax gap (£ millions)	2015/16	1600.0	£	↔
Illicit tobacco: share of cigarette sales (%)	2015/16	13.0	%	↔
Fatalities from accidental fires ignited by smoking materials	2015/16	41.4	%	↔
Accidental fires ignited by smoking related materials	2015/16	7.8	%	↑
Attitudes to smoking in 15 year olds - 'smoking causes harm to others' (WAY survey)	2014/15	90.9	%	-

Domain: smoking quitters

Indicator	Period	Value	Unit	Change from previous
No. setting a quit date	2015/16	5091.8	Crude rate per 100,000 smokers	↓
Successful quitters at 4 weeks	2015/16	2598.1	Crude rate per 100,000 smokers	↓
Successful quitters (CO validated) at 4 weeks	2015/16	1854.2	Crude rate per 100,000 smokers	↓
Completeness of NS-SEC recording by Stop Smoking Services	2015/16	91.1	%	↑
Cost per quitter	2015/16	479.0	£ per capita	*