



Public Health
England

Protecting and improving the nation's health

Laboratory surveillance of uncommon pathogens causing bacteraemia in England, Wales and Northern Ireland: 2016

Health Protection Report
Volume 15 Number 45
15 December 2017

Laboratory surveillance of uncommon pathogens causing bacteraemia in England, Wales and Northern Ireland: 2016

The analysis presented in this report is based on data extracted from the Public Health England (PHE) voluntary surveillance database (Second Generation Surveillance System) on 31 October 2017 for the period between 1 January 2012 and 31 December 2016. Data from Wales and Northern Ireland were extracted separately from DataStore (on 24 February 2017) and CoSurv (on 28 September 2017), respectively. This report describes uncommon pathogens causing clinically significant infections (bacterial genera with fewer than 50 reports in 2016) identified from blood cultures. Data in this report may differ slightly from data in earlier publications due to inclusion of late reports and reclassification of organisms.

There were 4,037 cases of bacteraemia caused by uncommon bacterial genera reported between 2012 to 2016. From these cases 153 uncommon genera were identified, 56.3% of which were Gram-negative. Of these episodes of bacteraemia in 2016 (n=1,228), 55.7% were due to Gram-negative bacteria (see the [web appendix](#) associated with this report). By definition of inclusion in this analysis, small numbers of reports preclude robust or meaningful analysis of trends, but of note are continuing decreases between 2012 and 2016 in reports of the Gram-negative genera *Kluyvera* and *Sphingobacterium*. In contrast, an increase was noted for the following Gram-positive genera: *Abiotrophia*, *Actinotignum*, *Anaerococcus*, *Arthrobacter*, *Atopobium*, *Dermacoccus*, *Eggerthella*, *Finegoldia*, *Globicatella*, *Gordonia*, *Microbacterium*, *Solobacterium*; and Gram-negative *Aggregatibacter*, *Brucella*, *Capnocytophaga*, *Delftia*, *Dialister*, *Eikenella*, *Kingella*, *Leptotrichia*, *Myroides*, *Parabacteroides*, *Paracoccus*, and *Sphingomonas* species.

Discussion

The purpose of this review is to describe unusual bacterial genera not routinely included in the monthly bacteraemia reports published in the Health Protection Report.

Examining trends in these pathogens can provide a means of identifying emerging or re-emerging infections [1], providing an alert to allow opportunities for preventative measures or education of frontline clinical staff.

There has been a general improvement in the identification of cultured organisms by increased use of automated biochemical identification systems, molecular techniques such as 16S ribosomal RNA, and the introduction of MALDI-TOF mass spectrometry in some laboratories. This has increased the accuracy of species identified, and we now have a better understanding of the relative importance of these hitherto difficult to identify species causing significant disease. However, there are concerns that MALDI-TOF cannot currently be used for accurate speciation of *Elizabethkingia* and accurate speciation within the *Elizabethkingia* genus requires sequence-based methods [2]. It should be borne in mind that findings by MALDI-TOF reflect organisms that are present in the database, therefore identification is expected to improve with expansion of the database.

Although these bacteria only account for a very low proportion of total bloodstream infections, they can be associated with important clinical consequences, such as endocarditis [3].

Abiotrophia defectiva can be isolated from the oral cavity and bacteraemia may follow dental treatment, potentially leading to endocarditis [4]. *Capnocytophaga canimorsus* are also part of the normal oral microflora of humans and dogs and bloodstream infections have been described following wounds or bites from dogs [5].

Actinotignum schaalii (previously *Actinobaculum schaalii* [6]) bacteraemia have been associated with urinary tract infection in the elderly [7] whereas *Kingella kingae* bacteraemia have been reported more frequently in children than adults [8].

Infections imported from endemic regions, such as *Brucella* species [9] although rarely diagnosed in this country can cause severe illness in those affected. Other bacteraemia

have been associated with the use of catheters in patients with underlying morbidity or who are immunocompromised e.g. *Delftia acidovorans* [10].

A number of genera were reported in 2016 that were not seen in the previous 4 years: *Actinotignum*, *Agromyces*, *Curtobacterium*, *Deinococcus*, *Dietzia*, *Erwinia*, *Geobacillus*, *Leifsonia*, *Lysinibacillus*, *Murdochiella*, *Oceanobacillus*, *Pseudoclavibacter*, *Salana*, *Weissella* (Gram-positive) and *Kosakonia*, *Megasphaera*, *Methylopila*, *Pseudoflavonifractor*, *Pseudoxanthomonas*, *Rhodobacter*, *Selenomonas*, *Suttonella*, *Xanthomonas* (Gram-negative).

Whilst the bloodstream infections reported to this voluntary surveillance system should, according to national reporting guidelines, reflect clinically significant disease, some of these reports may reflect skin colonisers or contaminants due to difficulties in blood culture sampling or contamination in laboratory processing [11].

If confirmation of unusual bacterial pathogens is required, isolates can be sent to the relevant laboratory within the Bacteriology Reference Department, National Infection Service, Colindale, Public Health England.

Acknowledgements

These reports would not be possible without the weekly contributions from microbiology colleagues in laboratories across England, Wales and Northern Ireland, without which there would be no surveillance data. In addition, the support from colleagues within Public Health England, Public Health Wales and HSC Public Health Agency (Northern Ireland) in particular, is valued in the preparation of the report. Please send any comments or feedback to: hcai.amrdepartment@phe.gov.uk.

References

1. Al Akhrass F, Al Wohoush I, Chaftari AM, Reitzel R, Jiang Y, Ghannoum M, et al (2012). *Rhodococcus* bacteremia in cancer patients is mostly catheter related and associated with biofilm formation. *PLoS ONE* **7**(3).
2. Kenna DTD, Fuller A, Martin K, Perry C, Pike R, Burns PJ, et al (2017). *rpoB* gene sequencing highlights the prevalence of an *E. miricola* cluster over other *Elizabethkingia* species among UK cystic fibrosis patients. *Diagn Microbiol Infect Dis* (online, 26 Oct).
3. Brouqui P, Raoult D (2001). Endocarditis due to rare and fastidious bacteria. *Clin Microbiol Rev* **14**(1):177-207 (Jan).
4. Kiernan TJ, O'Flaherty N, Gilmore R, Ho E, Hickey M, Tolan M, et al (2008). *Abiotrophia defectiva* endocarditis and associated hemophagocytic syndrome--a first case report and review of the literature. *Int J Infect Dis* **12**(5):478-82 (Sep).
5. van Dam AP, Jansz A (2011). *Capnocytophaga canimorsus* infections in the Netherlands: a nationwide survey. *Clin Microbiol Infect* **17**(2): 312-5 (Feb).
6. Yassin AF, Sproer C, Pukall R, Sylvester M, Siering C, Schumann P (2015). Dissection of the genus *Actinobaculum*: Reclassification of *Actinobaculum schaalii* Lawson et al. (1997) and *Actinobaculum urinale* Hall et al. (2003) as *Actinotignum schaalii* gen. nov., comb. nov. and *Actinotignum urinale* comb. nov., description of *Actinotignum sanguinis* sp. nov. and emended descriptions of the genus *Actinobaculum* and *Actinobaculum suis*; and re-examination of the culture deposited as *Actinobaculum massiliense* CCUG 47753T (= DSM 19118T), revealing that it does not represent a strain of this species. *Int J Syst Evol Microbiol* **65**(Pt 2): 615-24 (Feb).
7. Gomez E, Gustafson DR, Rosenblatt JE, Patel R (2011). *Actinobaculum* bacteremia: a report of 12 cases. *J Clin Microbiol* **49**(12): 4311-3 (Dec).
8. Yagupsky P. *Kingella kingae*: carriage, transmission, and disease (2015). *Clin Microbiol Rev* **28**(1): 54-79 (Jan).
9. Ramin B, Macpherson P (2010). Human brucellosis. *BMJ* **10**; 341:c4545 (Sep).
10. Chotikanatis K, Backer M, Rosas-Garcia G, Hammerschlag MR (2011). Recurrent intravascular-catheter-related bacteremia caused by *Delftia acidovorans* in a hemodialysis patient. *J Clin Microbiol* **49**(9): 3418-21 (Sep).
11. Melish M, Campbell KA (1998). Coagulase-positive Staphylococcal infections. In: Feigin RD, Cherry JD, editors. Textbook of Paediatric Infectious Diseases: WB Saunders (Philadelphia USA), pp. 1039-65.

About Public Health England

Public Health England exists to protect and improve the nation's health and wellbeing, and reduce health inequalities. We do this through world-class science, knowledge and intelligence, advocacy, partnerships and the delivery of specialist public health services. We are an executive agency of the Department of Health, and are a distinct delivery organisation with operational autonomy to advise and support government, local authorities and the NHS in a professionally independent manner.

About Health Protection Report

Health Protection Report is a national public health bulletin for England and Wales, published by Public Health England. It is PHE's principal channel for the dissemination of laboratory data relating to pathogens and infections/communicable diseases of public health significance and of reports on outbreaks, incidents and ongoing investigations.

Public Health England, Wellington House, 133-155 Waterloo Road, London SE1 8UG

Tel: 020 7654 8000 www.gov.uk/phe

Twitter: [@PHE_uk](https://twitter.com/PHE_uk) Facebook: www.facebook.com/PublicHealthEngland

Queries relating to this document should be directed to:

HCAI-AMR Department, National Infection Service,

PHE Colindale,

61 Colindale Avenue, London NW9 5EQ.

immunisation@phe.gov.uk

© Crown copyright 2017

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](https://www.ogcl.gov.uk) or email psi@nationalarchives.gsi.gov.uk. Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

Published December 2017

PHE publications gateway number: 2017661

