

GP OOHSS

GP Out-of-Hours Surveillance System: England

Data to: 24 September 2017

26 September 2017 Year: 2017 Week: 38

In This Issue:

Key Messages.

Weekly summary.

Total contacts.

Syndromic indicators.

Notes and caveats.

Further information.

Acknowledgements.

Key messages

GP out of hours consultations for respiratory indicators continued to rise during week 38, particularly in children aged 0-14 years (figures 2a & 5a).

Syndromic indicators at a glance:


Number of contacts and percentage of Read coded contacts.

Key indicator	No. of contacts	% Week 38	% Week 37	Trend*
All OOH contacts, all causes	159,100			
Acute respiratory infection	11,206	13.57	11.61	^
Influenza-like illness	123	0.15	0.10	^
Bronchitis/bronchiolitis	154	0.19	0.17	^
Difficulty breathing/wheeze/asthma	2,142	2.59	2.24	^
Pharyngitis	70	0.08	0.09	←→
Gastroenteritis	3,146	3.81	3.92	←→
Diarrhoea	777	0.94	1.09	$oldsymbol{\Psi}$
Vomiting	1,179	1.43	1.44	←→
Myocardial infarction	853	1.03	1.05	←→

^{*}Trend: reports on the trend seen over previous weeks in the percentage of Read coded contacts.

1: Total out-of-hours contacts:

Daily total number of out-of-hours and unscheduled contacts and 7 day average (adjusted for bank holidays).





2: Acute Respiratory Infection daily contacts.

35


30


Percentage of Read coded contacts

10

5

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.






3: Influenza-like illness daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.


*7-day moving average adjusted for bank holidays.






4: Bronchitis/ bronchiolitis daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.





5: Difficulty breathing/ wheeze/asthma daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.



5a: Difficulty
breathing/wheeze/
asthma daily
contacts by age
group.







Intentionally left blank


6: Acute pharyngitis and persistent sore throat.


Shown as a percentage of the total contacts with a Read code and as a 7 day average*.

7: Gastroenteritis daily contacts


Shown as a percentage of the total contacts with a Read code and as a 7 day average*.

*7-day moving average adjusted for bank holidays.






7a: Gastroenteritis weekly contacts by age group.




8: Diarrhoea daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.



8a: Diarrhoea weekly contacts by age group.




*7-day moving average adjusted for bank holidays.




9: Vomiting daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.




9a: Vomiting weekly contacts by age group.



10: Myocardial Infarction daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 day average*.



*7-day moving average adjusted for bank holidays.



Notes and caveats:

- This bulletin presents data from the Public Health England (PHE) GP Out
 -of-hours\Unscheduled Care Surveillance System (GP OOHSS).
- Fully anonymised data from GP out-of-hours (OOH) and unscheduled care service providers in England are being transferred to the PHE for analysis and interpretation by the PHE Real-time Syndromic Surveillance Team (ReSST).
- This system supplements existing PHE syndromic surveillance systems by monitoring data on general practitioner consultations outside of routine surgery opening times (evenings, weekends and bank holidays) and unplanned contacts within NHS primary care.
- The key indicators presented within this bulletin are derived by grouping selected Read coded consultations.
- GP OOH consultation data are analysed on a daily basis to identify
 national and regional trends. A statistical algorithm underpins each
 system, routinely identifying activity that has increased significantly or is
 statistically significantly high for the time of year. Results from these daily
 analyses are assessed by the ReSST, along with analysis by age group,
 and anything deemed of public health importance is alerted by the team.
- Baselines represent seasonally expected levels of activity and are constructed from historical data. Furthermore, they take into account any known substantial changes in data collection, population coverage or reporting practices. Baselines are refreshed using the latest data on a regular basis.

Further information:

The GP Out-of-Hours Surveillance System Bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance:

https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses

Acknowledgements:

We are grateful to Advanced Health and Care and the GP out-of-hours and unscheduled care service providers who have kindly agreed to participate in this system.

PHE Out-of-Hours/Unscheduled Care Surveillance

Produced by: PHE Real-time Syndromic Surveillance Team 6th Floor, 5 St Philip's Place, Birmingham, B3 2PW

Tel: 0344 225 3560 > Option 4 > Option 2 Fax: 0121 236 2215

Web: https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses