



Weekly All-cause Mortality Surveillance

07 May 2015 – Week 19 report (up to week 18 data)

In week 18 2015, no statistically significant excess all-cause mortality by week of death was seen through the EuroMOMO algorithm in England overall and by age group and across the devolved administrations. Since week 40 2014, significant excess mortality has been observed in England in weeks 50-7 predominantly in 65+ year olds, peaking in week 2 2015. This period of significant excess coincided with circulating influenza and cold snaps.

Excess overall all-cause mortality, England and Wales

-In week 17 2015, an estimated 10,599 all-cause deaths were registered in England and Wales (source: [Office for National Statistics](#)). This is less than the 11,639 estimated death registrations in week 16, and is just above the 95% upper limit of expected death registrations for the time of year as calculated by PHE (Figure 1). Weeks 52, 1 and 14 correspond to a week when there were bank holidays and fewer days when deaths were registered. Therefore the decrease in deaths seen is likely to be artificial and result in subsequent increases in following weeks.

Excess all-cause mortality in subpopulations, UK

-Since week 40 2014 up to week 18 2015 in England, excess mortality by date of death above the upper 2 z-score threshold was seen in England after correcting ONS disaggregate data for reporting delay with the standardised [EuroMOMO](#) algorithm in 65+ year olds in weeks 50-7 2015, 15-64 year olds in weeks 51-2, and weeks 2 and 5 in under five year olds (Figure 2, Table 1). This period of statistically significant excess coincided with circulating influenza and cold snaps. This data is provisional due to the time delay in registration; numbers may vary from week to week.

-In the devolved administrations, up to week 18 2015, excess mortality above the threshold was seen in weeks 51-4 and 6-9 in Scotland, weeks 42 and 1-3 in Wales and weeks 3-4 and 8-9 in Northern Ireland (Table 2).

Figure 1: Observed & predicted all-cause death registrations, E&W




Table 1: Excess mortality by age group, England*

Age group (years)	Excess detected in week 18 2015?	Weeks with excess in 2014/15
<5	x	2,5
5-14	x	NA
15-64	x	51-2
65+	x	50-7

* Excess mortality is calculated as the observed minus the expected number of deaths in weeks above threshold

Figure 2: Excess mortality in 65+ year olds by week of death, EuroMOMO, England




Table 2: Excess mortality by UK country*

Country	Excess detected in week 18 2015?	Weeks with excess in 2014/15
England	x	50-7
Wales	x	42, 1-3
Scotland	x	51-4, 6-9
Northern Ireland	x	3-4, 8-9

* Excess mortality is calculated as the observed minus the expected number of deaths in weeks above threshold

NB. Separate total and age-specific models are run for England which may lead to discrepancies between Tables 1 + 2

Produced by the Respiratory Diseases Department, Public Health England.

- Seasonal mortality is seen each year in England and Wales, with a higher number of deaths in winter months compared to the summer. Additionally, peaks of mortality above this expected higher level typically occur in winter, most commonly the result of factors such as cold snaps and increased circulation of respiratory viruses, in particular influenza.
- RDD's weekly mortality surveillance aims to detect and report acute significant weekly excess mortality above normal seasonal levels in a timely fashion. Excess mortality is defined as a significant number of deaths reported over that expected for a given point in the year, allowing for weekly variation in the number of deaths. This triggers further investigation of spikes and informs any public health responses.
- The aim is not to assess general mortality trends or precisely estimate the excess attributable to different factors, although some end-of-winter estimates and more in-depth analyses (by age, geography etc.) are undertaken.
- Separate to the calculations presented in this report, excess winter deaths (EWD), comparing the number of deaths in the winter period compared to the non-winter period, are calculated by [ONS](#) and presented in an [atlas](#) down to local authority level.