




Weekly All-cause Mortality Surveillance

4 December 2014 – Week 49 report (up to week 48 data)

In week 49 2014, no excess all-cause mortality by week of death was seen across the UK through the EuroMOMO algorithm.

Excess overall all-cause mortality, England and Wales

-In week 47 2014, an estimated 9,472 all-cause deaths were registered in England and Wales (source: [Office for National Statistics](#)). This is less than the 10,036 estimated death registrations in week 46 and remains below the 95% upper limit of expected death registrations for this time of year as calculated by PHE (Figure 1).



Excess all-cause mortality in subpopulations, UK

-In week 48 2014, no excess mortality by date of death above the upper 2 z-score threshold was seen in 65+ year olds in England after correcting ONS disaggregate data for reporting delay with the standardised [EuroMOMO](#) algorithm (Figure 2, Table 1), in other age groups or subnationally. This data is provisional due to the time delay in registration and so numbers may vary from week to week.

-No excess mortality above the threshold through the same standardised algorithm was seen across Wales, Scotland or Northern Ireland in week 48 (Table 2).

Table 1: Excess mortality by age group, England*

Age group (years)	Excess detected in week 48 2014?	Weeks with excess in 2014/15
<5	x	NA
5-14	x	NA
15-64	x	NA
65+	x	NA


* Excess mortality is calculated as the observed minus the expected number of deaths in weeks above threshold

Table 2: Excess mortality by UK country*

Country	Excess detected in week 48 2014?	Weeks with excess in 2014/15
England	x	NA
Wales	x	NA
Scotland	x	NA
Northern Ireland	x	NA

* Excess mortality is calculated as the observed minus the expected number of deaths in weeks above threshold

NB. Separate total and age-specific models are run for England which may lead to discrepancies between Tables 1 + 2



Produced by the Respiratory Diseases Department, Public Health England.

- Seasonal mortality is seen each year in England and Wales, with a higher number of deaths in winter months compared to the summer. Additionally, peaks of mortality above this expected higher level typically occur in winter, most commonly the result of factors such as cold snaps and increased circulation of respiratory viruses, in particular influenza.
- RDD's weekly mortality surveillance aims to detect and report acute significant weekly excess mortality above normal seasonal levels in a timely fashion. Excess mortality is defined as a significant number of deaths reported over that expected for a given point in the year, allowing for weekly variation in the number of deaths. This triggers further investigation of spikes and informs any public health responses.
- The aim is not to assess general mortality trends or precisely estimate the excess attributable to different factors, although some end-of-winter estimates and more in-depth analyses (by age, geography etc.) are undertaken.
- Separate to the calculations presented in this report, excess winter deaths (EWD), comparing the number of deaths in the winter period compared to the non-winter period, are calculated by [ONS](#) and presented in an [atlas](#) down to local authority level.