



Public Health
England

Protecting and improving the nation's health

Laboratory confirmed cases of pertussis (England): January to March 2018

Health Protection Report
Volume 12 Number 22
22 June 2018

Laboratory confirmed cases of pertussis (England): January to March 2018

In England there were 644 laboratory confirmed cases of pertussis (culture, PCR, serology or oral fluid) reported to the Public Health England (PHE) pertussis enhanced surveillance programme in the first quarter of 2018, from January to March 2018 (table 1). Total cases were 28% lower than those reported in the same quarter of 2017 (896 cases) and 49% lower than the 1264 cases reported in the first quarter of 2016.

A national outbreak of pertussis [1] was declared by the HPA in April 2012 and, as a response to the ongoing outbreak, the Department of Health (DH) introduced a temporary immunisation programme for pregnant women from October 2012 [2]. In June 2014 the Joint Committee on Vaccination and Immunisation (JCVI) recommended that the programme should continue for a further five years [3] based on UK evidence of impact, high effectiveness and safety and continuing high levels of disease [4,5,6,7]. From 1 April 2016 the recommended gestational age for vaccination was revised to between 20-32 weeks but can be given as early as 16 weeks [3].

Following the peak in 2012 an overall decrease in pertussis was observed between 2013 and 2015. A relative increase in pertussis activity occurred in 2016 consistent with pre-existing epidemiological trends of 3-4 yearly cyclical peaks (Figure 1).

In the first quarter of 2018, the greatest number of laboratory confirmed cases in England continues in individuals aged 15 years and over although the highest disease incidence persists in infants <3 months. Pertussis activity in all infants <1 year of age was lower in the first of 2018 (16 cases) than the equivalent periods in 2012 to 2017 (table 2).

Confirmed cases aged 6-11 months were higher (34 cases) in 2016 than in any year since the introduction of enhanced surveillance in 1994. Laboratory confirmed cases in this age group were 50% lower (17 cases) in 2017 and three confirmed cases were reported in the first quarter of 2018. This infant age group is known to have high levels of protection after completion of the primary immunisation programme.

Overall activity remains higher in all age groups from 1 year and older, relative to years preceding the pre-2012 peak. Ascertainment in those aged 5 to <17 years has improved with availability of oral fluid testing since 2013. From 1 May 2018 the availability of oral fluid testing was extended to all children aged 2 to <17 years. (See the guidelines for the public health management of pertussis for [8] for details of appropriate laboratory investigation of suspected cases of pertussis which is affected by the age of the suspect case and time since onset of their symptoms).

Pertussis vaccine coverage for pregnant women averaged 73.6% across October and December, 1.4% lower than coverage for the same period in 2016 but continuing at the higher levels seen since April 2016 [9]. Extended eligibility criteria for the vaccine may have contributed to the increase in uptake observed over the last couple of years [10].

There have been no reported deaths in infants with pertussis confirmed between January and March 2018 and there were no deaths in infants with pertussis confirmed in 2017. Of the eighteen infants who have died following confirmed pertussis disease and who were born after the introduction of the maternal programme (on 1 October 2012), 16 were born to mothers who had not been immunised against pertussis during pregnancy.

Surveillance data in young infants following the introduction of the pertussis immunisation in pregnancy programme continues to demonstrate that a relatively low incidence has been maintained in this age group, with expected seasonal increases. It is important to be aware, however, that raised levels of pertussis persist in groups aged 1 year and older. Women should continue to be supported in accessing immunisation against pertussis during pregnancy (ideally between 20-32 weeks) to optimise protection for their babies from birth.

Table 1. Laboratory-confirmed cases of pertussis by age and testing method in England, January – March 2018.

| Age group | Culture* | PCR | Serology | Oral fluid only | Total |
|--------------|-----------|-----------|------------|-----------------|------------|
| <3 months | 2 | 7 | 0 | 0 | 9 |
| 3-5 months | 0 | 4 | 0 | 0 | 4 |
| 6-11 months | 1 | 2 | 0 | 0 | 3 |
| 1-4 years | 0 | 1 | 10 | 2 | 13 |
| 5-9 years | 1 | 0 | 20 | 7 | 28 |
| 10-14 years | 0 | 1 | 49 | 15 | 65 |
| 15+ years | 6 | 2 | 510 | 4 | 522 |
| Total | 10 | 17 | 589 | 28 | 644 |

* Culture confirmed cases may additionally have tested positive by any other method, PCR confirmed cases may have additionally tested positive by serology or OF and serology confirmed cases may also have been confirmed by OF. Submission of all presumptive *B. pertussis* isolates is encouraged for confirmation of identity and to allow further characterisation for epidemiological purposes.

Figure 1. Total number of laboratory-confirmed pertussis cases per quarter in England, 2008 – 2018 (Q1).




Table 2: Laboratory-confirmed cases of pertussis by age and year England, January to March: 2012 – 2018

| Age group | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|--------------|------------|-------------|------------|------------|-------------|------------|------------|
| <3 months | 70 | 26 | 12 | 16 | 35 | 31 | 9 |
| 3-5 months | 11 | 7 | 5 | 6 | 9 | 6 | 4 |
| 6-11 months | 2 | 0 | 3 | 2 | 5 | 0 | 3 |
| 1-4 years | 4 | 20 | 6 | 14 | 15 | 11 | 13 |
| 5-9 years | 13 | 29 | 24 | 39 | 77 | 37 | 28 |
| 10-14 years | 98 | 175 | 79 | 82 | 121 | 89 | 65 |
| 15+ years | 504 | 1368 | 473 | 622 | 1002 | 722 | 522 |
| Total | 702 | 1625 | 602 | 781 | 1264 | 896 | 644 |

References

1. *HPR* 6(15), 13 April 2012.
2. Department of Health: Pregnant women to be offered whooping cough vaccine (news story, 28 September 2012).
3. Joint Committee on Vaccination and Immunisation.
4. G Amirthalingam, N Andrews, H Campbell, S Ribeiro, E Kara, K Donegan, et al (2014). Effectiveness of maternal pertussis vaccination in England: an observational study. *The Lancet*.
5. Dabrera G, Amirthalingam G, Andrews N et al (2014). A case-control study to estimate the effectiveness of maternal pertussis vaccination in protecting newborn infants in England and Wales, 2012–2013. *Clin Infect Dis*.
6. Amirthalingam G, Campbell H, Ribeiro S, Fry NK, Ramsay M, Miller E, Andrews N (2016). Sustained effectiveness of the maternal pertussis immunization program in England 3 Years following introduction. *Clin Infect Dis*.
7. Donegan K, King B, Bryan P (2014). Safety of pertussis vaccination in pregnant women in UK: observational study. *BMJ* 349:g4219.
8. PHE website: Guidelines for the public health management of pertussis: <https://www.gov.uk/government/publications/pertussis-guidelines-for-public-health-management>
9. *HPR* 12(15), 27 April 2018.
10. *HPR* 11(34), 29 September 2017.

About Public Health England

Public Health England exists to protect and improve the nation's health and wellbeing, and reduce health inequalities. We do this through world-class science, knowledge and intelligence, advocacy, partnerships and the delivery of specialist public health services. We are an executive agency of the Department of Health, and are a distinct delivery organisation with operational autonomy to advise and support government, local authorities and the NHS in a professionally independent manner.

About Health Protection Report

Health Protection Report is a national public health bulletin for England and Wales, published by Public Health England. It is PHE's principal channel for the dissemination of laboratory data relating to pathogens and infections/communicable diseases of public health significance and of reports on outbreaks, incidents and ongoing investigations.

Public Health England, Wellington House, 133-155 Waterloo Road, London SE1 8UG

Tel: 020 7654 8000 www.gov.uk/phe

Twitter: [@PHE_uk](https://twitter.com/PHE_uk) Facebook: www.facebook.com/PublicHealthEngland

Queries relating to this document should be directed to:

Immunisation, Hepatitis and Blood Safety Department,

National Infection Service, PHE Colindale,

61 Colindale Avenue, London NW9 5EQ.

immunisation@phe.gov.uk

© Crown copyright 2018

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](https://www.ogcl.gov.uk) or email psi@nationalarchives.gsi.gov.uk. Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

Published June 2018

PHE publications gateway number: 2018151

