

GP In Hours

Data to: 15 July 2018

Syndromic Surveillance System: England

7 July 2018

Year: 2018 Week: 28

In This Issue:

Key messages.

Diagnostic indicators at a glance.

GP practices and denominator population.

National syndromic indicators.

Notes and further information.

Appendix.

Key messages

During week 28, consultations for gastrointestinal syndromes have shown small increases, mainly in the under 1 year age group (Figures 7&7a, 8&8a, 9&9a).

GP consultations for heat/sun stroke remain elevated but within levels expected for the recent hot weather (figure 21).

A Heat-Health Watch system operates in England from 1 June to 15 September each year. As part of the Heatwave Plan for England, the PHE Real-time Syndromic Surveillance team will be routinely monitoring the public health impact of hot weather using syndromic surveillance data during this period.

Heat-health watch level (current reporting week): Level 1/2 Summer preparedness/ Alert & readiness/ http://www.metoffice.gov.uk/weather/uk/heathealth/

Diagnostic indicators at a glance:

Indicator	Trend	Level
Upper respiratory tract infection	decreasing	below baseline levels
Influenza-like illness	no trend	similar to baseline levels
Pharyngitis	decreasing	below baseline levels
Scarlet fever	decreasing	below baseline levels
Lower respiratory tract infection	decreasing	below baseline levels
Pneumonia	decreasing	below baseline levels
Gastroenteritis	increasing	similar to baseline levels
Vomiting	no trend	similar to baseline levels
Diarrhoea	increasing	similar to baseline levels
Asthma	decreasing	similar to baseline levels
Conjunctivitis	no trend	below baseline levels
Mumps	no trend	below baseline levels
Measles	no trend	above baseline levels
Rubella	no trend	above baseline levels
Pertussis	no trend	below baseline levels
Chickenpox	no trend	below baseline levels
Herpes zoster	no trend	below baseline levels
Cellulitis	no trend	similar to baseline levels
Impetigo	no trend	below baseline levels
Allergic rhinitis	decreasing	below baseline levels
Heat/sunstroke	decreasing	above baseline levels
Insect Bites	no trend	above baseline levels

GP practices and denominator population:


Year	Week	GP Practices Reporting**	Population size**
2018	28	2,293	19.3 million

^{**}based on the average number of practices and denominator population in the reporting working week.




1: Upper respiratory tract infection (URTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




2: Influenza-like illness (ILI)

Daily incidence rates (and 7-day moving average*) per 100,000 population (all England, all ages).



3: Pharyngitis or scarlet fever




^{* 7-}day moving average adjusted for bank holidays.

GP In Hours

7 July 2018 Year: 2018 Week: 28


4: Scarlet fever

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients)



4a: Scarlet fever by age

Average daily incidence rate by week per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients).




Intentionally left blank

^{* 7-}day moving average adjusted for bank holidays.


5: Lower respiratory tract infection (LRTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



6: Pneumonia

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




Intentionally left blank

^{* 7-}day moving average adjusted for bank holidays.




7: Gastroenteritis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




7a: Gastroenteritis by age

Average daily incidence rate by week per 100,000 population (all England).



8: Vomiting




^{* 7-}day moving average adjusted for bank holidays.




8a: Vomiting by age

Average daily incidence rate by week per 100,000 population (all England).




9: Diarrhoea

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



9a. Diarrhoea by age

Average daily incidence rate by week per 100,000 population (all England).




* 7-day moving average adjusted for bank holidays.




10: Asthma

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




10a: Asthma by age

Average daily incidence rate by week per 100,000 population (all England).



11: Conjunctivitis




^{* 7-}day moving average adjusted for bank holidays.



12: Mumps

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




13: Measles

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



14: Rubella

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




* 7-day moving average adjusted for bank holidays.




15: Pertussis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




16: Chickenpox

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




17: Herpes zoster






18: Cellulitis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




19: Impetigo

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




20: Allergic rhinitis






20a: Allergic rhinitis by age

Weekly incidence rate by week per 100,000 population (all England since April 2012).




21: Heat/sunstroke

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



22: Insect bites

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



* 7-day moving average adjusted for bank holidays.



Notes and further information

- The Public Health England GP in hours surveillance system is a syndromic surveillance system monitoring community-based morbidity recorded by GP practices.
- GP consultation data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team.
- This system captures anonymised GP morbidity data from two GP clinical software systems, EMIS, from version 1 of the QSurveillance® database, and TPP SystmOne.
- Baselines represent seasonally expected levels of activity and are constructed from
 historical data. Furthermore, they take into account any known substantial changes in data
 collection, population coverage or reporting practices. Baselines are refreshed using the
 latest data on a regular basis.

Maps:

- From week 40 2017 the levels of influenza-like illness (ILI) rates are illustrated in the bulletin appendix maps. The ILI intensity levels are calculated using the "Moving Epidemic Method" (MEM).¹ MEM is used as a standard methodology for setting influenza surveillance thresholds across Europe.²
- The current ILI thresholds are based upon previous influenza seasons from 2012/13
 onwards and therefore illustrate activity levels in relation to previous ILI activity recorded in
 the GPIH system. IILI thresholds presented in the maps should be interpreted with caution
 and reference made to other GP surveillance systems incorporating more historical
 data, which are available in the PHE National Influenza Report.
- The ILI thresholds have been calculated separately for each of the nine PHE Centres to allow for differences between areas e.g. background ILI rates are historically higher in London than other areas of England. However, upper tier Local Authority (utLA) ILI consultation rates are compared to Centre-level thresholds only and therefore utLAs with higher background rates than the Centre may appear to have higher ILI activity.
- ILI consultation rates presented for each utLA in the maps should be interpreted in context
 of regional and national ILI activity. The small numbers reported at this local level can often
 result in short-lived fluctuations in rates causing threshold exceedances that are out of
 context with national and regional activity. utLA ILI data should therefore be interpreted
 with caution and interpreted in context with the national influenza report which can
 be found here:

https://www.gov.uk/government/statistics/weekly-national-flu-reports

- The maps on the following pages contains Ordnance Survey data © Crown copyright and database right 2015. Contains National Statistics data © Crown copyright and database right 2015.
 - ¹ Vega T et al. Influenza Other Respir Viruses. 2013;**7**(4):546-58.

Acknowledgements:

We thank and acknowledge the University of Nottingham, ClinRisk® and the contribution of EMIS and EMIS practices. Data source: version 1 of the QSurveillance® database.

We thank TPP, ResearchOne and the SystmOne GP practices contributing to this surveillance system.

Contact ReSST:

syndromic.surveillance @phe.gov.uk

GP In Hours Syndromic Surveillance System Bulletin.

Produced by: PHE Real-time Syndromic Surveillance Team

1st Floor, 5 St Philip's Place, Birmingham, B3 2PW

Web: https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses

² Green HK et al. *Epidemiol Infect.* 2015;**143**(1):1-12.