



TB outcome at 12 months after treatment start for drug sensitive cases with expected treatment duration < 12 months*, 2012

Year	Completed		Died		Lost to follow-up		Still on treatment		Stopped		Not evaluated**		Total
	n	%	n	%	n	%	n	%	n	%	n	%	n
2003	4,561	69.4	404	6.1	285	4.3	253	3.9	64	1.0	1,008	15.3	6,575
2004	4,876	70.3	390	5.6	337	4.9	315	4.5	80	1.2	940	13.6	6,938
2005	5,287	70.2	431	5.7	371	4.9	416	5.5	88	1.2	937	12.4	7,530
2006	5,622	74.9	409	5.5	401	5.4	483	6.4	91	1.2	496	6.6	7,502
2007	5,724	77.3	409	5.5	335	4.5	510	6.9	83	1.1	344	4.7	7,405
2008	6,019	79.1	397	5.2	344	4.5	468	6.2	74	1.0	309	4.1	7,611
2009	6,420	81.0	382	4.8	328	4.1	474	6.0	88	1.1	236	3.0	7,928
2010	6,133	82.1	366	4.9	317	4.2	428	5.7	67	0.9	158	2.1	7,469
2011	6,459	81.4	361	4.6	388	4.9	505	6.4	77	1.0	143	1.8	7,933
2012	6,438	82.8	345	4.4	307	4.0	446	5.7	79	1.0	159	2.1	7,774

* Excludes initial and amplified to rifampicin resistant TB and MDR-TB cases and MDR-TB treated cases and those with CNS, spinal, miliary or cryptic disseminated TB

** Not evaluated includes missing, unknown and transferred out