



07 November 2018

Year: 2018 Week: 44

In This Issue:

- Key messages.
- Diagnostic indicators at a glance.
- GP practices and denominator population.
- National syndromic indicators.
- Winter 2018/19 appendix
- Notes and further information.

Key messages

Data to: 04 November 2018

Nothing new to report in week 44.

A Cold Watch System operates in England from 1 November to 31 March each year. As part of the Public Health England Cold Weather Plan for England the PHE Real-time Syndromic Surveillance team will be monitoring the impact of cold weather on syndromic surveillance data during this period.

Cold weather alert level (current reporting week): Level 1 **Winter preparedness**

<http://www.metoffice.gov.uk/weather/uk/coldweatheralert/>

Diagnostic indicators at a glance:

| Indicator | Trend | Level |
|-----------------------------------|------------|----------------------------|
| Upper respiratory tract infection | no trend | below baseline levels |
| Influenza-like illness | no trend | pre-epidemic threshold* |
| Pharyngitis | no trend | below baseline levels |
| Scarlet fever | no trend | below baseline levels |
| Lower respiratory tract infection | no trend | below baseline levels |
| Pneumonia | no trend | below baseline levels |
| Gastroenteritis | no trend | below baseline levels |
| Vomiting | no trend | below baseline levels |
| Diarrhoea | no trend | below baseline levels |
| Asthma | no trend | below baseline levels |
| Conjunctivitis | no trend | below baseline levels |
| Mumps | no trend | similar to baseline levels |
| Measles | no trend | similar to baseline levels |
| Rubella | no trend | similar to baseline levels |
| Pertussis | no trend | below baseline levels |
| Chickenpox | no trend | similar to baseline levels |
| Herpes zoster | no trend | below baseline levels |
| Cellulitis | no trend | below baseline levels |
| Impetigo | increasing | similar to baseline levels |

* Moving Epidemic Method (MEM) influenza activity threshold (see notes)

GP practices and denominator population:

| Year | Week | GP Practices Reporting** | Population size** |
|------|------|--------------------------|-------------------|
| 2018 | 44 | 3,578 | 30.6 million |


**based on the average number of practices and denominator population in the reporting working week.

07 November 2018

Year: 2018 Week: 44


1: Upper respiratory tract infection (URTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



1a: Upper respiratory tract infection (URTI) by age

Average daily incidence rate by week per 100,000 population (all England).



Intentionally left blank


* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


2: Influenza-like illness (ILI)

Daily incidence rates (and 7-day moving average*) per 100,000 population (all England, all ages).




2a: Influenza-like illness by age

Average daily incidence rate by week per 100,000 population (all England).



2b: Influenza-like illness by PHE Centre

Average daily incidence rate by week per 100,000 population (all ages).




* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


2c: Daily influenza-like illness (winter 2018/19) with MEM influenza activity thresholds (see notes)

Daily incidence rates (and 7-day moving average*) per 100,000 population (all England, all ages).



2d: Daily influenza-like illness by week with MEM influenza activity thresholds and comparison to previous seasons (see notes)

Average daily incidence rates* per 100,000 population (all England, all ages).



Intentionally left blank


* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


3: Pharyngitis or scarlet fever

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




4: Scarlet fever

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients)



4a: Scarlet fever by age

Average daily incidence rate by week per 100,000 population (all England, based on a denominator population of approximately 5.5 million patients).



* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


5: Lower respiratory tract infection (LRTI)

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



6: Pneumonia

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



Intentionally left blank


* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


7: Gastroenteritis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




7a: Gastroenteritis by age

Average daily incidence rate by week per 100,000 population (all England).



8: Vomiting

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



* 7-day moving average adjusted for bank holidays.




07 November 2018


Year: 2018 Week: 44

8a: Vomiting by age


Average daily incidence rate by week per 100,000 population (all England).

**9: Diarrhoea**

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).

**9a. Diarrhoea by age**

Average daily incidence rate by week per 100,000 population (all England).




* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


10: Asthma

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




10a: Asthma by age

Average daily incidence rate by week per 100,000 population (all England).



11: Conjunctivitis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


12: Mumps

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




13: Measles

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



14: Rubella

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44


15: Pertussis

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




16: Chickenpox

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).



17: Herpes zoster

Daily incidence rate (and 7-day moving average*) per 100,000 population (all England, all ages).




* 7-day moving average

07 November 2018

Year: 2018 Week: 44


18: Cellulitis

Daily incidence rate
(and 7-day moving
average*) per 100,000
population (all England,
all ages).



19: Impetigo

Daily incidence rate
(and 7-day moving
average*) per 100,000
population (all England,
all ages).



Intentionally left blank

* 7-day moving average
adjusted for bank
holidays.



07 November 2018

Year: 2018 Week: 44

Winter 2018/19 appendix

20. Influenza-like illness maps (England) illustrating ILI activity across each of the nine PHE Centres

Week 41




Week 42



Week 43



Week 44



* 7-day moving average adjusted for bank holidays.

07 November 2018
Year: 2018 Week: 44

21. Influenza-like illness by age (England) MEM heat map

Average daily incidence rate by week per 100,000 population (all England).

| PHE Centre | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 1 | 2 | 3 | 4 |
|----------------------|------|------|------|------|------|----|----|----|----|----|----|----|----|---|---|---|---|
| North West | 0.75 | 1.03 | 0.77 | 0.83 | 0.87 | | | | | | | | | | | | |
| Yorkshire and Humber | 0.55 | 0.40 | 0.35 | 0.32 | 0.42 | | | | | | | | | | | | |
| North East | 0.62 | 0.60 | 0.42 | 0.51 | 0.42 | | | | | | | | | | | | |
| West Midlands | 0.71 | 0.79 | 0.69 | 0.65 | 0.74 | | | | | | | | | | | | |
| East Midlands | 0.26 | 0.47 | 0.42 | 0.38 | 0.43 | | | | | | | | | | | | |
| East of England | 0.46 | 0.43 | 0.37 | 0.35 | 0.34 | | | | | | | | | | | | |
| South West | 0.55 | 0.80 | 0.82 | 0.61 | 0.82 | | | | | | | | | | | | |
| London | 1.05 | 0.84 | 0.94 | 0.90 | 1.12 | | | | | | | | | | | | |
| South East | 0.66 | 0.71 | 0.77 | 0.57 | 0.68 | | | | | | | | | | | | |

| PHE Centre | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
|----------------------|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|
| North West | | | | | | | | | | | | | | | | |
| Yorkshire and Humber | | | | | | | | | | | | | | | | |
| North East | | | | | | | | | | | | | | | | |
| West Midlands | | | | | | | | | | | | | | | | |
| East Midlands | | | | | | | | | | | | | | | | |
| East of England | | | | | | | | | | | | | | | | |
| South West | | | | | | | | | | | | | | | | |
| London | | | | | | | | | | | | | | | | |
| South East | | | | | | | | | | | | | | | | |

22. Influenza-like illness PHE Centre (all age) MEM heat map

Average daily incidence rate by week per 100,000 population (all England).

| Age band (years) | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 1 | 2 | 3 | 4 |
|------------------|------|------|------|------|------|----|----|----|----|----|----|----|----|---|---|---|---|
| Under 1 | 1.00 | 0.46 | 0.80 | 0.76 | 0.56 | | | | | | | | | | | | |
| 1 to 4 | 0.48 | 0.46 | 0.46 | 0.58 | 0.62 | | | | | | | | | | | | |
| 5 to 14 | 0.28 | 0.24 | 0.18 | 0.14 | 0.26 | | | | | | | | | | | | |
| 15 to 44 | 0.82 | 0.88 | 0.76 | 0.72 | 0.86 | | | | | | | | | | | | |
| 45 to 64 | 0.80 | 0.92 | 0.82 | 0.76 | 0.84 | | | | | | | | | | | | |
| 65 to 74 | 0.46 | 0.52 | 0.64 | 0.56 | 0.64 | | | | | | | | | | | | |
| 75 and over | 0.46 | 0.46 | 0.58 | 0.40 | 0.54 | | | | | | | | | | | | |

| Age band (years) | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
|------------------|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|
| Under 1 | | | | | | | | | | | | | | | | |
| 1 to 4 | | | | | | | | | | | | | | | | |
| 5 to 14 | | | | | | | | | | | | | | | | |
| 15 to 44 | | | | | | | | | | | | | | | | |
| 45 to 64 | | | | | | | | | | | | | | | | |
| 65 to 74 | | | | | | | | | | | | | | | | |
| 75 and over | | | | | | | | | | | | | | | | |

23. GPIH ILI MEM threshold values

| PHE Centre | Pre-epidemic | Medium | High | Very high | Age band (years) | Pre-epidemic | Medium | High | Very high |
|----------------------|--------------|--------|------|-----------|------------------|--------------|--------|------|-----------|
| North West | 1.87 | 3.02 | 7.51 | 11.24 | Under 1 | 1.28 | 1.49 | 2.03 | 2.33 |
| Yorkshire and Humber | 0.94 | 2.09 | 4.55 | 6.41 | 1 to 4 | 1.49 | 1.87 | 2.88 | 3.48 |
| North East | 1.28 | 2.37 | 6.23 | 9.56 | 5 to 14 | 0.94 | 1.77 | 4.07 | 5.87 |
| West Midlands | 1.85 | 3.00 | 6.87 | 9.91 | 15 to 44 | 1.82 | 3.4 | 6.28 | 8.24 |
| East Midlands | 1.08 | 2.00 | 4.23 | 5.90 | 45 to 64 | 1.87 | 3.71 | 8.44 | 12.14 |
| East of England | 1.11 | 2.11 | 3.87 | 5.05 | 65 to 74 | 1.16 | 2.35 | 5.86 | 8.78 |
| South West | 1.84 | 3.41 | 8.11 | 11.90 | 75 and over | 1.01 | 1.97 | 5.96 | 9.71 |
| London | 2.29 | 3.82 | 5.87 | 7.11 | | | | | |
| South East | 1.73 | 3.36 | 7.25 | 10.19 | | | | | |

* 7-day moving average adjusted for bank holidays.

07 November 2018

Year: 2018 Week: 44

Notes and further information

- The Public Health England GP in hours surveillance system is a syndromic surveillance system monitoring community-based morbidity recorded by GP practices.
- GP consultation data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team.
- This system captures anonymised GP morbidity data from two GP clinical software systems, EMIS, from version 1 of the QSurveillance® database, and TPP SystmOne.
- Baselines represent seasonally expected levels of activity and are constructed from historical data since April 2012. They take into account any known substantial changes in data collection, population coverage or reporting practices. Gastroenteritis, diarrhoea and vomiting baselines also account for changes since the introduction of rotavirus vaccine in July 2013. Baselines are refreshed using the latest data on a regular basis.

Moving Epidemic Method:

- During winter 2018/19 we are presenting Moving Epidemic Method (MEM) influenza thresholds on selected indicators.
- The moving epidemic method or MEM is a standard methodology used for setting influenza thresholds across many European nations.¹
- MEM is used for GP ILI thresholds at a national level and at PHE Centre level and stratified by age band.
- MEM thresholds should be interpreted using 7 day moving averages rather than daily data.**
- MEM thresholds currently use five years of historic data (2013-2018). The thresholds are re-calculated every year.
- 'Pre-epidemic thresholds' are used alongside other surveillance systems to identify the start of influenza circulating in the community; 40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high.

¹Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.

Maps:

- From week 40 2018 the levels of influenza-like illness (ILI) rates are illustrated in the bulletin appendix maps. The ILI intensity levels are calculated using MEM.
- The current ILI thresholds are based upon previous influenza seasons from 2012/13 onwards and therefore illustrate activity levels in relation to previous ILI activity recorded in the GPIH system. **ILI thresholds presented in the maps should be interpreted with caution and reference made to other GP surveillance systems incorporating more historical data, which are available in the PHE National Influenza Report.**

<https://www.gov.uk/government/statistics/weekly-national-flu-reports>

- The ILI thresholds have been calculated separately for each of the nine PHE Centres to allow for differences between areas e.g. background ILI rates are historically higher in London than other areas of England.

Acknowledgements:

We thank and acknowledge the University of Nottingham, ClinRisk® and the contribution of EMIS and EMIS practices. Data source: version 1 of the QSurveillance® database.

We thank TPP, ResearchOne and the SystmOne GP practices contributing to this surveillance system.

Contact ReSST:
syndromic.surveillance
@phe.gov.uk

GP In Hours Syndromic Surveillance System Bulletin.

Produced by: PHE Real-time Syndromic Surveillance Team
1st Floor, 5 St Philips Place, Birmingham, B3 2PW

Tel: 0344 225 3560 > Option 4 > Option 2 **Fax:** 0121 236 2215

Web: <https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses>