



Treating your infection



Patient Name Doctor or Nurse Practitioner recommends that you self-care

Your infection	Usually lasts	How to treat yourself better for these infections, now and next time	When should you get help: Contact your GP practice or contact NHS 111 (England), NHS 24 (Scotland dial 111), or NHS Direct (Wales dial 0845 4647)
<input type="checkbox"/> Middle-ear infection	4 days	<ul style="list-style-type: none"> Have plenty of rest. Drink enough fluids to avoid feeling thirsty. Ask your local pharmacist to recommend medicines to help your symptoms or pain (or both). Fever is a sign the body is fighting the infection and usually gets better by itself in most cases. You can use paracetamol if you or your child are uncomfortable as a result of a fever. Use a tissue and wash your hands well to help prevent spread of your infection to your family, friends and others you meet. Other things you can do suggested by GP or nurse: 	<p>1. to 8. are possible signs of serious illness and should be assessed urgently. Phone for advice if you are not sure how urgent the symptoms are.</p> <ol style="list-style-type: none"> If you develop a severe headache and are sick. If your skin is very cold or has a strange colour, or you develop an unusual rash. If you feel unwell or have slurred speech or are very drowsy. If you have difficulty breathing. Signs can include: <ul style="list-style-type: none"> breathing quickly turning blue around the lips and the skin below the mouth skin between or above the ribs getting sucked or pulled in with every breath. If you develop chest pain. If you have difficulty swallowing or are drooling. If you cough up blood. If you are feeling a lot worse. <p>Less serious signs that can usually wait until the next available GP appointment:</p> <ol style="list-style-type: none"> If you are not starting to improve a little by the time given in the 'Usually lasts' column. In children with middle-ear infection: if fluid is coming out of their ears for more than 10 days or if they have new deafness. Other
<input type="checkbox"/> Sore throat	7 days		
<input type="checkbox"/> Common cold	10 days		
<input type="checkbox"/> Sinusitis	18 days		
<input type="checkbox"/> Cough or bronchitis	3 weeks		
<input type="checkbox"/> Other infection: days		

Antibiotics may be required after days only if you are not starting to feel a little better or you feel worse. Contact your GP practice if this is the case.

- Colds, most coughs, sinusitis, ear infections, sore throats, and other infections often get better without antibiotics, as your body can usually fight these infections on its own.
- The more we use antibiotics, the greater the chance that bacteria will become resistant to them so that they no longer work on our infections.
- Antibiotics can cause side effects such as rashes, thrush, stomach pains, diarrhoea, reactions to sunlight, other symptoms, or being sick if you drink alcohol with metronidazole.

Never share antibiotics and always return any unused antibiotics to a pharmacy for safe disposal

