

PHE Influenza Surveillance graphs

- Influenza activity in the United Kingdom is monitored using a variety of <u>surveillance systems</u>. The <u>National Influenza Report</u> which interprets the data is published weekly.
- National and subnational outputs from the various surveillance systems are grouped according to the following categories:
 - Community influenza surveillance
 - <u>Influenza-like illness GP consultation rates</u>
 - Influenza-confirmed hospitalisations
 - All-cause mortality
 - Microbiological surveillance
 - Influenza vaccination


Influenza Section Immunisations and Countermeasures Division Centre for Infectious Disease Surveillance and Control Public Health England respscidsc@phe.gov.uk





Community influenza surveillance

10 January 2019





Number of acute respiratory outbreaks by PHE Centre

	Outbreaks reported in week 52	Cumulative number of outbreaks since week 40
PHE North East	2	9
PHE North West	12	71
PHE Yorkshire and Humber	6	24
PHE East Midlands	8	40
PHE West Midlands	14	43
PHE East of England	18	80
PHE London	4	17
PHE South East	18	46
PHE South West	10	56




Influenza-like illness GP consultation rates

10 January 2019



RCGP ILI consultation rate, England



Return to main menu



GP ILI consultations in the UK for all ages with MEM thresholds applied

GP ILI consultation	Week number																
rates (all ages)	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4
England (RCGP)	4.2	3.9	4.5	3.6	3.6	5.3	5.2	6.4	6.2	7.6	8.1	9.4	8.4	14.8			
Wales	7.0	3.6	4.2	6.6	6.3	6.4	4.5	4.7	6.5	3.2	4.5	9.0	9.5	14.7			
Scotland	7.1	5.1	3.6	4.5	2.9	7.3	4.2	5.2	5.9	4.1	6.8	10.5	7.1	17.0			
Northern Ireland	3.8	3.5	3.8	3.6	3.8	5.0	6.3	4.5	5.6	6.0	8.4	8.9	9.0	13.5			

*The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for GP ILI consultations for the start of influenza activity (based on 10 seasons excluding 2009/10), in a standardised approach across Europe. For MEM threshold values for each country, please visit: <u>https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#clinical-surveillance-through-primary-care</u>.



Influenza-confirmed hospitalisations

10 January 2019


Public Health England trust catchment pc

Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population , England, since week 40 2018

Number of cases


----Rate of hospital admission/100,000 (2018/19)

--- Rate of hospital admission/100,000 (2017/18)




*The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for ICU/HDU admission rates for the start of influenza activity (based on 6 seasons) in a standardised approach across Europe. For MEM threshold values, please visit: https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#disease-severity-and-mortality-data

Return to main menu






Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population by PHE Centre, East Midlands, West Midlands and East of England since week 40 2018






Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population by PHE Centre, South East, South West and London, since week 40 2018






Influenza confirmed ICU admissions (USISS mandatory) by age and influenza subtype, since week 40 2018







Weekly hospitalised influenza case rate per 100,000 trust catchment population , England, since week 40 2018



*The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for ICU/HDU admission rates for the start of influenza activity (based on 6 seasons) in a standardised approach across Europe. For MEM threshold values, please visit: <u>https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#disease-severity-and-mortality-data</u>

Return to main menu






NB: due to technical issues the PHE region breakdown is not available for week 01

Return to main menu

Public Health England by age and influenza subtype, since week 40 2018








All-cause mortality




Excess mortality (all causes) by week of death through the EuroMOMO algorithm in 65+ year olds, England








Excess mortality (all causes) by week of death through the Public Health England
EuroMOMO algorithm in all ages, PHE Midlands and East of England






Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE London






Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE South of England






Microbiological surveillance


10 January 2019










DataMart: Number and proportion of samples positive for influenza, by type, PHE North West of England









Week number (of sample)











DataMart: Number and proportion of samples positive for influenza, by type, PHE East of England
























Influenza vaccination







Links

• Sources of UK flu surveillance data:

<u>https://www.gov.uk/sources-of-uk-flu-data-</u> influenza-surveillance-in-the-uk

• Weekly flu report

https://www.gov.uk/government/statistics/wee kly-national-flu-reports-2018-to-2019-season