

PHE Influenza Surveillance graphs

- Influenza activity in the United Kingdom is monitored using a variety of <u>surveillance systems</u>. The <u>National Influenza Report</u> which interprets the data is published weekly.
- National and subnational outputs from the various surveillance systems are grouped according to the following categories:
 - Community influenza surveillance
 - Influenza-like illness GP consultation rates
 - Influenza-confirmed hospitalisations
 - All-cause mortality
 - Microbiological surveillance
 - Influenza vaccination


Influenza Section
Immunisations and Countermeasures Division
Centre for Infectious Disease Surveillance and Control
Public Health England
respscidsc@phe.gov.uk



Community influenza surveillance



Number of acute respiratory outbreaks by institution, UK





Number of acute respiratory outbreaks by PHE Centre


| | Outbreaks reported in week 11 | Cumulative number of outbreaks since week 40 |
|--------------------------|-------------------------------|--|
| PHE North East | 4 | 36 |
| PHE North West | 3 | 167 |
| PHE Yorkshire and Humber | 4 | 84 |
| PHE East Midlands | 2 | 109 |
| PHE West Midlands | 2 | 118 |
| PHE East of England | 15 | 178 |
| PHE London | 0 | 56 |
| PHE South East | 2 | 163 |
| PHE South West | 6 | 172 |



Influenza-like illness GP consultation rates



RCGP ILI consultation rate, England





GP ILI consultations in the UK for all ages with MEM thresholds applied

| GP ILI consultation | | Week number | | | | | | | | | | | | | | | | | | | | | | |
|---------------------|-----|-------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|-----|------|------|------|------|------|------|------|------|------|------|-----|
| rates (all ages) | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| England (RCGP) | 4.2 | 3.9 | 4.5 | 3.6 | 3.6 | 5.3 | 5.2 | 6.4 | 6.2 | 7.6 | 8.1 | 9.4 | 8.4 | 14.8 | 19.2 | 19.6 | 17.5 | 19.7 | 23.1 | 21.1 | 15.7 | 12.1 | 9.1 | 6.2 |
| Wales | 7.0 | 3.6 | 4.2 | 6.6 | 6.3 | 6.4 | 4.5 | 4.7 | 6.5 | 3.2 | 4.5 | 9.0 | 9.5 | 14.7 | 20.5 | 22.9 | 15.7 | 20.5 | 21.4 | 17.2 | 17.5 | 8.2 | 8.7 | 7.5 |
| Scotland | 7.1 | 5.8 | 4.0 | 3.8 | 2.8 | 7.6 | 4.0 | 4.7 | 5.6 | 4.0 | 6.5 | 10.1 | 6.9 | 17.7 | 26.7 | 18.0 | 28.4 | 32.7 | 32.3 | 27.2 | 20.8 | 10.2 | 10.2 | 6.3 |
| Northern Ireland | 3.8 | 3.5 | 3.8 | 3.6 | 3.8 | 5.0 | 6.3 | 4.5 | 5.6 | 6.0 | 8.4 | 8.9 | 9.0 | 13.5 | 18.9 | 14.4 | 12.4 | 14.5 | 16.2 | 14.5 | 11.1 | 9.0 | 5.9 | 5.6 |


^{*}The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for GP ILI consultations for the start of influenza activity (based on 10 seasons excluding 2009/10), in a standardised approach across Europe. For MEM threshold values for each country, please visit: https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#clinical-surveillance-through-primary-care.



Influenza-confirmed hospitalisations




Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population, England, since week 40 2018



^{*}The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for ICU/HDU admission rates for the start of influenza activity (based on 6 seasons) in a standardised approach across Europe. For MEM threshold values, please visit: https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#disease-severity-and-mortality-data




Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population by PHE Centre, North East, North West and Yorkshire and Humber, since week 40 2018






Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population by PHE Centre, East Midlands, West Midlands and East of England since week 40 2018






Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population by PHE Centre, South East, South West and London, since week 40 2018






Influenza confirmed ICU admissions (USISS mandatory) by age and influenza subtype, since week 40 2018






Weekly hospitalised influenza case rate per 100,000 trust catchment population, England, since week 40 2018



^{*}The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for ICU/HDU admission rates for the start of influenza activity (based on 6 seasons) in a standardised approach across Europe. For MEM threshold values, please visit: https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#disease-severity-and-mortality-data




Weekly hospitalised influenza admission rate per 100,000 trust catchment population by PHE region, since week 40 2018





Influenza confirmed hospital admissions (USISS sentinel) by age and influenza subtype, since week 40 2018






All-cause mortality

Return to main menu




Excess mortality (all causes) by week of death through the EuroMOMO algorithm in 65+ year olds, England






Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE North of England






Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE Midlands and East of England






Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE London





Excess mortality (all causes) by week of death through the EuroMOMO algorithm in all ages, PHE South of England






Microbiological surveillance




DataMart: Number and proportion of samples positive for influenza, by type, England






DataMart: Number and proportion of samples positive for influenza, by type, PHE North East of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE North West of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE Yorkshire and Humber






DataMart: Number and proportion of samples positive for influenza, by type, PHE East Midlands of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE West Midlands of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE East of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE London






DataMart: Number and proportion of samples positive for influenza, by type, PHE South East of England






DataMart: Number and proportion of samples positive for influenza, by type, PHE South West of England






DataMart: Proportion of samples positive for influenza, by age group





DataMart: Proportion of samples positive for other viruses






Influenza vaccination




Seasonal influenza vaccine uptake in targeted groups, 2018/19, England





Seasonal influenza vaccine uptake in targeted groups, 2018/19, England





Links

Sources of UK flu surveillance data:

https://www.gov.uk/sources-of-uk-flu-datainfluenza-surveillance-in-the-uk

Weekly flu report

https://www.gov.uk/government/statistics/weekly-national-flu-reports-2018-to-2019-season