

Protecting and improving the nation's health

## Food and Environmental Proficiency Testing Unit

The final date for return of results is indicated on the accompanying request/report form

#### **Contact details:**

The Organisers - FEPTU Public Health England 61 Colindale Avenue, London, NW9 5EQ, UK. Fax: +44 (0) 20 8200 8264 Tel: +44 (0) 20 8327 7119 e-mail: foodeqa@phe.gov.uk

Results to be returned via an online survey as per email sent to your laboratory

 $\label{eq:https://www.gov.uk/government/collections/external-quality-assessment-eqa-and-proficiency-testing-pt-for-food-water-and-environmental-microbiology#mycobacterium-spp.-in-water-scheme$ 

# Mycobacterium spp. in Water Scheme – Instruction Sheet

The LENTICULE® discs are provided in screw-cap plastic vials (with a desiccant). Each LENTICULE disc rests on the filter support containing the desiccant. The **desiccant** should be orange in colour. Please contact us if this is not the case. Each LENTICULE disc requires reconstitution by a process of rehydration and dispersion prior to examination, as described below.

Refer also to the Safety Data Sheet: <a href="http://www.gov.uk/government/publications/safety-data-sheet-for-lenticules">www.gov.uk/government/publications/safety-data-sheet-for-lenticules</a>

#### Storage:

- Store the samples at -20 ± 5°C on receipt.
- Allow the LENTICULE discs to reach ambient temperature (5 10 minutes) before reconstituting in diluent.

### Reconstitution:

- Open the sample container and transfer the LENTICULE disc into approximately 9mL 0.1% peptone saline (maximum recovery diluent (MRD)) that has been allowed to reach ambient temperature.
- Leave to stand at ambient temperature for 10 -12 minutes to rehydrate.
- Tighten the cap of the bottle and shake to disperse the micro-organisms.
- Transfer all the inoculated MRD to a sample bottle containing 1L sterile deionised or distilled water at ambient temperature.
- Rinse with approximately 2mL from the sample bottle, ensuring all liquid is transferred back to the 1L sample.
- Disperse the inoculum by inverting approximately 30 times.

#### **Examination:**

- Each reconstituted sample is equivalent to 1L water.
- Undertake the sample examinations between 30 45 minutes of reconstitution.
- Examine in accordance with your routine procedures.

#### Return of Results:

- Return your results on-line using the web-based reporting system.
- The distribution will be closed on the deadline date for return of results.
- Results cannot be accepted after the distribution has closed.