

# **Remote Health Advice**

Syndromic Surveillance System: England

#### 30 July 2019

In This Issue:

Key messages.

a glance.

Data summary.

Total calls. Cold/flu.

Fever.

Cough.

Sore throat.

Diarrhoea. Vomiting.

Eve problems.

Introduction to charts. Moving Epidemic Method (MEM). Notes and further information. Acknowledgements.

Syndromic indicators at

Indicators by syndrome.

Difficulty Breathing.

Year: 2019

Week: 30

## Key messages

Data to: 28 July 2019

During week 30 there was an increase in calls due to the impact of heat (figure 10) in line with the hot weather. Increases were also seen in fever and vomiting calls (particularly in children) and in insect bite calls (figures 3, 8, 8a & 11). These increases are similar in scale to those seen in previous periods of hot weather.

A Heat-Health Watch system operates in England from 1 June to 15 September each year. As part of the Heatwave Plan for England, the PHE Real-time Syndromic Surveillance team will be routinely monitoring the public health impact of hot weather using syndromic surveillance data during this period.

Heat-health watch level (current reporting week): Level 1/3 Summer preparedness / Heatwave action http://www.metoffice.gov.uk/weather/uk/heathealth/

# Syndromic indicators at a glance:

IndicatorTrendLevelCold/fludecreasingpre-epidemic threshold*Feverincreasingsimilar to baseline levelsCoughdecreasingsimilar to baseline levels
Fever increasing similar to baseline levels
Ŭ
Cough decreasing similar to baseline levels
Difficulty breathing increasing similar to baseline levels
Sore throat decreasing similar to baseline levels
Diarrhoea no trend below baseline levels
Vomiting increasing below baseline levels
Eye problems decreasing similar to baseline levels
Heat/sun impact increasing above baselines levels
Insect bites increasing above baselines levels

\* Moving Epidemic Method (MEM) influenza activity threshold (see notes)

# Data summary:


Year	Week	Total calls
2019	30	252,301

# With Public Health England

#### 30 July 2019

#### 1: Total calls.

The total number of syndromic calls recorded each day by NHS 111.




#### 2: Cold/flu

Daily 'cold/flu' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



Daily 'fever' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



weekend

bank holiday

fever

-

-- baseline

## **Remote Health Advice**

7 day average (adjusted for bank holidays)

#### 30 July 2019

#### 4: Cough


Daily 'cough' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

#### 5: Difficulty breathing

Daily 'difficulty breathing' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

# 5a: Difficulty breathing calls by age group

Difficulty breathing calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.





# **Remote Health Advice**

#### **X** Public Health England

#### 30 July 2019

#### 6: Sore throat

Daily 'sore throat' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

#### 7. Diarrhoea

Daily 'diarrhoea' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

#### 7a: Diarrhoea calls by age group

Daily 'diarrhoea' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

3

2

1

0 29/07/18

26/08/18

23/09/18

<1

21/10/18

1 to 4

18/11/18

- 5 to 14

16/12/18

13/01/19

-15 to 44

10/02/19

45 to 64

10/03/19

07/04/19

65 to 74


05/05/19

->74

02/06/19

- All ages

30/06/19



# **Remote Health Advice**

28/07/19

#### 鯋 Public Health England


#### 30 July 2019


#### 8: Vomiting calls

Daily 'vomiting' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

#### 8a: Vomiting calls by age group


Vomiting calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.





#### 9: Eye problems

Daily 'eye problems' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.




### **Remote Health Advice**

Public Health England

繱

2013.

# **Remote Health Advice**



# **Remote Health Advice**

30 July 2019	Year: 2019 Week: 30
Introduction to charts:	<ul> <li>Weekends and bank holidays are marked by vertical grey lines (bank holidays darker grey).</li> <li>A 7-day moving average (adjusted for bank holidays) is overlaid on the daily data reported in each chart, unless specified.</li> <li>Baselines represent seasonally expected levels of activity and are constructed from historical data since September 2013. They take into account any known substantial changes in data collection, population coverage or reporting practices. Baselines are refreshed using the latest data on a regular basis.</li> <li>NHS 111 call data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team.</li> </ul>
Moving Epidemic Method (MEM):	<ul> <li>During each winter we present Moving Epidemic Method (MEM) influenza thresholds on selected indicators.</li> <li>The moving epidemic method or MEM is a standard methodology used for setting influenza thresholds across many European nations.<sup>1</sup></li> <li>MEM is used for NHS 111 cold/flu thresholds at a national level.</li> <li>MEM thresholds should be interpreted using 7 day moving averages rather than daily data.</li> <li>MEM thresholds currently use five years of historic data. The thresholds are re-calculated every year.</li> <li>'Pre-epidemic thresholds' are used alongside other surveillance systems to identify the start of influenza circulating in the community.</li> <li>40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high.</li> <li>'Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.</li> </ul>
Notes and further information:	<ul> <li>Further information about NHS 111 can be found at: <u>https://www.nhs.uk/using-the-nhs/nhs-services/urgent-and-emergency-care/nhs-111/</u></li> <li>The Remote Health Advice Syndromic Surveillance bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance: <u>https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses</u></li> </ul>
Acknowledgements:	We are grateful to NHS 111 and to NHS Digital for their assistance and support in providing the anonymised call data that underpin the Remote Health Advice Syndromic Surveillance System.
<b>Contact ReSST:</b> syndromic.surveillance @phe.gov.uk	Produced by: PHE Real-time Syndromic Surveillance Team 1 <sup>st</sup> Floor, 5 St Philips Place, Birmingham, B3 2PW Tel: 0344 225 3560 > Option 4 > Option 2 Fax: 0121 236 2215 Web: <u>https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses</u> 7