

Syndromic Surveillance System: England

Data to: 06 October 2019

09 October 2019 Year: 2019 Week: 40

#### In This Issue:

Key messages.

Syndromic indicators at a glance.

Data summary.

Indicators by syndrome.

Total calls.

Cold/flu.

Fever.

Cough.

Difficulty Breathing.

Sore throat.

Diarrhoea.

Vomiting.

Eye problems.

Introduction to charts.

Moving Epidemic Method (MEM).

Notes and further information.

Acknowledgements.

## Key messages

During week 40 NHS 111 cough calls increased, following seasonally expected trends (Figure 4). Cough calls increased particularly in the 1-4 years age group (Figure 4a).

## Syndromic indicators at a glance:

Indicator	Trend	Level
Cold/flu	no trend	pre-epidemic threshold*
Fever	no trend	below baseline levels
Cough	increasing	above baselines levels
Difficulty breathing	no trend	similar to baseline levels
Sore throat	no trend	above baselines levels
Diarrhoea	no trend	below baseline levels
Vomiting	no trend	below baseline levels
Eye problems	increasing	above baseline levels

## **Data summary:**

Year	Week	Total calls
2019	40	250,149

<sup>\*</sup> Moving Epidemic Method (MEM) influenza activity threshold (see notes)

Public Health England 09 October 2019 2019 Week: 40 100000 1: Total calls. 90000 The total number of 80000 syndromic calls recorded each day by 70000 NHS 111. 60000 50000 40000 30000

### 2: Cold/flu

Daily 'cold/flu' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.

07/10/18

04/11/18

02/12/18

30/12/18

bank holiday

27/01/19

weekend

24/02/19

24/03/19

total calls

21/04/19

19/05/19

16/06/19


■7 day average (adjusted for bank holidays)

14/07/19

11/08/19


08/09/19

06/10/19



### 2b: Daily cold/flu calls (winter 2019/20) with MEM influenza activity thresholds (see notes)

Daily 'cold/flu' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.




09 October 2019 Year: 2019 Week: 40


2c: Daily cold/flu calls by week with MEM influenza activity thresholds and comparison to previous seasons (see notes)

Average daily calls as percentage of total calls by week (7 day moving average, All England, All ages).



Daily 'fever' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.






### Intentionally left blank

09 October 2019 Year: 2019 Week: 40

### 4: Cough

Daily 'cough' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.




# 4a: Cough calls by age group

Cough calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.



### 5: Difficulty breathing


Daily 'difficulty breathing' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



09 October 2019 Year: 2019 Week: 40


# 5a: Difficulty breathing calls by age group

Difficulty breathing calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.




#### 6: Sore throat

Daily 'sore throat' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



# 6a: Sore throat calls by age group


Sore throat calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.



09 October 2019 Year: 2019 Week: 40


#### 7. Diarrhoea

Daily 'diarrhoea' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.




### 8: Vomiting calls

Daily 'vomiting' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



### 9: Eye problems


Daily 'eye problems' calls as a percentage of total calls. Baselines are constructed from historical data since 2013.



09 October 2019 Year: 2019 Week: 40

# 9a: Eye problems calls by age group

'Eye problems calls as a percentage of total calls within each age group, shown as a 7 day moving average adjusted for bank holidays.



### Intentionally left blank

### Intentionally left blank

09 October 2019 Year: 2019 Week: 40

#### Introduction to charts:

- Weekends and bank holidays are marked by vertical grey lines (bank holidays darker grey).
- A 7-day moving average (adjusted for bank holidays) is overlaid on the daily data reported in each chart, unless specified.
- Baselines represent seasonally expected levels of activity and are constructed from historical data since September 2013. They take into account any known substantial changes in data collection, population coverage or reporting practices. Baselines are refreshed using the latest data on a regular basis.
- NHS 111 call data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team.

### Moving Epidemic Method (MEM):

- During each winter we present Moving Epidemic Method (MEM) influenza thresholds on selected indicators.
- The moving epidemic method or MEM is a standard methodology used for setting influenza thresholds across many European nations.<sup>1</sup>
- MEM is used for NHS 111 cold/flu thresholds at a national level.
- MEM thresholds should be interpreted using 7 day moving averages rather than daily data.
- MEM thresholds currently use six years of historic data (2013-2019). The thresholds are re-calculated every year.
- 'Pre-epidemic thresholds' are used alongside other surveillance systems to identify the start of influenza circulating in the community.
- 40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high.

<sup>1</sup>Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.

# Notes and further information:

- Further information about NHS 111 can be found at: <a href="https://www.nhs.uk/using-the-nhs/nhs-services/urgent-and-emergency-care/nhs-111/">https://www.nhs.uk/using-the-nhs/nhs-services/urgent-and-emergency-care/nhs-111/</a>
- The Remote Health Advice Syndromic Surveillance bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance: <a href="https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses">https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses</a>

#### **Acknowledgements:**

We are grateful to NHS 111 and to NHS Digital for their assistance and support in providing the anonymised call data that underpin the Remote Health Advice Syndromic Surveillance System.

Remote Health Advice Syndromic Surveillance System Bulletin.

**Produced by:** PHE Real-time Syndromic Surveillance Team 1<sup>st</sup> Floor, 5 St Philips Place, Birmingham, B3 2PW

Contact ReSST: syndromic.surveillance @phe.gov.uk