


Unconditional amendments to EP (UK) 2 322 153:

1A. A compound that inhibits hypoxia inducible factor (HIF) prolyl hydroxylase activity for use in treating or preventing functional iron deficiency associated with anemia in a subject, wherein the compound is a structural mimetic of 2-oxoglutarate.

2. The compound according to claim 1, wherein the compound is of Formula (I):



wherein

A is 1,2-arylidene, 1,3-arylidene, 1,4-arylidene; or (C₁-C₄)-alkylene, optionally substituted by one or two halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, (C₁-C₆)-fluoroalkoxy, (C₁-C₈)-fluoroalkenyloxy, (C₁-C₈)-fluoroalkynyloxy, -OCF₂Cl, -O-CF₂-CHFCl; (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, phenyl, benzyl, phenoxy, benzyloxy, anilino, N-methylanilino, phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl, N,N-di-(C₁-C₄)-alkylsulfamoyl; or by a substituted (C₆-C₁₂)-aryloxy, (C₇-C₁₁)-aralkyloxy, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl radical, which carries in the aryl moiety one to five identical or different substituents selected from halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)Hal_g, -OCF₂Cl, -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₆)-alkoxycarbonyl, carbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N,N-di-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-alkylcarbonyloxy, (C₃-C₈)-cycloalkyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl, N,N-di-(C₁-

C₄)-alkylsulfamoyl; or wherein A is -CR⁵R⁶ and R⁵ and R⁶ are each independently selected from hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, aryl, or a substituent of the α-carbon atom of an α-amino acid, wherein the amino acid is a natural L-amino acid or its D-isomer.

B is -CO₂H, -NH₂, -NHSO₂CF₃, tetrazolyl, imidazolyl, 3-hydroxyisoxazolyl, -CONHCOR^{'''}, -CONHSOR^{'''}, CONHSO₂R^{'''}, where R^{'''} is aryl, heteroaryl, (C₃-C₇)-cycloalkyl, or (C₁-C₄)-alkyl, optionally monosubstituted by (C₆-C₁₂)-aryl, heteroaryl, OH, SH, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-thioalkyl, (C₁-C₄)-sulfinyl, (C₁-C₄)-sulfonyl, CF₃, Cl, Br, F, I, NO₂, -COOH, (C₂-C₅)-alkoxycarbonyl, NH₂, mono-(C₁-C₄)-alkyl)-amino, di-(C₁-C₄)-alkyl)-amino, or (C₁-C₄)-perfluoroalkyl; or wherein B is a CO₂-G carboxyl radical, where G is a radical of an alcohol G-OH in which G is selected from (C₁-C₂₀)-alkyl radical, (C₃-C₈) cycloalkyl radical, (C₂-C₂₀)-alkenyl radical, (C₃-C₈)-cycloalkenyl radical, retinyl radical, (C₂-C₂₀)-alkynyl radical, (C₄-C₂₀)-alkenylnyl radical, where the alkenyl, cycloalkenyl, alkynyl, and alkenylnyl radicals contain one or more multiple bonds; (C₆-C₁₆)-carbocyclic aryl radical, (C₇-C₁₆)-carbocyclic aralkyl radical, heteroaryl radical, or heteroaralkyl radical, wherein a heteroaryl radical or heteroaryl moiety of a heteroaralkyl radical contains 5 or 6 ring atoms; and wherein radicals defined for G are substituted by one or more hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyl, (C₅-C₈)-cycloalkenyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₂-C₁₂)-alkenyl, (C₂-C₁₂)-alkynyl, (C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, (C₁-C₈)-hydroxyalkyl, -O-[CH₂]_x-C_fH_(2f+1-q)-F_q, -OCF₂Cl, -OCF₂-CHFCl, (C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl, (C₆-C₁₂)-arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl, cinnamoyl, (C₂-C₁₂)-alkenylcarbonyl, (C₂-C₁₂)-alkynylcarbonyl, (C₁-C₁₂)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₁₂)-alkenyloxycarbonyl, (C₂-C₁₂)-alkynyloxycarbonyl, acyloxy, (C₁-C₁₂)-

alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyloxy, (C₆-C₁₂)-
aryloxy-carbonyloxy, (C₇-C₁₆)-aralkyloxy-carbonyloxy, (C₃-C₈)-cycloalkoxy-carbonyloxy,
(C₂-C₁₂)-alkenyloxy-carbonyloxy, (C₂-C₁₂)-alkynyloxy-carbonyloxy, carbamoyl, N-(C₁-
C₁₂)-alkyl-carbamoyl, N,N-di(C₁-C₁₂)-alkyl-carbamoyl, N-(C₃-C₈)-cycloalkyl-carbamoyl,
N-(C₆-C₁₆)-aryl-carbamoyl, N-(C₇-C₁₆)-aralkyl-carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-
aryl-carbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkyl-carbamoyl, N-((C₁-C₁₀)-alkoxy-(C₁-
C₁₀)-alkyl)-carbamoyl, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-((C₇-C₁₆)-
aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-
alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-
(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, carbamoyloxy, N-(C₁-
C₁₂)-alkyl-carbamoyloxy, N,N-di-(C₁-C₁₂)-alkyl-carbamoyloxy, N-(C₃-C₈)-
cycloalkyl-carbamoyloxy, N-(C₆-C₁₂)-aryl-carbamoyloxy, N-(C₇-C₁₆)-
aralkyl-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-aryl-carbamoyloxy, N-(C₁-C₁₀)-alkyl-
N-(C₇-C₁₆)-aralkyl-carbamoyloxy, N-((C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₆-C₁₂)-
aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-
carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-
(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-
((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, amino, (C₁-C₁₂)-alkyl-amino, di-(C₁-
C₁₂)-alkyl-amino, (C₃-C₈)-cycloalkyl-amino, (C₂-C₁₂)-alkenyl-amino, (C₂-C₁₂)-
alkynyl-amino, N-(C₆-C₁₂)-aryl-amino, N-(C-C₁₁)-aralkyl-amino, N-alkyl-aralkyl-amino, N-
alkyl-aryl-amino, (C₁-C₁₂)-alkoxy-amino, (C₁-C₁₂)-alkoxy-N-(C₁-C₁₀)-alkyl-amino, (C₁-
C₁₂)-alkyl-carbonyl-amino, (C₃-C₈)-cycloalkyl-carbonyl-amino, (C₆-C₁₂)-
aryl-carbonyl-amino, (C₇-C₁₆)-aralkyl-carbonyl-amino, (C₁-C₁₂)-alkyl-carbonyl-N-(C₁-C₁₀)-
alkyl-amino, (C₃-C₈)-cycloalkyl-carbonyl-N-(C₁-C₁₀)-alkyl-amino, (C₆-C₁₂)-aryl-carbonyl-
N-(C₁-C₁₀)-alkyl-amino, (C₇-C₁₁)-aralkyl-carbonyl-N-(C₁-C₁₀)-alkyl-amino, (C₁-C₁₂)-
alkyl-carbonyl-amino-(C₁-C₈)-alkyl, (C₃-C₈)-cycloalkyl-carbonyl-amino-(C₁-C₈)-alkyl, (C₆-
C₁₂)-aryl-carbonyl-amino-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkyl-carbonyl-amino-(C₁-C₈)-alkyl,
amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkyl-amino-(C₁-C₁₀)-alkyl, N,N-di-(C₁-C₁₀)-

alkylamino-(C₁-C₁₀)-alkyl, (C₃-C₈)cycloalkylamino-(C₁-C₁₀)-alkyl, (C₁-C₁₂)-alkylmercapto, (C₁-C₁₂)-alkylsulfinyl, (C₁-C₁₂)-alkylsulfonyl, (C₆-C₁₆)-arylmercapto, (C₆-C₁₆)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl, (C₇-C₁₆)-aralkylsulfonyl, sulfamoyl, N-(C₁-C₁₀)-alkylsulfamoyl, N,N-di(C₁-C₁₀)-alkylsulfamoyl, (C₃-C₈)-cycloalkylsulfamoyl, N-(C₆-C₁₂)-alkylsulfamoyl, N-(C₇-C₁₆)-aralkylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylsulfamoyl, (C₁-C₁₀)-alkylsulfonamido, N-((C₁-C₁₀)-alkyl)-(C₁-C₁₀)-alkylsulfonamido, (C₇-C₁₆)-aralkylsulfonamido, or N-((C₁-C₁₀)-alkyl)-(C₇-C₁₆)-aralkylsulfonamido; wherein radicals which are aryl or contain an aryl moiety, may be substituted on the aryl by one to five identical or different hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyl, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, (C₁-C₈)-hydroxyalkyl, (C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkyl-carbonyl, (C₆-C₁₂)-arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl, (C₁-C₁₂)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₁₂)-alkenyloxycarbonyl, (C₂-C₁₂)-alkynyloxycarbonyl, (C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkylcarbonyloxy, (C₆-C₁₂)-arylcarbonyloxy, (C₇-C₁₆)-aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenylcarbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy, (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy, (C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxycarbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy, (C₂-C₁₂)-alkynyloxycarbonyloxy, carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-


(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N.N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyloxy, N(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, amino, (C₁-C₁₂)-alkylamino, di-(C₁-C₁₂)-alkylamino, (C₃-C₈)-cycloalkylamino, (C₃-C₁₂)-alkenylamino, (C₃-C₁₂)-alkynylamino, N-(C₆-C₁₂)-arylamino, N-(C₇-C₁₁)-aralkylamino, N-alkylaralkylamino, N-alkyl-arylamino, (C₁-C₁₂)-alkoxyamino, (C₁-C₁₂)-alkoxy-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkylcarbonylamino, (C₃-C₈)-cycloalkylcarbonylamino, (C₆-C₁₂)-arylcarbonylamino, (C₇-C₁₆)-alkylcarbonylamino, (C₁-C₁₂)-alkylcarbonyl-N-(C₁-C₁₀)-alkylamino, (C₃-C₈)-cycloalkylcarbonyl-N-(C₁-C₁₀)-alkylamino, (C₆-C₁₂)-arylcarbonyl-N-(C₁-C₁₀)-alkylamino, (C₇-C₁₁)-aralkylcarbonyl-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkylcarbonylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cycloalkylcarbonylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-arylcarbonylamino-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkylcarbonylamino-(C₁-C₈)-alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, N.N-di-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkylamino-(C₁-C₁₀)-alkyl, (C₁-C₁₂)-alkylmercapto, (C₁-C₁₂)-alkylsulfinyl, (C₁-C₁₂)-alkylsulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl, or (C₇-C₁₆)-aralkylsulfonyl;

X is O or S;

Q is O, S, NR', or a bond;


where, if Q is a bond, R⁴ is halogen, nitrile, or trifluoromethyl;

or where, if Q is O, S, or NR', R⁴ is hydrogen, (C₁-C₁₀)-alkyl radical, (C₂-C₁₀)-alkenyl radical, (C₂-C₁₀)-alkynyl radical, wherein alkenyl or alkynyl radical contains one or two C-C multiple bonds; unsubstituted fluoroalkyl radical of the formula -[CH₂]_x-C_fH_(2f+1-g)-F_g, (C₁-C₈)-alkoxy-(C₁-C₆)-alkyl radical, (C₁-C₆)-alkoxy-(C₁-C₄)-alkoxy-(C₁-C₄)-alkyl radical, aryl radical, heteroaryl radical, (C₇-C₁₁)-aralkyl radical, or a radical of the formula Z



where

E is a heteroaryl radical, a (C₃-C₈)-cycloalkyl radical, or a phenyl radical of the formula F



v is 0-6,

w is 0 or 1,

t is 0-3, and

R⁷, R⁸, R⁹, R¹⁰, and R¹¹ are identical or different and are hydrogen, halogen, cyano, nitro, trifluoromethyl, (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyl, (C₁-C₆)-alkoxy, -O-[CH₂]_x-C_fH_(2f+1-g)-F_g, -OCF₂-Cl, -O-CF₂-CHFCl, (C₁-C₆)-alkylmercapto, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy-(C₁-C₆)-alkoxy, (C₁-C₆)-alkoxy-(C₁-C₆)-alkyl, (C₁-C₆)-alkylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylcarbonyl, (C₁-C₈)-alkoxycarbonyl, carbamoyl, N-(C₁-C₈)-alkylcarbamoyl, N,N-di-(C₁-C₈)-alkylcarbamoyl, or (C₇-C₁₁)-aralkylcarbamoyl, optionally substituted by fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkoxy, N-(C₃-C₈)-cycloalkylcarbamoyl, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylcarbamoyl, (C₁-C₆)-

alkylcarbonyloxy, phenyl, benzyl, phenoxy, benzyloxy, NR^YR^Z wherein R^Y and R^Z are independently selected from hydrogen, (C₁-C₁₂)-alkyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₃-C₁₀)-cycloalkyl, (C₃-C₁₂)-alkenyl, (C₃-C₁₂)-alkynyl, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl, (C₁-C₁₂)-alkoxy, (C₇-C₁₂)aralkoxy, (C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl, (C₆-C₁₂)arylcarbonyl, (C₇-C₁₆)-aralkylcarbonyl; or further wherein R^Y and R^Z together are -[CH₂]_h, in which a CH₂ group can be replaced by O, S, N-(C₁-C₄)-alkylcarbonylimino, or N-(C₁-C₄)-alkoxycarbonylimino; phenylmercapto, phenylsulfonyl, phenylsulfinyl, sulfamoyl, N-(C₁-C₈)-alkylsulfamoyl, or N, N-di-(C₁-C₈)-alkylsulfamoyl; or alternatively R⁷ and R⁸, R⁸ and R⁹, R⁹ and R¹⁰, or R¹⁰ and R¹¹, together are a chain selected from -[CH₂]_n- or -CH=CH-CH=CH-, where a CH₂ group of the chain is optionally replaced by O, S, SO, SO₂, or NR^Y; and n is 3, 4, or 5; and if E is a heteroaryl radical, said radical can carry 1-3 substituents selected from those defined for R⁷-R¹¹, or if E is a cycloalkyl radical, the radical can carry one substituent selected from those defined for R⁷-R¹¹;


or where, if Q is NR', R⁴ is alternatively R'', where R' and R'' are identical or different and are hydrogen, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl, (C₁-C₈)-alkyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-alkylcarbonyl, optionally substituted (C₇-C₁₆)-aralkylcarbonyl, or optionally substituted (C₆-C₁₂)-arylcarbonyl; or R' and R'' together are -[CH₂]_h, in which a CH₂ group can be replaced by O, S, N-acylimino, or N-(C₁-C₁₀)-alkoxycarbonylimino, and h is 3 to 7.

Y is N or CR³;

R¹, R² and R³ are identical or different and are hydrogen, hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₁-C₂₀)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₈)cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkyloxy-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-

cycloalkyloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₇-C₁₆)-aralkenyl, (C₇-C₁₆)-aralkynyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₁-C₂₀)-alkoxy, (C₂-C₂₀)-alkenyloxy, (C₂-C₂₀)-alkynyloxy, retinyloxy, (C₁-C₂₀)-alkoxy-(C₁-C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy-(C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxy, (C₁-C₁₆)-hydroxyalkyl, (C₆-C₁₆)-aryloxy-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₂)-aralkyloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkenyloxy-(C₁-C₆)-alkyl, (C₂-C₂₀)-alkynyloxy-(C₁-C₆)-alkyl, retinyloxy-(C₁-C₆)-alkyl, -O-[CH₂]_xCfH_(2f+1-g)F_g, -OCF₂Cl, -OCF₂-CHFCl, (C₁-C₂₀)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl, (C₆-C₁₂)-arylcabonyl, (C₇-C₁₆)-aralkylcarbonyl, cinnamoyl, (C₂-C₂₀)-alkenylcarbonyl, (C₂-C₂₀)-alkynylcarbonyl, (C₁-C₂₀)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl, (C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₂-C₂₀)-alkenyloxycarbonyl, retinyloxycarbonyl, (C₂-C₂₀)-alkynyloxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₁-C₁₂)-alkylcarbonyloxy, (C₃-C₈)-cycloalkylcarbonyloxy, (C₆-C₁₂)-arylcabonyloxy, (C₇-C₁₆)-aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenylcarbonyloxy, (C₂-C₁₂)-alkynylcarbonyloxy, (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy, (C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-C₈)-cycloalkoxycarbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy, (C₂-C₁₂)-alkynyloxycarbonyloxy, carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di-(C₁-C₁₂)-alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)-carbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)-carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcabamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcabamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl,

N-((C₁-C₁₈)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl; CON(CH₂)_h, in which a CH₂ group can be replaced by O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino, N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino, and h is from 3 to 7; a carbamoyl radical of the formula R



in which

R^x and R^y are each independently selected from hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, aryl, or the substituent of an α-carbon of an α-amino acid, to which the L- and D-amino acids belong,

s is 1-5,

T is OH, or NR^{*}R^{**}, and R^{*}, R^{**} and R^{***} are identical or different and are selected from hydrogen, (C₆-C₁₂)-aryl, (C₇-C₁₁)-aralkyl, (C₁-C₈)-alkyl, (C₃-C₈)-cycloalkyl, (+)-dehydroabietyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-alkanoyl, optionally substituted (C₇-C₁₆)-aralkanoyl, optionally substituted (C₆-C₁₂)-aroyl; or R^{*} and R^{**} together are -[CH₂]_h, in which a CH₂ group can be replaced by O, S, SO, SO₂, N-acylamino, N-(C₁-C₁₀)-alkoxycarbonylimino, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino, N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino, and h is from 3 to 7;

carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₂)-arylcarmamoyloxy, N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarmamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)-carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyloxyamino, (C₁-C₁₂)-alkylamino, di-(C₁-C₁₂)-alkylamino, (C₃-C₈)-cycloalkylamino, (C₃-C₁₂)-alkenylamino, (C₃-C₁₂)-alkynylamino, N-(C₆-C₁₂)-arylamino, N-(C₇-C₁₁)-aralkylamino, N-alkyl-aralkylamino, N-alkyl-arylamino, (C₁-C₁₂)-alkoxyamino, (C₁-C₁₂)-alkoxy-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkanoylamino, (C₃-C₈)-cycloalkanoylamino, (C₆-C₁₂)-aroylamino, (C₇-C₁₆)-aralkanoylamino, (C₁-C₁₂)-alkanoyl-N-(C₁-C₁₀)-alkylamino, (C₃-C₈)-cycloalkanoyl-N-(C₁-C₁₀)-alkylamino, (C₆-C₁₂)-aroyl-N-(C₁-C₁₀)-alkylamino, (C₇-C₁₁)-aralkanoyl-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cycloalkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-aroylamino-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-(C₁-C₈)-alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, N,N-di(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkylamino(C₁-C₁₀)-alkyl, (C₁-C₂₀)-alkylmercapto, (C₁-C₂₀)-alkylsulfinyl, (C₁-C₂₀)-alkylsulfonyl, (C₆-C₁₂)-arylmercapto, (C₆-C₁₂)-arylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl, (C₇-C₁₆)-aralkylsulfonyl, (C₁-C₁₂)-alkylmercapto-(C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfinyl-(C₁-C₆)-alkyl, (C₁-C₁₂)-alkylsulfonyl-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylmercapto-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylsulfinyl-(C₁-C₆)-alkyl, (C₆-C₁₂)-arylsulfonyl-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylmercapto-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylsulfinyl-(C₁-C₆)-alkyl, (C₇-C₁₆)-aralkylsulfonyl-(C₁-C₆)-alkyl, sulfamoyl, N-(C₁-C₁₀)-alkylsulfamoyl, N,N-di-(C₁-C₁₀)-alkylsulfamoyl, (C₃-C₈)-cycloalkylsulfamoyl, N-(C₆-C₁₂)-arylsulfamoyl, N-(C₇-C₁₆)-aralkylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylsulfamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylsulfamoyl, (C₁-C₁₀)-alkylsulfonamido, N-((C₁-C₁₀)-alkyl)-(C₁-C₁₀)-

alkylsulfonamido, (C₇-C₁₆)-aralkylsulfonamido, and N-((C₁-C₁₀)-alkyl-(C₇-C₁₆)-
aralkylsulfonamido; where an aryl radical may be substituted by 1 to 5 substituents
selected from hydroxyl, halogen, cyano, trifluoromethyl, nitro, carboxyl, (C₂-C₁₆)-alkyl,
(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkoxy, (C₃-C₈)-
cycloalkyl-(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkoxy-(C₁-C₁₂)-alkyl, (C₃-C₈)-cycloalkoxy-
(C₁-C₁₂)-alkoxy, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl-(C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl(C₁-
C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-cycloalkoxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₃-C₈)-
cycloalkoxy-(C₁-C₈)-alkoxy-(C₁-C₈)-alkoxy, (C₆-C₁₂)-aryl, (C₇-C₁₆)-aralkyl, (C₂-C₁₆)-
alkenyl, (C₂-C₁₂)-alkynyl, (C₁-C₁₆)-alkoxy, (C₁-C₁₆)-alkenyloxy, (C₁-C₁₂)-alkoxy-(C₁-
C₁₂)-alkyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxy, (C₁-C₁₂)-alkoxy(C₁-C₈)-alkoxy-(C₁-C₈)-
alkyl, (C₆-C₁₂)-aryloxy, (C₇-C₁₆)-aralkyloxy, (C₆-C₁₂)-aryloxy-(C₁-C₆)-alkoxy, (C₇-C₁₆)-
aralkoxy-(C₁-C₆)-alkoxy, (C₁-C₈)-hydroxyalkyl, (C₆-C₁₆)-aryloxy-(C₁-C₈)-alkyl, (C₇-C₁₆)-
aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, (C₇-C₁₂)-
aralkyloxy-(C₁-C₈)-alkoxy-(C₁-C₆)-alkyl, -O-[CH₂]_xC_fH_(2f+1-g)F_g, -OCF₂Cl, -OCF₂-CHFCl,
(C₁-C₁₂)-alkylcarbonyl, (C₃-C₈)-cycloalkylcarbonyl, (C₆-C₁₂)-arylcabonyl, (C₇-C₁₆)-
aralkylcarbonyl, (C₁-C₁₂)-alkoxycarbonyl, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-alkoxycarbonyl,
(C₆-C₁₂)-aryloxycarbonyl, (C₇-C₁₆)-aralkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl,
(C₂-C₁₂)-alkenyloxycarbonyl, (C₂-C₁₂)-alkynyloxycarbonyl, (C₆-C₁₂)-aryloxy-(C₁-C₆)-
alkoxycarbonyl, (C₇-C₁₆)-aralkoxy-(C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkyl-(C₁-C₆)-
alkoxycarbonyl, (C₃-C₈)-cycloalkoxy-(C₁-C₆)-alkoxycarbonyl, (C₁-C₁₂)-
alkylcarbonyloxy, (C₃-C₈)-cycloalkylcarbonyloxy, (C₆-C₁₂)-arylcabonyloxy, (C₇-C₁₆)-
aralkylcarbonyloxy, cinnamoyloxy, (C₂-C₁₂)-alkenylcarbonyloxy, (C₂-C₁₂)-
alkynylcarbonyloxy, (C₁-C₁₂)-alkoxycarbonyloxy, (C₁-C₁₂)-alkoxy-(C₁-C₁₂)-
alkoxycarbonyloxy, (C₆-C₁₂)-aryloxycarbonyloxy, (C₇-C₁₆)-aralkyloxycarbonyloxy, (C₃-
C₈)-cycloalkoxycarbonyloxy, (C₂-C₁₂)-alkenyloxycarbonyloxy, (C₂-C₁₂)-
alkynyloxycarbonyloxy, carbamoyl, N-(C₁-C₁₂)-alkylcarbamoyl, N,N-di(C₁-C₁₂)-
alkylcarbamoyl, N-(C₃-C₈)-cycloalkylcarbamoyl, N,N-dicyclo-(C₃-C₈)-alkylcarbamoyl,
N-(C₁-C₁₀)-alkyl-N-(C₃-C₈)-cycloalkylcarbamoyl, N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-

alkyl)carbamoyl, N-(C₁-C₆)-alkyl-N-((C₃-C₈)-cycloalkyl-(C₁-C₆)-alkyl)carbamoyl, N-(+)-dehydroabietylcarbamoyl, N-(C₁-C₆)-alkyl-N-(+)-dehydroabietylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, N-(C₇-C₁₆)-aralkylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₆)-arylcarbamoyl, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyl, N-((C₁-C₁₆)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₆-C₁₆)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyl, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)-carbamoyl, CON(CH₂)_h, in which a CH₂ group can be replaced by, O, S, N-(C₁-C₈)-alkylimino, N-(C₃-C₈)-cycloalkylimino, N-(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylimino, N-(C₆-C₁₂)-arylimino, N-(C₇-C₁₆)-aralkylimino, N-(C₁-C₄)-alkoxy-(C₁-C₆)-alkylimino, and h is from 3 to 7; carbamoyloxy, N-(C₁-C₁₂)-alkylcarbamoyloxy, N,N-di-(C₁-C₁₂)-alkylcarbamoyloxy, N-(C₃-C₈)-cycloalkylcarbamoyloxy, N-(C₆-C₁₆)-arylcarbamoyloxy, N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₆-C₁₂)-arylcarbamoyloxy, N-(C₁-C₁₀)-alkyl-N-(C₇-C₁₆)-aralkylcarbamoyloxy, N-((C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-((C₁-C₁₀)-alkyl)-N-((C₁-C₁₀)-alkoxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₆-C₁₂)-aryloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, N-(C₁-C₁₀)-alkyl-N-((C₇-C₁₆)-aralkyloxy-(C₁-C₁₀)-alkyl)carbamoyloxy, amino, (C₁-C₁₂)-alkylamino, di-(C₁-C₁₂)-alkylamino, (C₃-C₈)-cycloalkylamino, (C₃-C₁₂)-alkenylamino, (C₃-C₁₂)-alkynylamino, N-(C₆-C₁₂)-arylamino, N-(C₇-C₁₁)-aralkylamino, N-alkyl-aralkylamino, N-alkyl-arylamino, (C₁-C₁₂)-alkoxyamino, (C₁-C₁₂)-alkoxy-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkanoylamino, (C₃-C₈)-cycloalkanoylamino, (C₆-C₁₂)-aroylamino, (C₇-C₁₆)-aralkanoylamino, (C₁-C₁₂)-alkanoyl-N-(C₁-C₁₀)-alkylamino, (C₃-C₈)-cycloalkanoyl-N-(C₁-C₁₀)-alkylamino, (C₆-C₁₂)-aroyl-N-(C₁-C₁₀)-alkylamino, (C₇-C₁₁)-aralkanoyl-N-(C₁-C₁₀)-alkylamino, (C₁-C₁₂)-alkanoylamino-(C₁-C₈)-alkyl, (C₃-C₈)-cycloalkanoylamino-(C₁-C₈)-alkyl, (C₆-C₁₂)-aroylamino-(C₁-C₈)-alkyl, (C₇-C₁₆)-aralkanoylamino-(C₁-C₈)-alkyl, amino-(C₁-C₁₀)-alkyl, N-(C₁-C₁₀)-alkylamino-(C₁-C₁₀)-alkyl, N,N-di-(C₁-C₁₀)-

alkylamino-(C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkylamino-(C₁-C₁₀)-alkyl, (C₁-C₁₂)-alkylmercapto, (C₁-C₁₂)-alkylsulfinyl, (C₁-C₁₂)-alkylsulfonyl, (C₆-C₁₆)-arylmercapto, (C₆-C₁₆)-arylsulfinyl, (C₆-C₁₆)-arylsulfonyl, (C₇-C₁₆)-aralkylmercapto, (C₇-C₁₆)-aralkylsulfinyl, or (C₇-C₁₆)-aralkylsulfonyl;


or wherein R¹ and R², or R² and R³ form a chain [CH₂]_o, which is saturated or unsaturated by a C=C double bond, in which 1 or 2 CH₂ groups are optionally replaced by O, S, SO, SO₂, or NR', and R' is hydrogen, (C₆-C₁₂)-aryl, (C₁-C₈)-alkyl, (C₁-C₈)-alkoxy-(C₁-C₈)-alkyl, (C₇-C₁₂)-aralkoxy-(C₁-C₈)-alkyl, (C₆-C₁₂)-aryloxy-(C₁-C₈)-alkyl, (C₁-C₁₀)-alkanoyl, optionally substituted (C₇-C₁₆)-aralkanoyl, or optionally substituted (C₆-C₁₂)-aroyl; and o is 3, 4 or 5;

or wherein the radicals R¹ and R², or R² and R³, together with the pyridine or pyridazine carrying them, form a 5,6,7,8-tetrahydroisoquinoline ring, a 5,6,7,8-tetrahydroquinoline ring, or a 5,6,7,8-tetrahydrocinnoline ring;

or wherein R¹ and R², or R² and R³ form a carbocyclic or heterocyclic 5- or 6-membered aromatic ring;

or where R¹ and R², or R² and R³, together with the pyridine or pyridazine carrying them, form an optionally substituted heterocyclic ring system selected from thienopyridines, furanopyridines, pyridopyridines, pyrimidinopyridines, imidazopyridines, thiazolopyridines, oxazolopyridines, quinoline, isoquinoline, and cinnoline;

or wherein the radicals R¹ and R², together with the pyridine carrying them, form a compound of Formula Id;



where V is S, O, or NR^k, and R^k is selected from hydrogen, (C₁-C₆)-alkyl, aryl, or benzyl; where an aryl radical may be optionally substituted by 1 to 5 substituents as defined above; and

R²⁴, R²⁵, R²⁶, and R²⁷ in each case independently of each other have the meaning of R¹, R² and R³;

f is 1 to 8;

g is 0 or 1 to (2f+1);

x is 0 to 3; and

h is 3 to 7;

or a physiologically active salt derived therefrom.

~~2. A compound that inhibits hypoxia inducible factor (HIF) hydroxylase activity for use in treating or preventing microcytosis associated with iron deficiency in a subject, wherein the compound is a structural mimetic of 2-oxoglutarate.~~

3. The compound of claim 2 for the use of that claim, wherein

A is C₁-alkylene;

B is -CO₂H;

Q is O;

R⁴ is hydrogen;

X is O;

Y is CR³;

and R¹, R² and R³ are as defined above;

including physiologically active salts derived therefrom.

~~3-4. The compound of claim 1 for the use of that claim, wherein the iron deficiency is functional iron deficiency.~~

~~4-5. The compound of claim 1 for the use of that claim, wherein the iron deficiency is associated with a disorder selected from the group consisting of anemia, iron deficiency anemia, microcytic anemia, inflammation, infection, immunodeficiency disorder and neoplastic disorder.~~

~~5-6A. The compound of claim 1 for the use of that claim, wherein the compound is for use in increasing iron absorption.~~

~~6-7A. The compound of claim 5-6A for the use of that claim, wherein the iron absorption is in the intestine, is in duodenal enterocytes, or is absorption of dietary iron.~~

~~7-8A. The compound of claim 1 for the use of that claim, wherein the compound is for use in increasing iron availability for erythropoiesis or red blood cell production.~~

~~8-9A. The compound of claim 1, 2 or 3 for the use of that claim, wherein the compound is for use in: increasing reticulocytes; increasing hematocrit; increasing hemoglobin; increasing red blood cell count; increasing mean corpuscular hemoglobin; or increasing mean corpuscular volume.~~

~~9-10A. The compound of claim 1, 2 or 3 for the use of that claim, wherein the compound is for use in: increasing serum iron or; increasing total iron binding~~

~~capacity; increasing transferrin saturation; increasing soluble transferrin receptor levels; or decreasing hepcidin expression.~~

11A. The compound of claim 1, 2 or 3 for the use of that claim, wherein the compound is for decreasing hepcidin expression.

~~40. The compound of any preceding claim for the use of that claim, wherein the compound inhibits HIF prolyl hydroxylase activity.~~

~~44~~ 12. The compound of any preceding claim for the use of that claim, wherein the compound is for oral administration.

Conditional amendment to claim 1 (in addition to the unconditional amendments):

1A. A compound that inhibits hypoxia inducible factor (HIF) prolyl hydroxylase activity for use in increasing serum iron in treating functional iron deficiency associated with anemia in a subject, wherein the compound is a structural mimetic of 2-oxoglutarate.

Claims 2-9, 11 and 12 are unchanged and Claim 10 is amended in the following manner:

10A. The compound of claim 1, 2 or 3 for the use of that claim, wherein the compound is for use in: ~~increasing serum iron or~~ increasing transferrin saturation.