



This report is published [online](#). A summary report is being published once a fortnight while influenza activity is low. For further information on the surveillance schemes mentioned in this report, please see information available [online](#).

Indicators for influenza show low levels of activity.



Primary care surveillance

- GP consultation rates for influenza-like illness (ILI) remain below baseline intensity threshold in England, Scotland and Northern Ireland. No data was available for Wales in weeks 30 and 31 (Table 1 & Figure 1).

Table 1: GP ILI consultations for all ages – week 30-31 2020, UK

GP ILI consultation rates (all ages)	Week number		Trend	Peak age group
	30	31		
England (RCGP)	0.4	0.5	↔	<1
Wales	-	-	-	-
Scotland	0.2	0.2	↔	75+
Northern Ireland	0.9	0.2	↔	5-14

Figure 1 : RCGP ILI consultation rates, England



*The Moving Epidemic Method (MEM) has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for GP ILI consultations for the start of influenza activity (based on 10 seasons excluding 2009/10) in a standardised approach across Europe. For MEM intensity threshold values for this season, please visit: <https://www.gov.uk/guidance/sources-of-uk-flu-data-influenza-surveillance-in-the-uk#clinical-surveillance-through-primary-care>

- Syndromic surveillance
 - Syndromic surveillance indicators for influenza remained stable or increased slightly in weeks 30 and 31 2020.
 - For further information, please see the Syndromic surveillance [webpage](#).


Virological surveillance

- Respiratory DataMart system (England)

In week 31 2020, out of the 211 respiratory specimens reported through the Respiratory DataMart System, no sample tested positive for influenza.

RSV positivity remained low at 0.0% in week 31. Rhinovirus positivity remained low but increased slightly at 4.8% in week 31. Adenovirus positivity remained low at 2.2% in week 31. Parainfluenza positivity remained low at 0.0% in week 31. Human metapneumovirus (hMPV) positivity remained low at 0.0% in week 31 2020.

Figure 2: Weekly ICU/HDU influenza admission rate per 100,000 trust catchment population, England, since week 40 2019



Outbreak Reporting

- 229 acute respiratory outbreaks have been reported in week 31 in the UK. 119 outbreaks were from care homes where 76 tested positive for SARS-CoV-2. 7 outbreaks were from educational settings where 2 tested positive for SARS-CoV-2. 41 outbreaks were from workplace settings where 38 tested positive for SARS-CoV-2. 12 outbreaks were from hospitals where 9 tested positive for SARS-CoV-2. 3 outbreak was from a prison where 1 tested positive for SARS-CoV-2. 47 outbreaks were from the Other Settings category where 35 tested positive for SARS-CoV-2.

All-cause mortality surveillance

- In week 31 2020 in England, no statistically significant excess mortality by week of death above the upper 2 z-score threshold was seen overall, by age group or sub nationally (all ages) after correcting GRO disaggregate data for reporting delay with the standardised [EuroMOMO](#) algorithm (Figure 3). This data is provisional due to the time delay in registration; numbers may vary from week to week.

Figure 3: Weekly observed and expected number of all-cause deaths in all ages, with the dominant circulating influenza A subtype, England, 2015 to week 31 2020




Table 2: Excess mortality by UK country, for all ages*

Country	Excess detected in week 31 2020?	Weeks with excess in 2019/20
England	x	50-02;12-23
Wales	x	01;13-20
Northern Ireland	x	49-51; 02-03; 13-22
Country	Excess detected in week 29 2020?	Weeks with excess in 2019/20
Scotland	x	41; 46; 49-51; 01-02; 13-22

* Excess mortality is calculated as the observed minus the expected number of deaths in weeks above threshold

*Note: Delays in receiving all registered deaths from April 2018, following changes in IT systems at ONS, may result in some delays in the model to adjust for most recent deaths.

International Surveillance

- **Influenza** updated on 05 August 2020 (based on data up to 19 July 2020)
 - In the temperate zone of the northern hemisphere, influenza activity remained at inter-seasonal levels. In the temperate zones of the southern hemisphere, the influenza activity remained at inter-seasonal levels. Worldwide, influenza A viruses accounted for the majority of detections.
 - In North America, influenza activity indicators were at very low levels.
 - In Europe, influenza activity remained at inter-seasonal levels.
 - In Central Asia, no influenza updates for this reporting period.
 - In Northern Africa, there were no influenza updates for this reporting period.
 - In Western Asia, there were no influenza detections across reporting countries.
 - In East Asia, influenza illness indicators and influenza activity remained at inter-seasonal levels across all countries.
 - In the Caribbean and Central American countries, sporadic influenza detections were reported in most reporting countries.
 - In tropical South American countries, there were no or sporadic influenza viruses detected across reporting countries.
 - In tropical Africa, there were no or sporadic influenza viruses detected across reporting countries.
 - In Southern Asia, no influenza detections were reported.
 - In South East Asia, no influenza detections were reported.
 - The WHO GISRS laboratories tested more than 302,586 specimens between 06 July 2020 and 19 July 2020. 45 were positive for influenza viruses, of which 26 (57.8%) were typed as influenza A and 19 (42.2%) as influenza B. Of the sub-typed influenza A viruses, 7 (100%) was influenza A(H1N1)pdm09. Of the characterised B viruses influenza B viruses 2 (100%) belonged to the B-Victoria lineage.
- **MERS-CoV** latest update on 05 August 2020
 - Up to 04 August 2020, a total of five cases of Middle East respiratory syndrome coronavirus, MERS-CoV, (three imported and two linked cases) have been confirmed in the UK. On-going surveillance has identified 1,816 suspected cases in the UK since September 2012 that have been investigated for MERS-CoV and tested negative.
 - From 1 April to 31 May 2020, the National IHR Focal Point of Saudi Arabia reported 9 new cases of MERS-CoV infection, including five deaths.
 - Globally, since September 2012, WHO has been notified of 2,562 laboratory-confirmed cases of infection with MERS-CoV, including 881 associated deaths. Further guidance on the management of possible cases in the UK is available [online](#). The latest ECDC MERS-CoV risk assessment can be found [here](#), where it is highlighted that risk of widespread transmission of MERS-CoV remains very low.
- **Avian/Zoonotic influenza** latest update on 22 July 2020
 - Between **9 May to 10 July 2020**, two new human infections with avian influenza A(H9N2) viruses were reported from China, one human infection with an influenza A(H1N1) variant virus was reported from Germany and one human infection with an influenza A(H1N2) variant virus was reported from Brazil.
 - For further updates please see the [WHO website](#) and for advice on clinical management in the UK please see information available [online](#).
- **Coronavirus Disease 2019 (COVID-19)** latest update 05 August 2020
 - Up to **04 August 2020**, a 306,293 total of cases of COVID-19 have been confirmed in the UK.
 - Globally, up to 04 August 2020, [WHO](#) has been notified of 18,142,718 confirmed cases of COVID-19 infection, including 691,013 related deaths.