

GP OOHSS

GP Out-of-Hours Surveillance System: England

Year: 2020 Week: 34 25 August 2020 Key messages In This Issue: Data to: 23 August 2020 During week 34, GP out of hours contacts for heatstroke and insect bites Key Messages. decreased, returning to baseline levels (figures 11 & 12). Weekly summary. Total contacts. Contacts for acute respiratory infection increased slightly, though remain below seasonally expected levels (figures 2 & 2a). Syndromic indicators. Notes and caveats. A Heat-Health Watch system operates in England from 1 June to 15 September each year. As part Further information. of the Heatwave Plan for England, the PHE Real-time Syndromic Surveillance team will be Acknowledgements. routinely monitoring the public health impact of hot weather using syndromic surveillance data during this period. Heat-health watch level (current reporting week): Level 1: Summer preparedness http://www.metoffice.gov.uk/weather/uk/heathealth % % No. of Syndromic indicators **Key indicator** contacts Week 34 Week 33 Trend* at a glance: All OOH contacts, all causes 164,110 Acute respiratory infection 3,469 5.74 5.52 $\mathbf{\Lambda}$ Number of contacts and Influenza-like illness 62 0.10 0.11 ←→ percentage of Read Bronchitis/bronchiolitis 21 0.03 0.03 ←→ coded contacts. Difficulty breathing/wheeze/asthma 855 1.41 1.42 $\leftarrow \rightarrow$ 24 0.04 Pharyngitis 0.02 Gastroenteritis 1,882 3.11 2.96 ←→ Diarrhoea 628 1.04 0.92 ←→ Vomiting 725 1.20 1.18 **4** -> Chest pain/myocardial infarction 1.06 783 1.29 $\leftarrow \rightarrow$ Heatstroke 0.00 1 0.07 Insect bites 1.236 2.04 2.76 *Trend: reports on the trend seen over previous weeks in the percentage of Read coded contacts. 60.000 contacts: 50,000 40.000 Number of contacts 30,000 20.000 10.000 25/08/19 22/09/19 20/10/19 17/11/19 15/12/19 12/01/20 09/02/20 08/03/20 05/04/20 03/05/20 31/05/20 28/06/20 26/07/20 23/08/20 Bank Holiday weekend All contacts 7 day moving average

1: Total out-of-hours


Daily total number of out-of-hours and unscheduled contacts and 7-day moving average (adjusted for bank holidays).


2: Acute Respiratory Infection daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7-day moving average*.

2a: Acute Respiratory Infection by age group.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.





Intentionally left blank


*7-day moving average adjusted for bank holidays.

GP OOHSS


Year: 2020 Week: 34


3: Influenza-like illness daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.



As a percentage of total contacts within each age group, shown as a 7-day moving average*.





Intentionally left blank

*7-day moving average adjusted for bank holidays.

GP OOHSS

WW Public Health England

25 August 2020

GP OOHSS

Year: 2020 Week: 34

4: Bronchitis/ bronchiolitis daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.


4a: Bronchitis/ bronchiolitis daily contacts by age group*.


As a percentage of total contacts within each age group, shown as a 7-day moving average*.


5: Difficulty breathing/ wheeze/asthma daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.

*7-day moving average adjusted for bank holidays.








GP OOHSS


Year: 2020 Week: 34



As a percentage of total contacts within each age group, shown as a 7-day moving average *.


6: Acute pharyngitis and persistent sore throat.

Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.



Shown as a percentage of the total contacts with a Read code and as a 7day moving average*.

*7-day moving average adjusted for bank holidays.



8: Diarrhoea daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.


8a: Diarrhoea daily contacts by age group*.

As a percentage of total contacts within each age group, shown as a 7-day moving average*.

9: Vomiting daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.

*7-day moving average adjusted for bank holidays.



GP OOHSS


Year: 2020 Week: 34

10: Chest pain/ myocardial infarction daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.


11: Heatstroke daily contacts.

Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.



Shown as a percentage of the total contacts with a Read code and as a 7 -day moving average*.

*7-day moving average adjusted for bank holidays.



GP OOHSS

Year: 2020 Week: 34

GP OOHSS

25 August 2020	Year: 2020 Week: 34
Notes and caveats:	 This bulletin presents data from the Public Health England (PHE) GP Out -of-hours\Unscheduled Care Surveillance System (GP OOHSS). Fully anonymised data from GP out-of-hours (OOH) and unscheduled care service providers in England are being transferred to the PHE for analysis and interpretation by the PHE Real-time Syndromic Surveillance Team (ReSST). This system supplements existing PHE syndromic surveillance systems by monitoring data on general practitioner consultations outside of routine surgery opening times (evenings, weekends and bank holidays) and unplanned contacts within NHS primary care. The key indicators presented within this bulletin are derived by grouping selected Read coded consultations. GP OOH consultation data are analysed on a daily basis to identify national and regional trends. A statistical algorithm underpins each system, routinely identifying activity that has increased significantly or is statistically significantly high for the time of year. Results from these daily analyses are assessed by the ReSST, along with analysis by age group, and anything deemed of public health importance is alerted by the team. Baselines represent seasonally expected levels of activity and are constructed from historical data since Nov 2009. They take into account any known substantial changes in data collection, population coverage or reporting practices. Gastroenteritis, diarrhoea and vomiting baselines also account for changes since the introduction of rotavirus vaccine in July 2013. Baselines are refreshed using the latest data on a regular basis.
Moving Epidemic Method (MEM):	 During winter we present Moving Epidemic Method (MEM) influenza thresholds on selected indicators. The moving epidemic method or MEM is a standard methodology used for setting influenza thresholds across many European nations.¹ MEM is used for GP OOH ILI thresholds at a national level. MEM thresholds should be interpreted using 7 day moving averages rather than daily data. MEM thresholds currently use six years of historic data (2013-2019). The thresholds are re-calculated every year. Baseline ('Pre-epidemic') thresholds are used alongside other surveillance systems to identify the start of influenza circulating in the community 40%, 95% and 97.5% intensity thresholds are used to identify when influenza activity moves from low to medium, high or very high. ¹Vega T et al. Influenza Other Respir Viruses. 2013;7(4):546-58.
Further information:	The GP Out-of-Hours Surveillance System Bulletin can also be downloaded from the PHE Real-time Syndromic Surveillance website which also contains more information about syndromic surveillance:
	https://www.gov.uk/government/collections/syndromic-surveillance-systems-and-analyses
Acknowledgements:	We are grateful to Advanced and the GP OOH and unscheduled care service providers who have kindly agreed to participate in this system.
	PHE Out-of-Hours/Unscheduled Care Surveillance
Contact ReSST: syndromic.surveillance	Produced by: PHE Real-time Syndromic Surveillance Team 1≝ Floor, 5 St Philips Place, Birmingham, B3 2PW Tel: 0344 225 3560 > Option 4 > Option 2 Fax: 0121 236 2215